

**New York**  
**State College of Agriculture**  
**At Cornell University**  
**Ithaca, N. Y.**

---

**Library**

Cornell University Library  
**QK 306.B113**

**Manual of British botany, containing the**



**3 1924 003 269 010**

mann



Cornell University  
Library

The original of this book is in  
the Cornell University Library.

There are no known copyright restrictions in  
the United States on the use of the text.

MANUAL  
OF  
BRITISH BOTANY,

CONTAINING THE  
FLOWERING PLANTS AND FERNS.

ARRANGED ACCORDING TO  
THE NATURAL ORDERS.

BY  
CHARLES CARDALE BABINGTON, M.A.,  
F.R.S., F.L.S., ETC. ETC.,  
PROFESSOR OF BOTANY IN THE UNIVERSITY OF CAMBRIDGE.

EIGHTH EDITION.  
CORRECTED THROUGHOUT.



LONDON:  
JOHN VAN VOORST, PATERNOSTER ROW.

MDCCCLXXXI.

“ Quod ad me attinet, ingenue fatear, me in rebus dubiis de specifica differentia numquam consulere Auctores, qui in herbariis plantis multis, sed eos modo, qui in natura plantis multum student. At iis, qui ad præceptas opiniones experientiam suam concinnant et in singulo externæ faciei lusu, neglectis notis essentialibus, formas transitorias vident, parum fido.”—FRIES.

## PREFACE.

---

IN this work it has been the Author's wish to adopt in all cases those names which have the claim of priority, unless good cause should be shown for a contrary proceeding; and with this object he has carefully examined nearly all the best European Floras, comparing our plants with the descriptions contained in them, and in most cases with foreign specimens of undoubted authenticity. In the adoption of genera and species an endeavour has been made, by the examination of the plants themselves, to determine which are to be regarded as truly distinct,—thus, it is hoped, taking Nature as a guide. Still, let it not be supposed that any claim is made to peculiar accuracy, or that the Author considers himself qualified to dictate to any student of botany; for he is well aware that there are many points upon which persons who have carefully studied the subject form different conclusions from those to which he has been led.

The progress of our knowledge has caused changes in the nomenclature in successive editions of this book and in the Author's views of the value of forms—as species or varieties. The inconvenience of these alterations to all, especially to statistical botanists, is fully admitted; but the Author does not know of any mode by which it can be avoided if each edition is to be brought up as completely as is in his power to the contemporary knowledge of our plants. No alterations have been admitted until careful

study has convinced the Author that they are required. He may have fallen into error, but has earnestly endeavoured to discover the truth.

Attempts have been made greatly to reduce the number of recognized species found in Britain ; but the results obtained seem to be, so totally opposed to the teaching of the plants themselves, and the evidence adduced in their favour is so seldom more than a statement of opinion, that they cannot safely be adopted ; nor does the plan of the present work admit of a discussion of the many questions raised by them. Also it has been laid down as a rule by some botanists, that no plant can be a species whose distinctive characters are not as manifest in an herbarium as when it is alive. We are told that our business as descriptive botanists is not "to determine what is a species," but simply to describe plants so that they may be easily recognized from the dry specimen. The Author cannot agree to this rule. Although he, in common with other naturalists, is unable to define what is a species, he believes that species exist, and that they may often be easily distinguished amongst living plants, although sometimes separated with difficulty when dried specimens alone are examined. He thinks that it is our duty as botanists to study the living plants whenever it is possible to do so, and to describe from them ; to write for the use and instruction of field- rather than cabinet-naturalists ; for the advancement of a knowledge of the plants rather than for the convenience of possessors of herbaria : also that the differences which we are able to describe as distinguishing plants being taken from their more minute organs, does not invalidate their claim to distinction. It seems to be our business to decide upon the probable distinctness of plants before we attempt to define them—to make the species afford the character, not the character define the species.



This volume being intended as a field-book or travelling companion for botanists, it is advisable to restrict the space allotted to each species as much as possible; and accordingly the characters and observations are only such as appear to be necessary for an accurate discrimination of the plants. Facts relating to their geographical distribution are therefore usually omitted. Synonyms have been almost wholly omitted; but the plates of the original *English Botany* or some other British plates are quoted. *Syme's English Botany* may well be used by those who desire full descriptions of the plants; and the plates in that work have often had valuable additions made to those of the old *English Botany*, from which most of them are taken. Localities are only given for new or rare plants; Mr. Watson's works and the numerous local floras render it unnecessary inconveniently to swell the present volume by their introduction. But in order to convey some idea of the distribution of plants throughout the United Kingdom, the letters E., S., or I. have been appended to the descriptions of such species as have, it is believed, been found in England, Scotland, or Ireland. The descriptions of a considerable number of plants which only occur in the Channel Islands, and are, therefore, not properly parts of the British Flora; or which, although included in our lists, there is reason to suppose have never been really detected in Britain; or, although naturalized, have very slender claims to be considered aboriginal natives; or which are now supposed to be lost by the alterations made in the places where they were found by our predecessors; are included within []; and notices of a few plants concerning which more accurate information is requisite, are distinguished in a similar manner. It is hoped that by this arrangement the truly indigenous species will be clearly distinguished from those which have little or no claim to be considered aboriginal

or even thoroughly naturalized. The attempt to do this is necessary for two seemingly contradictory reasons, namely :—the great tendency of many collectors to consider native any plant found growing upon a spot where it is not cultivated ; and the peculiar scepticism of some of our botanists concerning the claims of many local or thinly scattered species to be admitted as indigenous, even when their distribution upon the European continent is not unfavourable to the belief that they may inhabit Britain. It has been recommended that the descriptions of these excluded species should be placed in an Appendix or even omitted ; but as some of them are not unlikely to be observed by collectors, it is more convenient that they should be arranged with their allies. Those who desire to obtain a complete knowledge of the distribution of our plants should consult Watson's *Cybele Britannica*, and Moore & More's *Cybele Hibernica*.

Full characters of the Natural Orders are to be found in most of the best 'Introductions to Botany;' and it has therefore not been considered advisable to give them in detail in the present volume. In his definitions, the Author has endeavoured to point out the characteristic marks, more especially as far as British plants are concerned.

In using this book the student will find it convenient to pay attention to the *italicized* parts of the generic and specific characters, and, if they are found to agree with the plant under examination, then to compare it with the other parts of those characters, and also with those of allied genera and species.

It is most desirable that the students of our native flora should not confine their attention to books published in this country. Owing to such an unavoidable restriction we fell far behind our continental brethren during the earlier part of the present century. A few modern works may be

named which will assist them in their studies. Koch's *Synopsis Floræ Germanicæ*, ed. 2, and Grenier and Godron's *Flore de France* are strongly recommended,—also, although in a rather less degree, Lloyd's *Flore de l'Ouest de la France*, Brébisson's *Flore de la Normandie*, and Cosson and Germain's *Flore des environs de Paris*, ed. 2. Boreau's *Flore du Centre de la France*, ed. 3, and the scattered papers of M. Alexis Jordan of Lyons are valuable for the study of varieties; for many of their species can claim no higher rank. But, above all, the works of Fries deserve careful study—especially his *Novitiæ Floræ Suevicæ*, with its three *Mantissæ*, and *Summa Vegetabilium Scandinaviæ*. It is necessary to warn students against the very common error of supposing that they have found one of the plants described in a foreign Flora when in reality they have only gathered a variety of some well-known British plant. The risk of falling into such errors renders it necessary to consult such works as those of Messrs. Boreau and Jordan with great caution, lest we should be misled by descriptions, most accurate, indeed, but often rather those of individuals than species. Amongst plants so closely allied as are many of those called species in some continental works, it is scarcely possible to arrive at a certain conclusion without the inspection of authentic specimens.

The Author takes this opportunity of returning most sincere thanks to his botanical friends and correspondents (far too numerous to record by name) for the great assistance they have again rendered to him by the communication of valuable suggestions, observations, and specimens.

The book has been again carefully revised throughout, so as, if possible, to keep pace with the rapidly advancing knowledge of British plants.

As many as possible of the real English names are given. All the genera and species could not be thus named, owing

to the absence of any recognized English terms which have been applied to them. It does not seem desirable to invent or adopt new English names, known only to botanists, for the few genera which have them not, the Latin name being sufficient in those cases, and better in the original than in an Anglicized form.

As the portability of this volume is perhaps its most valuable quality, the Author cannot accede to the wishes of some young botanists by prefixing a short *Introduction to Botany*. Sufficient space is wanting to admit of its possessing that fulness of detail without which such an *Introduction* would be worse than useless. Oliver's *Lessons in Elementary Botany* will be found of much use; but students should afterwards make themselves acquainted with the contents of some more elaborate treatise, such as Balfour's *Manual*, Masters's edition of Henfrey's *Elementary Course*, or Gray's *Structural Botany*.

It is hoped that those who use this book will favour the author with information of any (even the slightest) addition, correction or alteration that may appear to be necessary, in order that it may be employed in the preparation of a future edition, as it is only through such assistance that the flora of an extensive country can attain to even a moderate degree of perfection.

Cambridge, July 15, 1881.

# A GLOSSARY

OF

## THE TERMS USED IN THE MANUAL.

---

- Accumbent*; used to express the application of the edges of the cotyledons to the radicle in the seeds of Crucifers.
- Achene*; a hard dry one-seeded superior pericarp.
- Acotyledonous*; without distinct cotyledons.
- Acuminate*; drawn out into a long point, but with the sides slightly hollowed.
- Acute*; sharp; forming an angle less than a right angle at the tip.
- Adhering*; uniting together of two different parts, as a calyx to an ovary.
- Adnate*; attached throughout their whole length. Adnate anthers have their lobes so attached to the filament. Stipules are often adnate to the petiole by one of their edges.
- Adpressed*; pressed close to any thing.
- Adpressed-serrate*; serrate with the teeth lying closely on each other or to the edge of the leaf.
- Æstival*; produced in summer.
- Albumen*; nutritious matter contained in the seed to feed the young plant: more correctly called *perisperm*.
- Alternate*; placed successively on the opposite sides of an axis, as in the case of leaves; or opposite to the spaces between the parts of the next whorl in flowers.
- Amplexicaul*; clasping the stem with their base.
- Anastomosing*; veins combining with each other at their ends.
- Annual plants* rise from the seed, flower, and die in the same year.
- Annular*; forming a ring.
- Anterior*; the part of a flower next the bract or in front.
- Anther*; the part of the stamen which contains the pollen.
- Apex*; the end furthest from the point of attachment.
- Apical*; at or relating to the apex.
- Apiculate*; having a very small hard point at the end, usually formed by the tip of the midvein.
- Apocarpous*; fruit formed of carpels which are quite separate.
- Approximate*; close together.
- Arching*; curved into the form of an arch.

- Arcuate* ; curved so as to form a considerable part of a circle.  
*Aril* ; an aftergrowth from the placenta or seedstalk surrounding the seed.  
*Arillode* ; an aftergrowth from the lips of the foramen (or terminal opening of seed).  
*Ascending* ; curving upwards into a vertical position.  
*Asperous* ; rough with short raised points.  
*Attenuate* ; narrowing gradually to a point.  
*Auricled* ; having *auricles*, or appendages at the base of the leaves.  
*Awn* ; a long-pointed bristlelike appendage, as the beard of Barley.  
*Awned* ; having awns.  
*Axil* ; the upper angle formed by the union of the stem and leaf.  
*Axillary* ; placed in an axil.  
*Axis* ; the line passing through the centre of any thing ; the common stalk of the flowers in a spikelet of Grasses.
- Baccate* ; pulpy like a berry.  
*Base* ; the end nearest to the point of attachment.  
*Beak* ; a long pointed projection.  
*Bearded* ; having long hair like a beard.  
*Berry* ; a pulpy fruit containing several seeds ; a true *bacca* when inferior, a *uva* when superior.  
*Biennial* plants spring from the seed in one year, flower in the following year, and then die.  
*Bifariouly* ; arranged in two rows, one on each side of any thing.  
*Bifid* ; divided halfway down into two parts.  
*Bipartite* ; divided nearly to its base into two parts.  
*Bipinnate* ; when the divisions of a pinnate leaf are themselves pinnate.  
*Bipinnatifid* ; when the divisions of a pinnatifid leaf are themselves pinnatifid.  
*Biternate* ; when the divisions of a ternate leaf are themselves ternate.  
*Boatshaped* ; resembling a small boat.  
*Bracteoles* ; minute bracts.  
*Bracts* ; small leaves somewhat different from the others, seated on the inflorescence.  
*Bulb* ; a leaf-bud with fleshy scales, usually placed underground.  
*Bulbiferous* ; bearing bulbs on its stem.  
*Bulblike* ; resembling a bulb in appearance, but solid.  
*Bulbous* ; having radical bulbs.  
*Bulbous* hairs have a round swelling at their base.
- Cæspitose* ; growing in tufts from the root.

- Calyx*; the outer whorl of leaflike organs forming the flower, usually green, called sepals.
- Capillary*; like very slender threads.
- Capitate*; growing in heads or close clusters; having a knob like the head of a pin.
- Capsular*; like a capsule.
- Capsule*; a dry usually many-seeded seed-vessel.
- Carpel*; the divisions of the ovary or capsule: sometimes one carpel forms an ovary, being rolled up so that its edges meet.
- Carpophore*; the stalk of the ovary or capsule within the outer whorls of the flower.
- Catkin*; a deciduous unisexual spike of crowded flowers in which the perianths are replaced by bracts.
- Cauline*; growing from the stem, not radical.
- Cellular tissue*; a collection of minute vesicles filled with fluid.
- Chaffy*; covered with minute membranous scales.
- Channelled*; hollowed somewhat like a gutter.
- Cilia*; hairs placed like eyelashes on the edge of any thing.
- Ciliate*; with cilia.
- Circinate*; rolled up from the top towards the base like a crosier.
- Clavate*; clubshaped.
- Claw*; the narrow base of a petal.
- Clawed*; having a claw.
- Cleft*; deeply cut, but not to the midrib.
- Clubshaped*; a long solid body which is slender at the base and gradually thickens upwards.
- Cluster*; a kind of dense cyme; also the patches of capsules in Ferns.
- Cæsious*; with a fine pale-blue bloom.
- Cohering*; the attachment to each other of similar parts; as the petals forming a gamopetalous corolla.
- Collapsing*; shrinking together. The submersed and much-divided leaves of aquatic plants often collapse into a form like a painter's pencil, when removed from the water.
- Columella*; a cylindrical central placenta.
- Commissure*; the inner faces of the carpels (mericarps) of Umbelliferæ, by which they join.
- Compound*; formed of many similar parts which ultimately and naturally separate from each other. *A compound umbel* has small umbels on its branches.
- Compressed*; when flattened laterally.
- Conduplicate*; folded upon each other lengthwise.
- Cone*; fruit of a fir-tree.
- Conical*; a solid figure narrowing to a point from a circular base.
- Connate*; when two similar parts, as leaves, are slightly connected round the stem.

*Connective* ; the continuation of the filament between the cells of an anther.

*Connivent* ; converging.

*Constricted* ; narrowed at some point as if by the pressure of a string.

*Contiguous* petals touch or overlap by their edges.

*Converging* ; their points gradually approaching.

*Convolute* ; rolled together lengthwise.

*Cordate* ; ovate, acute, with two rounded lobes at the base ; like the figure of the heart on cards : a *cordate-based* leaf is of any shape, but has the two lobes at its base.

*Coriaceous* ; leathery ; firm, dry, tough.

*Corm* ; a fleshy bulblike, but solid, not scaly underground stem.

*Corneous* ; like horn.

*Corolla* ; the whorl of floral leaves between the calyx and stamens, usually coloured, called petals.

*Corymb* ; a raceme with the peduncles becoming gradually shorter as they approach the top, so that all the flowers are about on a level.

*Corymbose* ; in the form of a corymb.

*Cotyledons* ; the seed-lobes, often forming the first leaves of the plant.

*Crenate* ; with rounded marginal teeth. When these are again crenate, the whole is *doubly crenate* : not *bicrenate*, which means having two such teeth.

*Crenatures* ; the blunt rounded teeth of a crenate leaf.

*Crenulate* ; minutely crenate.

*Crested* ; having an appendage like a crest.

*Crowned* ; having an appendage on the upper side at the base of the limb, as some petals.

*Cruciform* ; four parts, as petals, arranged so as to form a cross.

*Crustaceous* ; hard, thin and brittle.

*Cuneate* ; like a wedge, but attached by its point.

*Cuspidate* ; abrupt, but with a point starting suddenly from the middle of its end.

*Cuticle* ; the external skin.

*Cylindrical* ; nearly in the form of a cylinder.

*Cyme* ; inflorescence formed of a terminal flower, beneath which are lateral branches each having a terminal flower and lateral branches again similarly dividing, and so on. A *globose* cyme has flowers so placed as to form a globose mass. A *scorpioid* cyme produces only the external branch of each pair, except the first.

*Cymose* ; arranged in a cyme.

*Deciduous* ; falling off.

*Declining* ; straight, but pointing downwards.



- Decumbent*; lying on the ground, but tending to rise at the end.
- Decurrent*; when the limb of a leaf is prolonged down the stem below the point of attachment of the midrib.
- Decussate*; opposite leaves, but the successive pairs placed at right angles to each other.
- Deflexed*; curved downwards or towards the back.
- Dehiscence*; the mode in which an organ opens.
- Deltoid*; fleshy with a triangular transverse section.
- Dentate*; with short equilateral triangular teeth. When these are again dentate, the whole is *doubly dentate*: not *bidentate*, which means having two teeth.
- Denticulate*; finely dentate.
- Depressed*; when flattened vertically or at the top.
- Determinate* inflorescence ends in a flower.
- Dicotyledonous*; with two opposite cotyledons.
- Didymous*; formed of two similar parts attached to each other by a small portion of their margin.
- Diffuse*; widely spreading.
- Digitate*; fingered; of several leaves all starting from the top of the petioles.
- Dioecious*; with the sexes on different plants.
- Disk*; a fleshy space from which the stamens and pistils spring, or between them; the central part of a head (capitulum).
- Dissepiments*; vertical plates dividing an ovary into parts; septa.
- Distichous*; arranged above each other in two rows on opposite sides of an axis.
- Distinct*; separate from its neighbours.
- Divaricate*; spreading at an obtuse angle.
- Diverging*; gradually separating.
- Dorsal*; attached to, or on the back.
- Drupe*; a one-celled superior fruit, not bursting, fleshy externally, stony within, containing one or two seeds.
- Echinate*; armed with straight slender prickles like a hedgehog.
- Elliptic*; oval but acute at each end.
- Elongate*; much lengthened.
- Emarginate*; slightly notched at the end.
- Embryo*; the young plant as first seen in the seed.
- Entire*; not toothed nor lobed at the edge.
- Epidermis*; the skin.
- Epigynous*; apparently seated upon the ovary.
- Epipetalous*; borne on the petals.
- Epiphytes*; plants growing upon others, but not deriving nourishment from their juices.
- Equalling*; when the ends of organs rise to the same height even though their relative lengths are different.

*Equitant*; when a conduplicate organ covers the edges of another similarly folded, and that covers a third, and so on.

*Erect*; standing nearly perpendicular to that from which it grows, as a seed rising from the base of an ovary; at right angles to its support.

*Exceeding*; when an organ extends beyond an adjoining organ, but is not necessarily itself longer than it.

*Excurent*; extending beyond the edge or point.

*Exserted*; projecting beyond that which surrounds its base.

*Extrorse* anthers have the slit by which the pollen escapes directed from the ovary.

*Falcate*; like a sickle.

*Falling short of*; the reverse of exceeding.

*Fasciculate*; when several similar parts are collected into a bundle and spring from the same spot; often the developed leaves of an undeveloped axillary branch form a fascicle.

*Fastigate*; when all the branches are parallel and point upwards.

*Feathery*; like a feather in structure.

*Felted*; tomentose.

*Fibre*; a hair-like kind of elementary structure.

*Fibrous*; having many threadlike parts.

*Filament*; the stalk usually found supporting an anther.

*Filiform*; like a thread.

*Flaccid*; weak.

*Flexuose*; zigzag, usually changing its direction at each joining.

*Floccose*; with little tufts like wool.

*Follicle*; an inflated 1-celled carpel, opening by only one suture to which several seeds are attached.

*Forked*; like a fork of two prongs.

*Frond*; the leaflike part of Ferns.

*Fruit*; the seed-vessel with its ripe contents and any external appendages.

*Fruit-bearing*; the state of the inflorescence when the fruit is ripe or nearly so, contradistinguished from *flower-bearing*.

*Fruticose*; shrubby.

*Fugacious*; soon falling off.

*Funnel-shaped*; tubular below, but gradually enlarging upwards.

*Furcate*; forked.

*Fusiform*; spindle-shaped; thick, tapering to each end.

*Gamosepalous*; *gamopetalous*; when the sepals or petals are joined by their edges so as apparently to form one.

*Germen*; the ovary.

*Gibbous*; swollen on one side.

*Glabrous*; without hairs or other clothing.

*Gland*; a wartlike cellular secreting organ usually raised above the surface.

*Glandular*; having glands.

*Glandular-hairy*; having hairs tipped with glands.

*Glandular-serrate*; having short teeth tipped with glands.

*Glaucous*; green with a whitish-blue lustre.

*Globose*; round like a globe.

*Glumes*; the scales enclosing the spikelet of flowers in Grasses; the imbricate bracts enclosing the flowers of Sedges.

*Glumiferous*; having flowers covered by glumes.

*Granular*; covered with minute projecting points.

*Habit*; the general appearance of a plant.

*Haft*; a winged leafstalk; the linear part of a spatulate leaf or petal.

*Hastate*; enlarged at the base into two lobes directed nearly horizontally.

*Head*; a close terminal collection of flowers surrounded by an involucre.

*Helmet*; the hooded upper part of a flower.

*Helmet-shaped*; arched and concave like a helmet.

*Herbaceous*; the parts of plants which are not woody; also organs, or parts of them, of a green colour.

*Hermaphrodite*; having both sexes in one flower.

*Hilum* or *hile*; the mark on a seed which indicates its place of attachment.

*Hispid*; covered with stiff hairs.

*Hoary*; with greyish-white down.

*Hooded*; formed into a hood at the end.

*Horizontal*; spreading at right angles to their support, as leaves on a stem.

*Hybrid*; a mule.

*Hypogynous*; springing from below the base of the ovary and not attached to the calyx.

*Imbricate*; arranged over each other like the tiles of a roof.

*Imparipinnate*; pinnate with a single terminal leaflet.

*Incise*; deeply cut.

*Included*; not extending beyond the organs surrounding it.

*Incumbent*; when the radicle is applied to the back, not edges, of the cotyledons.

*Incurved*; curved inwards.

*Indefinite*; many but uncertain in number.

*Indehiscent*; not bursting.

- Indeterminate*; inflorescence having always a terminal leaf-bud.
- Induplicate*; when the edges of organs arranged in a valvate manner are folded inwards.
- Indusium*; a thin membrane often covering the clusters of capsules of Ferns.
- Inferior*; an inferior calyx or corolla is wholly free from the ovary; the reverse of superior.
- Inflexed*; curved inwards.
- Inflorescence*; arrangement of the flowers.
- Innate*; attached by their base to the apex of a stalk as are some anthers.
- Inserted*; growing upon.
- Internode*; the space between two nodes; a joint.
- Interruptedly pinnate*; when pairs of small pinnæ (leaflets) alternate with large pinnæ.
- Introrse*; anthers having the slit by which the pollen escapes directed towards the ovary.
- Inverse*; *inverted*. An embryo is so called when its radicle is directed towards a point at the opposite end of the seed from the hile.
- Involucels*; the involucre of secondary umbels.
- Involucre*; the whorled bracts at the base of an umbel or head; or sometimes below a single flower.
- Involute*; rolled from the back of any thing, as towards the upper side of a leaf.
- Joinings*; the places where the parts of the stem are attached to each other; the nodes.
- Joints*; the spaces between the knots, nodes, or joinings; the parts joined.
- Keel*; a prominent ridge. The two lower petals of a Pea-flower, within the others and united more or less by their anterior edge, form the keel.
- Kneed*; bent like the knee.
- Knots*; the joinings or nodes of the stem in Grasses.
- Label*; the terminal segment of the lip in Orchids.
- Lacinate*; divided into narrow irregular lobes.
- Lanceolate*; narrowly elliptic and tapering to each end.
- Lancet-shaped*; shortly and bluntly lanceolate.
- Lax*; loosely arranged.
- Leaflets*; the subdivisions of compound leaves.
- Legume*; a one-celled and two-valved seed-vessel with the seeds arranged along the inner angle, as the pod of a Pea.
- Lenticular*; like a doubly convex lens.

*Ligulate*; strap-shaped; not very narrow nor long, and with nearly parallel sides.

*Ligule*; a membrane at the base of the limb of the leaf of Grasses.

*Limb*; the flattened expanded part of a leaf or petal.

*Linear*; very narrow and long, with parallel sides until near the end.

*Lingulate*; tongue-shaped; long, fleshy, convex, blunt.

*Lipped*; applied to a corolla or calyx appearing to consist of two lips.

*Lobate*; lobed; with large divisions.

*Loculicidal*; opening down the back (or midrib) of the carpel.

*Lower part* of a floral whorl; that furthest from the main axis; anterior.

*Lunate*; shaped like the new moon.

*Lyrate*; a pinnatifid leaf with the lobes successively and gradually enlarging from the petiole, and ending in one still larger lobe.

*Marcrescent*; fading but remaining in its place.

*Medullary*; relating to the pith. Medullary rays are plates of cells which connect the pith with the growing part next to the bark.

*Membranous*; of the texture of membrane; thin and flexible.

*Mericarps*; the carpels of Umbelliferæ.

*Midrib*; the large vein extending along the middle of a leaf from its petiole nearly or quite to the other end.

*Moniliform*; cylindrical but constricted at regular intervals.

*Monocotyledonous*; having one sheathing cotyledon.

*Monœcious*; with the sexes in separate flowers on the same plant.

*Monosepalous*; *monopetalous*; when the sepals or petals are joined by their edges so as apparently to form one.

*Mucronate*; abruptly tipped with a short point of the same texture.

*Multifid*; divided into many parts.

*Muricate*; covered with short sharp points.

*Mono-, di-, &c. androus*; with 1, 2, &c. stamens.

*Mono-, di-, &c. gynous*; with 1, 2, &c. free styles or stigmas.

*Nectary*; an organ which secretes honey.

*Netted*; covered with lines connected together like network.

*Node*; a point in a stem where a leaf is produced; a joining.

*Nut*; a hard dry 1-seeded superior pericarp; also used for a *glans*, a hard dry 1- or few-seeded inferior pericarp not bursting and seated in a cup-like involucre, *e. g.* acorn.

*Ob*; in conjunction with terms means inverted; as *obovate* is ovate with the attachment at the narrow end.

*Oblong*; long oval, equally broad at each end.

*Ocrea*; a tubular membranous stipule surrounding the stem.

*Opaque*; not shining.

*Opposite*; when two similar organs grow one on each side of some body; or different organs are opposed to each other with a stem between them.

*Orbicular*; nearly round and flat.

*Oval*; an ellipse; a figure rounded at each end, not broader at one end than at the other; and about twice as long as broad.

*Ovary*; the young seed-vessel.

*Ovate*; eggshaped; a short flat figure (thin like a leaf) rather broader below the middle of its length.

*Ovoid*; a solid eggshaped figure.

*Ovule*; the young seed.

*Palate*; the prominent part of the base of the lower lip which closes the mouth of a ringent corolla.

*Pales*; the leaflike parts of the flower of Grasses, enclosing the stamens, pistils, and hypogynous scales.

*Palmate*; with lobes spreading like the fingers of a hand from the same point.

*Panicle*; a raceme with branching pedicels; hence *paniculate*.

*Papilionaceous*; like the flower of a Pea.

*Papillæ*; small elongated protuberances.

*Papillose*; with small long protuberances.

*Pappus*; the crest of the fruit in Composites, formed of the altered limb of the calyx.

*Parabolic*; starting from a broad base and gradually narrowing with curved sides to a blunt point, as the divisions of a calyx.

*Parallel* veins start several together from the base of a leaf, diverge slightly, then proceed parallel and simple, and converge at the apex.

*Parietal*; on the inner surface of an ovary.

*Patent*; spreading widely.

*Pedate*; palmate of three lobes with the lateral lobes having similar large lobes on their upper edge.

*Pedicel*; the branch of a peduncle.

*Peduncle*; flowerstalk.

*Pellucid*; nearly transparent.

*Peltate*; when its point of attachment is on the face, not at the edge of a leaf or other organ.

*Pendulous*; seeds hanging from the top of an ovary.

*Pentagonal*; with five angles having convex spaces between them.

- Pentangular* ; with five angles and five flat or concave faces.
- Perennial* plants live several years and flower more than once, usually many times.
- Perfect* flowers have both stamens and pistils in an efficient state.
- Perfoliate* ; when the leaf completely surrounds the stem so that the latter seems to pass through it.
- Perianth* ; the floral whorls when the calyx and corolla are not distinguishable.
- Pericarp* ; seed-vessel, including adhering calyx if present.
- Perigynous* ; when the corolla and stamens are borne on the calyx but free from the ovary.
- Persistent* ; not soon falling off.
- Personate* ; a gamopetalous two-lipped corolla of which the lower lip is pressed upwards so as to close the opening.
- Petal-like* ; resembling petals in texture and colour.
- Petals* ; the divisions of the corolla.
- Petiolate* ; having a petiole.
- Petiole* ; the stalk of a leaf: *petiolule* ; of a leaflet.
- Phænogamous* ; visibly furnished with stamens and pistils.
- Phanerogamous* ; phænogamous.
- Phyllaries* ; the scales or bracts of the involucre of Composites.
- Pilose* ; with scattered rather stiff hairs.
- Pinnæ* ; the segments of a pinnate leaf.
- Pinnate* ; when leaflets are arranged on opposite sides of a common stalk. A leaf is 2- or 3-pinnate when its primary or secondary divisions are pinnate.
- Pinnatifid* ; a leaf deeply cut into segments nearly to the midrib. A 2- or 3-pinnatifid leaf corresponds to a 2- or 3-pinnate leaf.
- Pinnules* ; the segments of a bipinnate leaf.
- Pistil* ; the ovary, style, and stigma taken together.
- Pith* ; a column of cellular tissue in the centre of the stem and branches of Dicotyledons.
- Pitted* ; covered with small depressed spots.
- Placenta* ; the part of the carpel from which the ovules spring.
- Plane* ; flat ; also an imaginary flat surface in which things are placed.
- Plicate* ; plaited.
- Plumule* ; the ascending leafy part of the embryo.
- Pod* ; a 1-celled and 2-valved seed-vessel with the seeds arranged along the inner angle.
- Pollen* ; the dust in the anther.
- Polygonal* ; with many angles.
- Polypetalous* ; with many separate petals.
- Polysepalous* ; with many separate sepals.

*Pome* ; a compound fleshy many-seeded fruit, an apple or fruit resembling it.

*Pores* ; small, often roundish, holes.

*Porrect* ; extending forwards.

*Posterior* ; the part of the flower nearest to the axis.

*Prickles* ; hardened epidermal appendages resembling thorns, but not woody.

*Primordial* ; the first flower of inflorescence.

*Procumbent, prostrate* ; lying on the ground.

*Prolonged* ; drawn out into a long point, like acuminate, but with no hollowing at the sides.

*Pubescence* ; closely adpressed down.

*Pubescent* ; with pubescence.

*Pulverulent* ; covered with fine powdery matter.

*Punctate* ; having minute spots like pin-holes, real or apparent.

*Pyramidal* ; nearly in the shape of a pyramid.

*Pyriform* ; pear-shaped.

*Quadrata* ; squarish.

*Raceme* ; a spike with stalked flowers : hence

*Racemose* ; flowering in a raceme.

*Rachis* ; the central stem of some kinds of inflorescence ; as the stalk common to several spikelets of Grasses ; the stalk of the frond of Ferns above the lowest pinnæ.

*Radiate flowers* ; those at the margin of a head or other inflorescence which are long and spreading like rays.

*Radical* ; springing from just above the root.

*Radicle* ; the end of the embryo from which the root grows ; also small roots.

*Raphides* ; minute needle-shaped crystals found in the cells of some plants.

*Rays* (see *Radiant*) ; parts diverging in a circle from a central point.

*Receptacle* ; the dilated top of the stalk bearing the flowers in Composites ; the common support of the parts of a flower.

*Reclinate and reclining* ; curved downwards.

*Recurved* ; bent moderately backwards.

*Reflexed* ; bent considerably backwards.

*Reniform* ; transversely oval, but broadly cordate at the base.

*Repand* ; with a rather wavy margin.

*Reticulate* ; forming a network.

*Retorse* ; directed from the point of an organ.

*Retuse* ; abruptly blunt with a notch in the middle.

*Revolute* ; rolled back, as towards the underside of a leaf.

*Rhizomatous* ; having rhizomes.



- Rhizome* ; a prostrate more or less subterranean stem producing roots and leafy shoots.
- Rhomboidal* ; approaching a quadrangular, not square, figure attached by one of its more acute angles.
- Ringent* ; a 2-lipped widely open corolla.
- Rootstock* ; a thick short rhizome or tuber.
- Rosette* ; a collection of leaves growing close together, like the petals of a double rose.
- Rosulate* ; arranged in a rosette.
- Rotate* ; a monopetalous corolla with a short tube and very spreading limb.
- Rudimentary* ; imperfectly developed.
- Rugose* ; covered with a network of lines enclosing convex spaces.
- Rugulose* ; finely rugose.
- Runcinate* ; where the lobes of leaves are directed towards the base.
- Runner* ; a prostrate shoot rooting at its end ; a stolon.
- Sagittate* ; like the barbed head of an arrow, the auricles or lobes pointing backwards.
- Salvershaped* ; a corolla with a long slender tube and flat limb.
- Scabrous* ; rough like a blacksmith's hand.
- Scales* ; minute rudimentary leaves ; very small flat semidetached parts of the cuticle.
- Scape* ; a leafless radical peduncle.
- Scarious* ; very thin, dry, and semitransparent.
- Scorpioid* ; said of the branches of a cyme curved in a circinate manner, and the flowers produced only on the upper side.
- Secund* ; all turned towards one side.
- Seed* ; the ovule arrived at maturity.
- Seedstalk* ; the stalk connecting the hilum of a seed with the placenta.
- Sepals* ; the divisions of the calyx.
- Septicidal* ; when a fruit splits through the middle of the septa or partitions.
- Septifragal* ; when a fruit splits by the separation of the backs of the carpels from the septa.
- Septum* ; the division of an ovary formed by the inflexed edges of the carpels.
- Serrate* ; toothed like a saw.
- Serratures* ; teeth like those of a saw.
- Serrulate* ; with very small sawlike teeth.
- Sessile* ; without a stalk.
- Seta* ; a bristle ; a bristle tipped with a gland ; a slender straight prickle.
- Setaceous* ; like a bristle.
- Setose* ; bearing bristles or setæ usually ending in glands.

- Sheath*; the lower part of a leaf or its petiole, which forms a vertical sheath surrounding the stem. It is sometimes found alone.
- Silicle*; a silique not four times as long as broad.
- Silique*; a long podlike fruit of Crucifers having its edges connected by an internal membrane.
- Simple*; not compound; not branched.
- Sinuate*; having many large blunt lobes and notches.
- Slashed*; with deep tapering incisions.
- Smooth*; free from all kinds of roughness.
- Sobole*; a creeping underground stem producing roots and leaf-buds at intervals; an underground stole.
- Soboliferous*; having soboles, or long underground shoots ending in suckers.
- Solitary*; growing singly.
- Spadix*; a succulent spike bearing many sessile closely placed flowers.
- Spath*; a large bract often enclosing a spadix.
- Spathulate*; oblong, with a long linear claw or haft.
- Spike*; a long simple axis with many sessile flowers; hence *spicate* flowers.
- Spikelet*; the small group of flowers in Grasses enclosed within one or more glumes.
- Spine*; a stiff sharp woody persistent thorn.
- Spinous*; furnished with spines.
- Spinulose*; with small, often very minute spines or prickles.
- Spiral vessels*; fine tubes composed of membrane with spirally twisted fibres internally.
- Sporules*; the seedlike reproductive bodies of flowerless plants.
- Spur*; a tubular extension of the lower part of a petal or gamopetalous corolla; a loose prolongation of the base of a leaf beyond its point of attachment.
- Spurred*; furnished with a spur.
- Squarrose*; covered with appendages spreading at right angles or more.
- Stamen*; the male organ of a flower, usually formed of a filament and anther.
- Staminode*; a scale on the inside of the upper lip of some Scrophulariaceæ.
- Standard*; the upper or posterior petal of a Pea-flower, which is outside the others in the bud.
- Starlike*; applied to flowers of which the petals are narrow and distant and radiant like a star.
- Stellate*; radiating from a centre like a star.
- Stellulate*; like minute stars.
- Stigma*; the cellular part at the top of a carpel or style to which the pollen adheres.

*Stigmatic disk* ; a broad surface at the top of the style, or forming the whole of it, upon which the stigmas are placed.

*Stipe* ; the stalk of Ferns up to the lowest pinna.

*Stipules* ; leaflike appendages at the base of the petiole.

*Stipulodes* ; spines beneath the whorls of branches in *Chara*.

*Stole* ; a lax trailing shoot from the crown of the root, rooting at intervals.

*Stoloniferous* ; having stoles.

*Stomates* ; minute organic openings in the skin of plants.

*Strapshaped* ; not very narrow nor long, and with nearly parallel sides.

*Streak* ; a straight line of peculiar colour or structure, or a furrow.

*Striæ* ; very slight furrows or ridges.

*Striate* ; with slender streaks or furrows.

*Striped* ; having coloured streaks.

*Stripes* ; the vittæ of Umbellifers.

*Style* ; the space between the ovary and stigma.

*Stylopode* ; a fleshy disk crowning the ovary and supporting the style of Umbellifers.

*Sub* ; in composition means a near approach to ; as subrotund is nearly round.

*Subulate* ; awlshaped, tapering from the base to a fine point, a long narrow triangle.

*Sucker* ; a stem produced at the end of an underground shoot.

*Superior* ; above any thing ; a calyx is superior when its tube is wholly attached to the ovary, half-superior when attached only to the lower half of it ; an ovary is superior when wholly free from the calyx ; a part of a flower placed next to the axis.

*Suspended* ovules hang down from near the top of the ovary.

*Suture* ; the line of junction of similar organs cohering.

*Swordshaped* ; very long, narrow, nearly parallel-sided, sharp-edged, acute.

*Sympode* ; a stem formed of a series of superposed branches so as to resemble a continuous axis.

*Syncarpous* ; fruit formed of cohering carpels.

*Syngenesious* flowers form a head and have 5 stamens with united anthers.

*Tailed* ; having a long slender point.

*Tassel-like* ; resembling a silken tassel or painter's camel's-hair pencil.

*Tendril* ; a twisting slender organ for laying hold of objects.

*Terete* ; having a nearly round transverse section.

*Ternate* ; growing in threes about the same point of a stem.

- Testa*; the outer coat of a seed.
- Testaceous*; brownish yellow.
- Tetragonous*; with four angles and four convex faces.
- Thorn*; an abortive branch with a sharp point; distinguished from a prickle by being woody.
- Three-veined*; having three veins, usually of nearly equal size, proceeding from the base. Sometimes leaves are *falsely three-veined* when the ends of a series of lateral veins combine to form a submarginal vein near each edge of the leaf.
- Throat*; the orifice of the tube of a gamopetalous corolla or gamosepalous calyx.
- Thyrsoïd*; having a close-branched raceme of which the middle is broader than the ends.
- Tomentose*; covered with cottony entangled hairs, forming a matted shagginess called *tomentum*; felted.
- Torulose*; uneven, alternately elevated and depressed like a knotted cord.
- Torus*; the part within the calyx to which the floral organs are attached.
- Transversely*; applied to forms like oval when attached by one of their longer sides.
- Triangular*; with three angles and three flat faces.
- Trichotomous*; in forks of three prongs.
- Trifarious*; arranged in three rows.
- Trifid*; dividing about halfway down into three parts.
- Trifurcate*; forked with three nearly equal prongs.
- Trigonal*; with three angles and three convex faces.
- Tripartite*; divided into three parts nearly to its base.
- Triquetrous*; having three angles and three concave faces.
- Truncate*; blunt as if cut off at the end.
- Tube*; the pipe formed by the cohesion of the parts of a floral whorl.
- Tuber*; a thickened underground fleshy part of the stem.
- Tubercles*; little round knobs.
- Tubercular*; *tubercled*; covered with little knobs.
- Tuberous*; like a tuber, but not part of the stem.
- Tubular*; hollow and nearly cylindrical.
- Tumid*; swollen.
- Turbinate*; topshaped, conical and attached by its long point.
- Two-edged*; compressed so as to have two sharp edges.
- Umbel*; when many stalked flowers spring from one point and reach about the same level. *Partial* umbels are umbels seated upon the branches of an umbel, when the whole forms a *compound* umbel.
- Umbilicate*; peltate, but having the attached organ hollowed to receive the top of the stalk.

*Unilateral*; turned to one side.

*Upper part of a floral whorl*; that next the main axis of the stem; posterior.

*Urceolate*; like a pitcher contracted at the mouth.

*Utricle*; a bladder-like covering as in *Chenopodiaceæ*, i. e. an achene with a membranous pericarp. The envelope of the nut of *Carex*.

*Valvate*; having *valves* or parts of an organ opening like little doors; or organs touching only along their edges.

*Veins*; bundles of vessels in leaves and their modifications.

*Ventricose*; swelling unequally on one side.

*Vernation*; the arrangement of leaves in a bud.

*Versatile*; swinging freely on its support, as an anther attached by one point of its back.

*Villose*; shaggy with loose long soft hair.

*Viscous*; clammy.

*Vittæ*; linear receptacles of oil in the fruits of Umbellifers stripes.

*Viviparous*; bearing young plants in the place of flowers.

*Wedgeshaped*; like a wedge, but attached by its point.

*Whorl*; formed of similar organs arranged in a circle round an axis.

*Whorled*; arranged in whorls.

*Winged*; having leaflike or membranous expansions.

*Wings*; the lateral petals of a Pea-flower; the flat membranous appendages of some seeds.

---

When two terms are combined, as *ovate-lanceolate*, it means that the form or structure is compounded of the two, or lies between them.

## SYNOPSIS

OF THE

## NATURAL ORDERS OF BRITISH PLANTS.

---

THIS Synopsis has been prepared for the purpose of facilitating the discovery of the ORDER to which an unknown British Plant belongs. *It must be used with caution, as a very slight error will totally mislead.* The student must always commence with the pair of characters numbered 1 in the left-hand margin; and having determined with which of these his plant agrees, proceed similarly with the group of characters referred to by the number on the right-hand side of the page, and so on.

For instance, having gathered a *Hawthorn*, he finds it to agree with the *second* character of number 1, the *first* of number 3, the *second* of number 4, the *third* of number 29, and the *first* of number 30. It therefore belongs to ORDER XXVI. ROSACEÆ and Suborder POMEÆ. Then turning to the body of the *Manual* (p. 97), he will examine the specimen by the characters given for that Order and its Suborder. Finding it to agree with them, a perusal of the generic definitions placed under POMEÆ will show that it is a *Cratægus*. After a little experience in the examination of plants, the eye becomes so familiar with the principal Orders as to render this process unnecessary, except in doubtful cases.

1	Leaves straight- or parallel-veined (belonging generally to plants having 3-parted floral whorls or a 6-parted perianth) * . . . . .	2
	Leaves net-veined (belonging generally to plants having 5- or 4-parted floral whorls) † . . . . .	3
2	Seeds in a seed-vessel . . . . .	62
	Seeds apparently naked on an axillary scale or in a fleshy cup and solitary. Male fl. in catkins. L. linear or subulate . . . . . lxxix. <i>Coniferæ</i> .	
3	Fl. with a calyx and corolla . . . . .	4
	Fl. with a perianth (calyx undistinguishable from corolla) or none . . . . .	50
4	Corolla polypetalous, inferior . . . . .	5
	Corolla polypetalous, superior . . . . .	29
	Corolla gamopetalous (petals cohering), superior . . . . .	34
	Corolla gamopetalous, inferior . . . . .	39
5	Ovaries many, distinct or united, each bearing a style; or solitary with one lateral placenta . . . . .	6
	Ovary solitary; placentas 2 or more, parietal or on the dissepiments, not forming a central axis . . . . .	13
	Ovary solitary; placentas central . . . . .	17
6	Corolla regular . . . . .	7
	Corolla irregular . . . . .	10
7	Sepals distinct. Stamens hypogynous . . . . .	8
	Sepals more or less combined below . . . . .	9
8	Stamens few. Anth. adnate; connective extending beyond them. . . . . lxxx. <i>Trilliaceæ</i> .	
	Stamens indefinite, usually many (when few, alternate with the petals). Anth. at top of filament, opening by two longitudinal clefts. . . . . i. <i>Ranunculaceæ</i> .	
	Stamens as many as and opposite to the petals. Anth. at top of filament. . . . . ii. <i>Berberidaceæ</i> .	

\* The leaves of these plants (*Monocotyledones*) are nearly always parallel-veined. A few exceptions are *Dioscoreaceæ* (*Tamus*), *Trilliaceæ* (*Paris*), and *Araceæ* (*Arum*), which will be found by following either series of characters. Care must be taken not to confound leaves having *parallel veins which are connected by simple transverse veins* with *net-veined* leaves. The broad leaves of *Alisma*, *Potamogeton*, and *Hydrocharis* are examples of the former. Pinnatifid leaves with linear lobes as in *Anemone Pulsatilla*, may be mistaken for parallel-veined leaves; also the submersed leaves of *Ranunculi*.

† The narrow leaves of some of these plants (*Dicotyledones*) are apparently parallel-veined. They are chiefly aquatics:

Leaves divided in a pinnatifid way.

xxx. *Haloragaceæ* (*Myriophyllum*).

Leaves repeatedly forked . . . . . lxxiii. *Ceratophyllaceæ*.

Leaves simple . . . . . xxx. *Haloragaceæ* (*Hippuris*).





- 23 Stamens opposite the petals. Sepals 2.  
xxxii. *Portulacææ*.
- Stamens opposite the petals. Sepals 3—5. . . . . 24
- 24 Stamens 10 or fewer. Stipules none.  
xiv. *Caryophyllacææ* (*Alsineæ*).
- Stamens 5. Stipules present. Petals distinct.  
xiv. *Caryophyllacææ* (*Polycarpeæ*).
- Stamens 5. Stipules present. Petals subulate.  
xxxiii. *Paronychiacææ*.
- 25 Stamens free . . . . . 27
- Stamens polyadelphous . . . . . xvii. *Hypericacææ*.
- Stamens monadelphous or inserted in an hypogynous ring . . . . . 26
- 26 Stamens 4—5. Stipules none . . . . . xxii. *Linacææ*.
- Stamens 10. Stipules none or united to the base of the petioles. Cells of ovary 5; each many-seeded.  
xxi. *Oxalidacææ*.
- Stamens 10. Stipules present. Cells of ovary 5; each 1-seeded . . . . . xix. *Geraniacææ*.
- 27 Style 1 . . . . . 28
- Styles 3—5. Anthers terminal . . . . . xiii. *Elatinacææ*.
- Styles 3—5. Filaments extending beyond the anthers.  
lxxx. *Trilliaceææ*.
- 28 Stamens 3. Petals 3. Fruit fleshy. lxxi. *Empetracææ*.
- Stamens and petals 4 or 5, inserted in an hypogynous disk. . . . . xxiii. *Celastracææ*.
- Stamens 5 or more. Petals 5. Caps. 1-celled, 3—4-valved . . . . . xi. *Frankeniaceææ*.
- Stamens usually 8. Petals 5. Fruit winged, separating into two capsules . . . . . xviii. *Aceracææ*.
- Stamens 8—10. Capsules 5-celled, 5-valved.  
xlvii. *Ericacææ* (*Pyroleææ* and *Monotropææ*).
- 29 Ovary 1-celled with one pendulous ovule. Stamens as many as and opposite to the petals.  
xl. *Loranthacææ*.
- Ovary 1-celled; placentas 2 parietal; ovules many. Stamens and petals 4—5, alternating.  
xxxv. *Ribesiaceææ*.
- Ovary 2—many-celled; placentas central. Or a pome. 30
- 30 Stamens many, indefinite. Fruit a 1—5-seeded pome.  
xxvi. *Rosacææ* (*Pomææ*).
- Stamens as many as and alternating with the petals, or twice as many . . . . . 31
- 31 Petals imbricate in the bud . . . . . 32
- Petals valvate in the bud . . . . . 33
- Petals twisted in the bud. Sepals valvate. Style 1.  
xxix. *Onagraceææ*.
- 32 Petals 5. Stamens 5. Styles 2. Inflorescence umbellate. Seeds solitary. . . . . xxxvii. *Umbelliferææ*.

- Petals 4—5. Stamens 5—10. Styles 2. (Calyx sometimes inferior.) Capsule 2-valved. Seeds many. (Fruit with two horns) . . . . .xxxvi. *Saxifragaceæ*.
- Petals 4. Stamens 4—8. Styles 4. Fruit of 4 hard nuts . . . . .xxx. *Haloragaceæ*.
- 33 Fruit a berry. Styles more than two. Leaves alternate. . . . .xxxviii. *Hederaceæ*.
- Fruit a drupe. Style 1. Leaves opposite. . . . .xxxix. *Cornaceæ*.
- Fruit dry. Styles 2. Leaves alternate. . . . .xxxvii. *Umbelliferæ* (in part).
- 34 Stamens inserted beneath an epigynous disk. . . . .xlvii. *Ericaceæ* (*Vacciniæ*).
- Stamens inserted with the corolla and free from it . . . . .35
- Stamens inserted on the tube of the corolla or between its lobes . . . . .36
- 35 Filaments free . . . . .xlvi. *Campanulaceæ*.
- Filaments united into 3 bundles . . . . .xxxi. *Cucurbitaceæ*.
- 36 Anthers united. Flowers in a head. . . . .xlv. *Compositæ*.
- Anthers free . . . . .37
- 37 Flowers in a head. Calyx double. . . . .xliv. *Dipsacaceæ*.
- Flowers corymbose or cymose . . . . .38
- 38 Fruit a double indehiscent pericarp, 2-celled, 2-seeded. . . . .xlii. *Rubiaceæ*.
- Fruit dry, with 1 perfect cell, 1-seeded; and often 2 empty cells. Stamens 1—3 . . . . .xliii. *Valerianaceæ*.
- Fruit fleshy, with 1 or several seeds. Stamens 4—5. . . . .xli. *Caprifoliaceæ*.
- 39 Ovary and fruit 4-lobed, separating into 4 small 1-seeded nuts. Style from base of ovary . . . . .40
- Fruit of several follicles. . . . .xxxiv. *Crassulaceæ* (Cotyledon).
- Ovary and fruit simple. Style terminal. . . . .41
- 40 Stamens 5. Corolla regular. Leaves alternate. . . . .liv. *Boraginaceæ*.
- Stamens 4, didynamous, or 2. Corolla 2-lipped. Leaves opposite . . . . .lviii. *Labiataæ*.
- 41 Ovary 1-celled, 1-seeded. Calyx tubular. Stamens 5. Styles 5 . . . . .lxii. *Plumbaginaceæ*.
- Ovary 1-celled, many-seeded . . . . .42
- Ovary with 2 or more cells (but fruit sometimes 1-celled) . . . . .44
- 42 Corolla scarious, regular, 4-parted. Stamens 4. . . . .lxiii. *Plantaginaceæ*.
- Corolla coloured . . . . .43
- 43 Corolla irregular. Stamens 2. . . . .lx. *Lentibulariaceæ*.
- Corolla regular. Stamens 4—5, opposite to the segments of the corolla . . . . .lxi. *Primulaceæ*.
- Corolla irregular. Stamens 4, didynamous. . . . .lvi. *Orobanchaceæ*.

- 44 Stamens hygogynous, scarcely attached to the corolla, distinct. . . . .xlvii. *Ericaceæ* (*Ericææ* and *Arbutææ*).  
Stamens upon the corolla; filaments connected.  
xii. *Polygalaceæ*.  
Stamens upon the corolla, distinct . . . . . 45
- 45 Stamens 2. Corolla regular. . . . .xlix. *Oleaceæ*.  
Stamens 2 or 4, and didynamous. Corolla irregular. . . . . 46  
Stamens 4 or 5, not didynamous . . . . . 47
- 46 Ovary 2-celled, not lobed; placentas central.  
lvii. *Scrophulariaceæ*.  
Ovary 2—4-celled, lobed . . . . .lix. *Verbenaceæ*.
- 47 Cells of ovary each with 1 or 2 ovules . . . . . 48  
Cells of ovary each with many ovules. . . . . 49
- 48 Fruit fleshy, not bursting. Stigmas sessile.  
xlviii. *Aquifoliaceæ*.  
Fruit a capsule, bursting. Styles manifest.  
liii. *Convolvulaceæ*.
- 49 Fruit a double follicle . . . . .l. *Apocynaceæ*.  
Fruit 2- or imperfectly 4-celled. Leaves alternate.  
lv. *Solanaceæ*.  
Fruit 1- or imperfectly 2-celled, 2-valved. Leaves opposite . . . . .li. *Gentianaceæ*.  
Fruit 3-celled, 3-valved . . . . .lii. *Polemoniaceæ*.
- 50 Flowers not in catkins . . . . . 51  
Male flowers in catkins . . . . .lxxviii. *Amentiferæ*.  
Flowers inconspicuous (rarely found). Plant formed of minute leaflike fronds, floating freely.  
xciv. *Lemnaceæ*.
- 51 Ovary one, superior. Perianth sometimes wanting . . . . . 54  
Ovary inferior . . . . . 52  
Ovaries many, distinct . . . . . 53
- 52 Stamens 1 or 8. Fruit not bursting, 1- or 4-celled, 1- or 4-seeded. Limb of perianth of female flower very minute . . . . .xxx. *Haloragaceæ*.  
Stamens 4 or 5. Fruit not bursting, 1-celled, 1-seeded. Limb of perianth manifest. . . . .lxix. *Santalaceæ*.  
Stamens 4. Style filiform. Ovary 4-celled, many-seeded . . . . .xxix. *Onagraceæ*.  
Stamens 5, syngenesious. . . . .xlv. *Compositæ*.  
Stamens 6, on base of perianth. Style 1, trifid. Fl. dicecious . . . . .lxxx1. *Dioscoreaceæ*.  
Stamens 6—12, epigynous. Style short. Stigma radiant. Ovary 3—6-celled, many-seeded.  
lxx. *Aristolochiaceæ*.
- Stamens 8—10. Styles 2. Ovary 1-celled, 2-beaked. many-seeded, opening like a cup.  
xxxvi. *Saxifragaceæ*.

- 53 Sepals petal-like, deciduous. Anthers extrorse.  
     i. *Ranunculaceæ* (*Caltha*).  
 Sepals petal-like, persistent. Anthers introrse.  
     i. *Ranunculaceæ* (*Pæonia*). 55
- 54 Fruit separating into several carpels. No perianth.... 56  
 Fruit not separating into carpels .....
- 55 Carpels 4, not bursting. Stamen 1.  
     lxxiv. *Callitrichaceæ*.  
 Carpels 3 or 2, opening, separating with elasticity.  
     lxxii. *Euphorbiaceæ*.
- 56 Leaves with stipules ..... 57  
 Leaves without stipules ..... 58
- 57 Stipules sheathing the stem .....lxxvi. *Polygonaceæ*.  
 Stipules attached to the petiole.  
     xxvi. *Rosaceæ* (*Sanguisorbææ*).  
 Stipules free, deciduous. Ovary 1-celled. Perianth  
 4—5-parted .....lxxv. *Urticaceæ*.  
 Stipules free, deciduous. Ovary 2-celled.  
     lxxvii. *Ulmaceæ*.  
 Stipules free, deciduous. Ovary 1-celled. Female  
 perianth scale-like, open ....lxxvi. *Cannabinaceæ*.
- 58 Flowers monœcious or diœcious ..... 59  
 Flowers perfect or polygamous .....
- 59 Fruit fleshy. Male fl. each of one 2-celled naked stamens, crowded together on a spadix.  
     xciii. *Araceæ* (*Arææ*).  
 Fruit fleshy. Stamens 2 or 3. Perianth of scales imbricated in several rows .....lxxi. *Empetraceæ*.  
 Fruit dry. Stamens 3 or more. Perianth tubular.  
     lxvii. *Elæagnaceæ*.  
 Fruit dry. Stamens 12—20. Perianth 10—12-cleft.  
     lxxiii. *Ceratophyllaceæ*.
- 60 Perianth hardened over the fruit.  
     xiv. *Caryophyllaceæ* (*Scleranthææ*).  
 Perianth not hardened over the fruit..... 61  
 Perianth none. Fruit compressed linear and leaflike at the end.....xlix. *Oleaceæ* (*Fraxinus*).
- 61 Perianth 3—5-cleft, herbaceous ..lxv. *Chenopodiaceæ*.  
 Perianth 3-cleft, scarious .....lxiv. *Amaranthaceæ*.  
 Perianth tubular. Stamens perigynous.  
     lxviii. *Thymelaceæ*.  
 Perianth 6—8-parted, herbaceous. Stamens hypogynous..... lxxx. *Trilliaceæ*.
- 62 Leaves net-veined. Floral envelopes whorled..... 63  
 Leaves net-veined. Perianth none. Fl. monœcious, on a spadix .....xciii. *Araceæ* (*Arææ*).

- Leaves parallel-veined, persistent. Floral envelopes  
whorled, or none . . . . . 64
- Leaves parallel-veined, persistent. Floral envelopes  
imbricate, bractlike . . . . . 75
- 63 Perianth inferior. Flowers perfect . . lxxx. *Trilliaceæ*.  
Perianth superior. Flowers dioecious.  
lxxx. *Dioscoreaceæ*.
- 64 Ovaries several, each bearing a style or stigma . . . . . 65  
Ovary 1, inferior . . . . . 66  
Ovary 1, superior . . . . . 69
- 65 Perianth 4-parted, inconspicuous or none.  
xcv. *Potamogetonaceæ*.  
Perianth 6-parted, conspicuous. Carpels not opening.  
lxxxvi. *Alismaceæ*.  
Perianth 6-parted, conspicuous. Carpels opening at  
their inner edge . . . . . lxxxix. *Melanthaceæ*.
- 66 Stamens and style united in a central column.  
lxxxiii. *Orchidaceæ*.  
Stamens free from the style . . . . . 67
- 67 Perianth wholly petal-like . . . . . 68  
Three outer segments of perianth resembling an herba-  
ceous calyx . . . . . lxxxii. *Hydrocharidaceæ*.  
All the 4 segments of perianth resembling an herba-  
ceous calyx . . . . . xl. *Loranthaceæ*.
- 68 Stamens 3. Anthers bursting outwards.  
lxxxiv. *Iridaceæ*.  
Stamens 6. Anthers bursting inwards.  
lxxxv. *Amaryllidaceæ*.
- 69 Perianth like a corolla of 6 petals. Flowers perfect . . 70  
Perianth more or less glumaceous, or partly coloured,  
or wanting . . . . . 71
- 70 Perianth petal-like, conspicuous. Fruit dry, bursting  
with 3 valves . . . . . lxxxviii. *Liliaceæ*.  
Perianth petal-like, conspicuous. Fruit succulent, not  
bursting . . . . . lxxxvii. *Asparagaceæ*.  
Perianth herbaceous, scale-like. Fruit baccate, not  
bursting . . . . . xciii. *Araceæ* (*Orontææ*).
71. Flowers perfect. Perianth 6-parted. Stamens 6.  
xc. *Juncaceæ*.  
Flowers monœcious . . . . . 72
- 72 Flowers on a spadix . . . . . 73  
Spadix wanting . . . . . 74  
Flowers solitary . . . . . xcvi. *Naiadaceæ*.
- 73 Flower on a thick spadix. Spath wanting. Anthers  
wedged-shaped, erect; filaments long.  
xcii. *Typhaceæ*.

Flowers on a thick spadix in a spath. Perianth none.

Anthers ovate; filaments very short.

xciii. *Araceæ* (Areæ).

74 Flowers capitate. Stamens 2—5 . . xci. *Eriocaulaceæ*.

Flowers 2 in a spath, one male, one female. (Floating.)

xciv. *Lemnaceæ*.

Flowers in two rows on one side of a spath, enclosed by  
a fold of the leaf . . . . .xcvi. *Naiadaceæ*.

75 Leaves with entire sheaths. Anthers entire at the  
ends . . . . .xcvii. *Cyperaceæ*.

Leaves with split sheaths. Anthers notched at both  
ends . . . . .xcviii. *Gramineæ*.

## CLASSES, DIVISIONS, AND ORDERS

OF

## BRITISH PLANTS.

Class I. DICOTYLEDONES<sup>1</sup>.

Stems when perennal composed of bark, wood, and pith. The wood furnished with medullary rays and increasing by the addition of concentric layers externally. Leaves usually net-veined. Cotyledons 2 or more, opposite or whorled.—Each floral whorl composed of 5 or 4 parts.

Division 1. THALAMIFLORÆ. Pet. distinct (rarely 0) and as well as the stam. growing separately from the sepals, hypogynous .....	1
Division 2. CALYCIFLORÆ. Pet. distinct and, as well as the stam., perigynous or epigynous .....	77
Division 3. COROLLIFLORÆ. Pet. united, at least at the base. Stam. mostly epipetalous .....	169
Division 4. MONOCHLAMYDÆ. Only a single perianth or none .....	297
Division 5. GYMNOSPERMÆ. Ovules and seeds apparently naked. Ovary and styles wanting. Perianth wanting.	325

## Division 1. THALAMIFLORÆ.

\* *Apocarpous.*

1. <i>Ranunculaceæ.</i> Stam. polyandrous (rarely pentandrous). Pistils usually many, of achenes or follicles.—Rarely (in <i>Actæa</i> ) a many-seeded berrylike carpel .....	1
2. <i>Berberidaceæ.</i> Stam. 6 or 4, opposite the petals; anth. opening by valves from the bottom .....	15

<sup>1</sup> The characters are drawn to suit our plants.

\*\* *Appearing syncarpous from the fleshy disk enveloping the ovaries.*

3. *Nymphaeaceæ*. Polyandrous. Pet. many, seated with the many stamens upon a fleshy disk.—Aquatic. Rhizome prostrate in mud. Leaves floating ..... 15

\*\*\* *Syncarpous with parietal placentas.*

4. *Papaveraceæ*. Polyandrous. Sep. 2, deciduous. Pet. 4. Pistil 1 ..... 16

5. *Fumariaceæ*. Stam. in two bundles of 3 each; lateral anth. of each bundle 1-celled. Sep. 2, minute. Pet. 4, irregular, one or more of the upper gibbous or spurred at the base. Fl. very irregular ..... 19

6. *Cruciferae*. Stam. tetradynamous. Sep. 4. Pet. 4.—Fl. cruciform..... 21

7. *Resedaceæ*. Stam. 10—24, from an hypogynous 1-sided disk. Sep. 4—8. Pet. 4—8, irregular. Caps. opening at the top ..... 39

8. *Cistaceæ*. Polyandrous. Sep. 3, equal, and 2 smaller external. Pet. 5, regular, very deciduous ..... 40

9. *Violaceæ*. Stam. 5; anth. on inner side of dilated filaments which slightly cohere. Pet. 5, irregular; lower spurred ..... 41

10. *Droseraceæ*. Stam. 5. Cal. 5-cleft. Pet. 5, regular.—L. bearing long hairs tipped with glands ..... 44

11. *Frankeniaceæ*. Stam. 4 or more. Cal. 4—5-cleft, furrowed. Pet. 4—5, regular.—L. with revolute edges, not glandular ..... 45

\*\*\*\* *Syncarpous with axile placentas.*

† Fl. irregular.

12. *Polygalaceæ*. Stam. 8, in 2 equal opposite bundles. Anth. 1-celled ..... 45

†† Fl. regular (except in Ord. 20).

13. *Elatinaceæ*. Pet. 3—5. Stam. as many or twice as many as the petals. Caps. 3—5-celled.—L. opposite. Plants aquatic ..... 47



14. *Caryophyllaceæ*. Pet. 5 or 4. Stam. 8—10, rarely 5 or 4. Caps. 1-celled with a free central placenta.—L. opposite . . . . . 47
15. *Malvaceæ*. Pet. 5, twisted. Stam. many, monadelphous. Anth. 1-celled. Ovary of several carpels round a common axis. L. alternate . . . . . 65
16. *Tiliaceæ*. Pet. 5. Stam. many, free or polyadelphous. Anth. 2-celled. Ovary 5-celled.—Trees with alternate leaves . . . . . 66
17. *Hypericaceæ*. Pet. 5, twisted. Stam. many, in 3—4 bundles. Anth. 2-celled . . . . . 67
18. *Aceraceæ*. Pet. 5 (in our plants), imbricate. Stam. distinct, generally 8, on an hypogynous disk. Fr. of 2, winged, 1-seeded nuts.—Trees with opposite leaves. 70
19. *Geraniaceæ*. Pet. 5, twisted. Stam. monadelphous. Fr. of 5 carpels cohering together with their styles round a long beaked persistent axis, from which they ultimately separate with elasticity . . . . . 71
20. *Balsaminaceæ*. Pet. 5, irregular, the lateral united in pairs, imbricate. Stam. 5. Anth. slightly cohering. Fr. of 5 carpels with elastic valves . . . . . 74
21. *Oxalidaceæ*. Pet. 5, twisted. Stam. 10. Anth. free. Caps. 5-valved.—L. usually ternate . . . . . 75
22. *Linaceæ*. Pet. 4—5, twisted. Stam. 4—5, with intermediate teeth. from an hypogynous ring. Carp. separating but leaving no central axis . . . . . 76
- (Ord. 48. *Agufoliaceæ* with sep. free, cor. 4—5-parted, stam. 4—5, an evergreen tree, is sometimes placed here.) 237
- (Ord. 71. *Empetraceæ* with diœcious flowers, a heath-like evergreen shrub, is sometimes placed here.) . . . . . 313

## Division 2. CALYCIFLORÆ.

\* *Stam. and petals mostly perigynous.*

23. *Celastraceæ*. Fl. greenish. Stam. alternating with the petals and inserted with them in a fleshy disk surrounding the ovary.—Trees or shrubs . . . . . 77

24. *Rhamnaceæ*. Fl. greenish. Stam. opposite the petals and inserted with them in a disk lining the calyx.—Shrubs ..... 78
25. *Leguminosæ*. Fl. coloured, irregular, papilionaceous; odd petal superior and external. Stam. 10, di- or rarely monadelphous. Fruit a legume..... 78
26. *Rosaceæ*. Fl. regular; odd petal inferior. Stam. usually many, perigynous. Carp. distinct when young, and superior, or connate, or adnate to calyx (falsely syncarpous and inferior) ..... 97
27. *Lythraceæ*. Cal. tubular. Pet. and stam. perigynous. Ovary free, syncarpous. Style 1 ..... 132
28. *Tamariscaceæ*. Sep. slightly connate below. Pet. perigynous. Stam. from the edge of hypogynous ring. Ovary free. Styles 3. Caps. 3-valved, loculicidal.—Shrub with narrow spurred leaves..... 132
29. *Onagraceæ*. Cal. adnate to ovary, with 2—4 lobes. Pet. 2—4. Stam. 2, 4, or 8. Style 1 ..... 133
30. *Haloragaceæ*. Cal. adnate to ovary, with a minute limb. Pet. minute or 0. Stam. 1 or 8. Styles 1—4.—Aquatic. Fl. usually unisexual. (*Callitriche*, p. 318, is often placed here.) ..... 137
31. *Cucurbitaceæ*. Dioecious (in our plant). Cal.-tube adnate to ovary. Stam. 5, resembling 3 by cohesion.—Climbing by tendrils ..... 138
32. *Portulacææ*. Cal.-tube short; sep. 2. Pet. connate or free. Stam. 3—5, free. Ovary 1-celled; placenta free, central ..... 138
33. *Paronychiaceæ*. Cal. 5-parted. Pet. minute resembling barren filaments, or none. Stam. 5, or rarely 10. Ovary free ..... 139
34. *Crassulaceæ*. Sep. and pet. each 5, rarely 3 or 4, or 10—12. Pet. free or connate at the base. Stam. as many or twice as many as petals. Carp. free, same number as petals. Fr. of follicles ..... 141
35. *Ribesiaceæ*. Sep., pet., and stam. 4—5. Ovary inferior; placentas parietal. Style 2—4-cleft. Fr. a berry 144
36. *Saxifragaceæ*. Cal. 4—5-cleft. Pet. 4, 5, or 0, perigynous. Ovary of 2 carpels combined below. Styles distinct ..... 145

\*\* *Stamens and petals epigynous, round an epigynous disk. Cal.-tube adnate to ovary.*

- 37. *Umbelliferae*. Fl. umbellate. Fr. of 2 dry carpels adhering by their faces to a common axis. Styles 2. Stam. 5.—L. alternate ..... 149
- 38. *Hederaceae*. Fl. umbellate. Fr. a 5-celled, 5-seeded berry. Style 1. Stam. 4.—L. opposite ..... 168
- 39. *Cornaceae*. Fl. regular, subumbellate. Fr. a 2-seeded berry. Style 1. Stam. 4.—L. opposite ..... 168

Division 3. COROLLIFLORÆ.

\* *Corolla epigynous. Stam. epipetalous.*

- 40. *Loranthaceae*. L. opposite. Stam. 4, adnate to and opposite to petals.—Pet. scarcely connected.—Parasitical..... 169
- 41. *Caprifoliaceae*. L. opposite. Stam. 4—10, alternate with cor.-lobes, single or in pairs ..... 170
- 42. *Rubiaceae*. L. whorled. Stam. 4—5, alternate with cor.-lobes. Fr. a didymous indehiscent pericarp .. 172
- 43. *Valerianaceae*. L. opposite. Stam. 1—3, fewer than the cor.-lobes. Fr. with 1 perfect fertile cell and 1 seed ..... 176
- 44. *Dipsacaceae*. L. opposite. Fl. in involucrate heads and with a free outer calyx and adnate inner one ..... 179
- 45. *Compositae*. Fl. in involucrate heads; no free outer calyx or involucl. Stam. 5. Anth. cohering as a tube round style ..... 180

\*\* *Corolla epigynous or hypogynous. Stam. not adhering to the corolla.*

- 46. *Campanulaceae*. Cal. superior. Stam. and cor.-lobes 5, alternate. Anth. opening longitudinally.—Anthers free or connate. Herbaceous ..... 227
- 47. *Ericaceae*. Cal. inferior or superior. Stam. 8—10 (5 in *Azalea*). Anth. opening by pores or slits.—Small shrubs ..... 231

\*\*\* *Corolla hypogynous. Stam. epipetalous.*

† Cor. nearly or quite regular.

48. *Aquifoliaceæ.* Sep. free. Cor. 4—5-parted. Stam. as many as and alternate with cor.-lobes.—An evergreen tree. . . . . 237
49. *Oleaceæ.* Cal. 4-fid, or 0. Cor. 4-cleft, or 0. Stam. 2. Anth. attached by back.—Trees and shrubs . . . . . 238
50. *Apocynaceæ.* Cal. 4—5-fid. Cor. 4—5-lobed. Stam. 4—5. Anth. attached by base. Ovaries 2, 1-celled. Fr. 2 follicles.—Prostrate. Herbaceous . . . . . 238
51. *Gentianaceæ.* Cal. 4—8-fid. Cor. 4—8-fid. Stam. 4—8. Anth. versatile. Ovary 1-celled, of 2 carpels . . . . 239
52. *Polemoniaceæ.* Cal. 5-parted. Cor. 5-lobed, regular. Stam. 5. Ovary 3-celled. Stig. 3-fid . . . . . 243
53. *Convolvulaceæ.* Sep. 5, free or connate. Cor. 5-parted. Stam. 5. Ovary 2—4-celled. Stig. 2 . . . . . 244
54. *Boraginaceæ.* Cal. 5-parted. Cor. 5-parted. Stam. 5. Ovary of 2, 2-parted carpels; each lobe 1-seeded. Style from base of lobes, simple.—L. alternate, rough. Fr. of 4 nutlets . . . . . 246
55. *Solanaceæ.* Cal. 5-parted. Cor. 5-parted. Stam. 5. Ovary 2- or sub-4-celled, many-seeded. Fr. a berry or capsule . . . . . 252
63. *Plantaginaceæ.* Cal. 4-parted. Cor. 4-parted, scarious. Stam. 4. Ovary 2—4-celled. Caps. opening transversely. (In *Littorella* the fl. are monœcious, fem.-cal. 3-parted, cor. urceolate, stam. hypogynous.) . . . . . 254
- †† Corolla more or less irregular.
56. *Orobanchaceæ.* Cor. 2-lipped. Stam. 4, didynamous. Ovary 1-celled. Caps. 2-valved, many-seeded.—Root-parasites . . . . . 254
57. *Scrophulariaceæ.* Cor. 2-lipped or personate or nearly regular. Stam. didynamous or rarely 2 or 5. Ovary 2-celled. Style terminal. Caps. 2-celled, many-seeded . . . . . 258
58. *Labiatae.* Cor. 2-lipped or nearly regular. Stam. didynamous or rarely 2 or 4. Ovary 4-lobed. Style from between the lobes. Fr. of 1—4, small, nutlike, 1-seeded parts.—L. opposite. Fl. cymose, in false whorls. . . 271

59. *Verbenaceæ*. Cor. scarcely 2-lipped (in our plant). Stam. didynamous. Ovary 2—4-celled. Style terminal. Fr. a capsule dividing into 4 nuts . . . . . 286
60. *Lentibulariaceæ*. Cor. 2-lipped. Stam. 2. Caps. 1-celled, 2-valved; placenta free, central; seeds many.—Floating plants . . . . . 286
61. *Primulaceæ*. Cor. regular. Stam. 5, rarely 4 or 7, opposite the petals. Caps. opening by 5—10 valves at the top, or by a deciduous cap; placenta central; seeds many. Style 1. . . . . 288
62. *Plumbaginaceæ*. Cor. regular, 5-fid, or rarely 5 free petals. Stam. 5, opposite the petals. Fr. a 1-seeded utricle. Styles 5 . . . . . 293
- \*\*\*\* *Cor. and stam. hypogynous.*
63. *Plantaginaceæ* (in part). The genus *Littorella* with monœcious fl., fem.-cal. 3-parted, cor. urceolate. . . . . 295

Division 4. MONOCHLAMYDEÆ.

64. *Amaranthaceæ*. Monœcious. Perianth 3—5-parted, scarious, persistent, inferior. Stam. opposite perianth-segments. Caps. 1-seeded, not bursting. Style 0; stig. 1 or more . . . . . 297
65. *Chenopodiaceæ*. Monœcious or perfect. Perianth 3—5-parted, herbaceous, persistent, inferior. Stam. opposite perianth-segments. Caps. 1-seeded, not bursting. Styles 2 or more . . . . . 297
66. *Polygonaceæ*. Usually with perfect flowers. Perianth 3—6-parted, herbaceous, or petal-like, inferior. Stam. 5—8, perigynous or hypogynous, opposite perianth-segments. Caps. 1-seeded, not bursting. Styles several.—Nut often enclosed in enlarged perianth. Stipules ocreæ . . . . . 304
67. *Elæagnaceæ*. Diccious. Male perianth 4-parted, female tubular with 2—5-toothed limb. Stam. alternate with perianth-segments, or twice as many, perigynous. Caps. crustaceous enclosed in persistent inferior perianth.—Shrub. . . . . 311
68. *Thymelaceæ*. Perianth tubular, 4—5-lobed, inferior. Stam. 8 in our plant, on the tube. Ovary 1-celled, with 1 pendulous ovule. Fr. a nut or drupe.—Small shrubs . . . . . 311

69. *Santalaceæ*. Perianth superior, 3—5-lobed. Stam. 3—5, opposite perianth-lobes. Ovary 1-celled; ovules several, pendulous from apex of central placenta. . . . . 312
70. *Aristolochiaceæ*. Perianth superior, tubular above, 3-lobed, or with a dilated irregular limb. Stam. 6—12. . . . . 313
71. *Empetraceæ*. Diccious. Perianth of hypogynous scales imbricate in several rows. Stam. 3, free. Ovary superior, 3—9-celled. Ovules solitary.—A small prostrate heathlike shrub. . . . . 313
72. *Euphorbiaceæ*. Mon- or di-ccious. Perianth 3-parted or 0. Stam. 1 or more. Ovary superior, 2—3-lobed and -celled; each cell with 1—2 suspended ovules. Caps. opening with elasticity.—In *Euphorbia* many fl., each of one naked stamen, surround a pendulous 3-lobed naked ovary, enclosed by a calyx-like involucre . . . . . 314
73. *Ceratophyllaceæ*. Monccious. Perianth of many divisions. Stam. many, sessile; anth. 2-celled. Ovary superior, 1-celled. Style 1.—Submerged plants with inconspicuous fl. and multifid leaves . . . . . 317
74. *Callitrichaceæ*. Monccious. Perianth 0, but often 2 white bracts. Stam. 1; filament long; anth. 1-celled. Ovary 4-angled, 4-celled. Styles 2. Fr. of 4, 1-seeded, indehiscent carpels.—Aquatic plants. . . . . 318
75. *Urticaceæ*. Mon- or di-ccious (really perfect). Perianth inferior, 4—5-parted; in fem. fl. tubular, notched. Stam. 4—5. Style filiform or 0 . . . . . 319
76. *Cannabinaceæ*. Diccious. Perianth inferior, of 5 sepals in male, open and hidden in scales of catkin in female. Stam. 5. Styles 2, long.—Long twining plant . . . . . 320
77. *Ulmaceæ*. Fl. 2-sexual, not in catkins. Perianth inferior, bellshaped, 4—8-cleft. Stam. 4—8, on tube, opposite lobes. Ovary 1—2-celled. Styles 2. Fr. dry, flat, winged.—Trees. L. alternate, distichous. Stip. deciduous . . . . . 320
78. *Amentiferæ*. Mon- or di-ccious. Males in catkins. Fem.-perianth 0, or adherent. Fr. bony or membranous (or drupaceous in *Myrica*).—Trees or shrubs . . . . . 321

## Division 5. GYMNOSPERMEÆ.

79. *Coniferæ*. Anth. and ovules on the scales of separate catkins or cones.—Trees or shrubs. . . . . 325

## Class II. MONOCOTYLEDONES.

Stem not separable into bark, wood, and pith; of cellular tissue with vascular bundles embedded irregularly in it. Leaves mostly alternate and sheathing, with parallel simple veins connected by smaller transverse veins, rarely net-veined. Cotyledon one, or, if more, alternate. Floral whorls composed of 3 or 6 parts.

Division 1. DICTYOGENÆ. Leaves net-veined. Floral envelopes whorled .....	337
Division 2. FLORIDÆ. Leaves parallel-veined. Floral envelopes whorled or wanting .....	338
Division 3. GLUMIFERÆ. Leaves parallel-veined, persistent. Floral envelopes rudimentary or wanting. Flowers covered by imbricate bractlike scales .....	387

## Division 1. DICTYOGENÆ.

80. <i>Trilliaceæ</i> . Perianth inferior. Fl. perfect .....	337
81. <i>Dioscoreaceæ</i> . Perianth superior. Fl. dioecious.—Twining plants .....	337

## Division 2. FLORIDÆ.

\* *Perianth superior, petal-like. Syncarpous.*

82. <i>Hydrocharidaceæ</i> . Dioecious. Fl. regular. Sep. herbaceous. Pet. coloured.—Floating plants .....	339
83. <i>Orchidaceæ</i> . Fl. irregular. Stam. 1—2, sessile, adnate to central column bearing the stigma .....	340
84. <i>Iridaceæ</i> . Fl. regular, of 6 petal-like parts. Stam. 3, bursting outwards. Ovary 3-celled, 3-valved .....	352
85. <i>Amaryllidaceæ</i> . Fl. regular, of 6 petal-like parts. Stam. 6, bursting inwards. Ovary 3-celled, 3-valved ..	355

\*\* *Perianth inferior, apocarpous, or carpels separable when ripe.*

86. <i>Alismaceæ</i> . Perianth of 3 sepal-like and 3 petal-like parts. Stam. 6. Fr. of many carpels .....	356
--	-----

\*\*\* *Perianth inferior, syncarpous.*

87. <i>Asparagaceæ</i> . Perianth petal-like. Stam. 6 or 4 or 8; anth. bursting inwards. Fr. a berry.—Rhizome creeping .....	359
--	-----

88. *Liliaceæ*. Perianth petal-like. Stam. 6; anth. bursting inwards. Fr. a dry capsule.—Bulbous or with root of fleshy fibres in *Simethis* ..... 361
89. *Melanthaceæ*. Perianth subherbaceous or petal-like. Stam. 6; anth. versatile from below their middle. Fr. of 3 follicles slightly connected or a 3-celled capsule.—Tuberous, or with a creeping rhizome ..... 367
90. *Juncaceæ*. Perianth of glumelike scarious segments. Stam. usually 6; anth. innate. Fr. capsular, 3-valved, loculicidal.—Rushes ..... 368
91. *Eriocaulaceæ*. Perianth membranous, of 4—5 parts. Stam. 4—6; anth. attached by back. Fr. capsular, membranous, 2—3-celled, loculicidal.—Aquatic plants. L. tufted. Fl. in heads on naked stalks ..... 375
- \*\*\*\* *Perianth wanting in our plants, or of 4 small scalelike parts.*
92. *Typhaceæ*. Monœcious. Fl. closely packed in cylindrical spikes, or in dense globose clusters, the males uppermost. Fr. dry or spongy.—Water-plants ..... 375
93. *Araceæ*. Monœcious with the male fl. uppermost, or perfect, on a spadix and usually in a spathe. Fr. succulent.—Spath usually conspicuous. L. often net-veined. 377
94. *Lemnaceæ*. Monœcious. Fl. 2 in a spathe (very rare).—Small, floating, green, leaflike, proliferous ..... 378
95. *Potamogetonaceæ*. Fl. perfect or monœcious, in small spikes or clusters. No spathe. Ovaries 4; stig. 1, sessile, subcapitate.—Aquatic plants ..... 379
96. *Naiadaceæ*. Mon- or di-œcious. Fl. solitary in a spathe. Ovary 1. Style 1; stig. filiform.—Aquatic plants with linear leaves ..... 386

## Division 3. GLUMIFERÆ.

97. *Cyperaceæ*. Anth. entire at the end, innate. Leaf-sheaths entire.—Sedges ..... 387
98. *Gramineæ*. Anth. notched at both ends, versatile. Leaf-sheaths split.—Grasses ..... 412



## Class III. CRYPTOGAMEÆ.

- \* *Vasculares*. Stems with a few ducts amongst the cellular tissue. Producing spores which develop into a prothallus which bears antheridia and arche-gonia.

99. *Equisetaceæ*. Capsules on the under side of the peltate scales of terminal cones. Stems jointed, with whorled branches.—Horse-tails . . . . . 448

100. *Filices*. Capsules on the edge or underside of fronds, or rarely on separate branch of frond.—Fronds usually circinate . . . . . 451

101. *Marsileaceæ*. Capsules in the cells of globular coriaceous sessile or shortly stalked axillary receptacles . . 465

102. *Lycopodiaceæ*. Capsules in the axils of imbricate leaves or terminal spikes, 2—3-valved . . . . . 465

- \*\* Stems formed of one or more parallel tubes, verticillately branched. Nucules and globules on the branches.

103. *Charuceæ*. Leafless, branched. Fructification of red globules and ovoid nucules with a spirally twisted coat, on the same or different plants . . . . . 468

NATURAL ORDERS OF BRITISH PLANTS.

Class I. *Dicotyledones.*

Division I. THALAMIFLORÆ.

Order I. RANUNCULACEÆ.

II. BERBERIDACEÆ.

III. NYMPHÆACEÆ.

IV. PAPAVERACEÆ.

V. FUMARIACEÆ.

VI. CRUCIFERÆ.

VII. RESEDACEÆ.

VIII. CISTACEÆ.

IX. VIOLACEÆ.

X. DROSERACEÆ.

XI. FRANKENIACEÆ.

XII. POLYGALACEÆ.

XIII. ELATINACEÆ.

XIV. CARYOPHYLLACEÆ.

XV. MALVACEÆ.

XVI. TILIACEÆ.

XVII. HYPERICACEÆ.

XVIII. ACERACEÆ.

XIX. GERANIACEÆ.

XX. BALSAMINACEÆ.

XXI. OXALIDACEÆ.

XXII. LINACEÆ.

Div. II. CALYCIFLORÆ.

XXIII. CELASTRACEÆ.

XXIV. RHAMNACEÆ.

XXV. LEGUMINOSÆ.

XXVI. ROSACEÆ.

XXVII. LYTHRACEÆ.

XXVIII. TAMARISCACEÆ.

Order XXIX. ONOGRACEÆ.

XXX. HALORAGACEÆ.

XXXI. CUCURBITACEÆ.

XXXII. PORTULACEÆ.

XXXIII. PARONYCHIACEÆ.

XXXIV. CRASSULACEÆ.

XXXV. RIBESIACEÆ.

XXXVI. SAXIFRAGACEÆ.

XXXVII. UMBELLIFERÆ.

XXXVIII. HEDERACEÆ.

XXXIX. CORNACEÆ.

Div. III. COROLLIFLORÆ.

XL. LORANTHACEÆ.

XLI. CAPRIFOLIACEÆ.

XLII. RUBIACEÆ.

XLIII. VALERIANACEÆ.

XLIV. DIPSACACEÆ.

XLV. COMPOSITÆ.

XLVI. CAMPANULACEÆ.

XLVII. ERICACEÆ.

XLVIII. AQUIFOLIACEÆ.

XLIX. OLEACEÆ.

L. APOCYNACEÆ.

LI. GENTIANACEÆ.

LII. POLEMONIACEÆ.

LIII. CONVOLVULACEÆ.

LIV. BORAGINACEÆ.

LV. SOLANACEÆ.

LVI. OROBANCHACEÆ.

LVII. SCROPHULARIACEÆ.

LVIII. LABIATÆ.

LIX. VERBENACEÆ.

## Order LX. LENTIBULARIACEÆ.

LXI. PRIMULACEÆ.

LXII. PLUMBAGINACEÆ.

LXIII. PLANTAGINACEÆ.

## Div. IV. MONOCHLAMYDEÆ.

LXIV. AMARANTHACEÆ.

LXV. CHENOPODIACEÆ.

LXVI. POLYGONACEÆ.

LXVII. ELÆAGNACEÆ.

LXVIII. THYMELACEÆ.

LXIX. SANTALACEÆ.

LXX. ARISTOLOCHIACEÆ.

LXXI. EMPETRACEÆ.

LXXII. EUPHORBIACEÆ.

LXXIII. CERATOPHYLLACEÆ.

LXXIV. CALLITRICHACEÆ.

LXXV. URTICACEÆ.

LXXVI. CANNABINACEÆ.

LXXVII. ULMACEÆ.

LXXVIII. AMENTIFERÆ.

## Div. V. GYMNOSPERMÆ.

LXXIX. CONIFERÆ.

Class II. *Monocotyledones.*

## Division I. DICTYOGENÆ.

LXXX. TRILLIACEÆ.

LXXXI. DIOSCOREACEÆ.

## Division II. FLORIDÆ.

LXXXII. HYDROCHARIDACEÆ.

LXXXIII. ORCHIDACEÆ.

LXXXIV. IRIDACEÆ.

LXXXV. AMARYLLIDACEÆ.

LXXXVI. ALISMACEÆ.

LXXXVII. ASPARAGACEÆ.

LXXXVIII. LILIACEÆ.

LXXXIX. MELANTHACEÆ.

XC. JUNCACEÆ.

XCI. ERIOCAULACEÆ.

XCII. TYPHACEÆ.

XCIII. ARACEÆ.

XCIV. LEMNACEÆ.

XCV. POTAMOGETONACEÆ.

XCVI. NAIADACEÆ.

## Division III. GLUMIFERÆ.

XCVII. CYPERACEÆ.

XCVIII. GRAMINEÆ.

Class III. *Cryptogameæ.*

XCIX. EQUISETACEÆ.

C. FILICES.

CI. MARSILEACEÆ.

CII. LYCOPODIACEÆ.

CIII. CHARACEÆ.

ABBREVIATIONS.

*In the descriptions.*

<i>anth.</i> . . . .	anther.	<i>pan.</i> . . . .	panicle.
<i>cal.</i> . . . .	calyx.	<i>ped.</i> . . . .	peduncle.
<i>caps.</i> . . . .	capsule.	<i>per.</i> . . . .	perianth.
<i>carp.</i> . . . .	carpel.	<i>pet.</i> . . . .	petal.
<i>cor.</i> . . . .	corolla.	<i>phyll.</i> . . . .	phyllary.
<i>fl.</i> . . . .	flower.	<i>segm.</i> . . . .	segment.
<i>fr.</i> . . . .	fruit.	<i>sep.</i> . . . .	sepal.
<i>ft.</i> . . . .	feet.	<i>st.</i> . . . .	stem.
<i>gl.</i> . . . .	glume.	<i>stam.</i> . . . .	stamen.
<i>in.</i> . . . .	inch.	<i>stig.</i> . . . .	stigma.
<i>interm.</i> . . . .	intermediate.	<i>stip.</i> . . . .	stipule.
<i>inv.</i> . . . .	involucre.	<i>t.</i> . . . .	plate.
<i>L., l., ls.</i> . . . .	leaves.	<i>term.</i> . . . .	terminal.
<i>lt., lts.</i> . . . .	leaflets.	<i>Tr.</i> . . . .	tribe.
<i>nect.</i> . . . .	nectary.	<i>var.</i> . . . .	variety.

*Books.*

<i>A.N.H.</i> . . . .	Annals of Natural History.
<i>Curt.</i> . . . .	Curtis's Flora Londinensis.
<i>E.B.</i> . . . .	English Botany.
<i>E.B.S.</i> . . . .	Supplement to E. B.
<i>J. of B.</i> . . . .	Journal of Botany.
<i>Sy. E.B.</i> . . . .	Engl. Botany, Syme's Edition.
<i>P.</i> . . . .	Parnell's Grasses of Britain.
<i>Phytol.</i> . . . .	The Phytologist.
<i>R.</i> . . . .	Reichenbach's Icones Floræ Germanicæ.
<i>R.I.</i> . . . .	Reichenbach's Iconographia Botanica.
<i>St.</i> . . . .	Sturm's Deutschlands Flora.

*Duration and Native Country.*

A. . . .	Annual.
B. . . .	Biennial.
P. . . .	Perennial.
Sh. . . .	Shrub.
T. . . .	Tree.
E. . . .	England.
S. . . .	Scotland.
I. . . .	Ireland.
† . . . .	Possibly introduced, but now appearing like a true native.
‡ . . . .	Probably introduced, but admitting of some slight doubt on the subject.
* . . . .	Certainly naturalized.
! . . . .	After the name of a plant shows that an authentic specimen has been seen.

I. II. III. &c. represent the months of flowering, viz. Jan., Feb., March, &c.; but they differ so much in different parts of the kingdom that only an approximation to the true time can be given.

When the Initial letter of the generic name is prefixed to that of a variety, it is intended to show that the author quoted considered it to be a species.

## ADDITIONS AND CORRECTIONS.

- Page  
 89, line 24, for *peduncles* read *pedicels*.  
 152, insert 24\*. SELINUM. Cal. inconspicuous. *Pet. oblong, notched*, with an incurved lobe. Carp. with 3 dorsal elevated thin ridges, and 2 marginal dilated rather broad wings. *Interstices with solitary stripes. Seed adhering to the carpel.*  
 163, insert 24\*. SELINUM Linn.  
 1. *S. Carvifolia* (L.); *st. angular furrowed*, rays of umbel scabrous.—*J. of B.* xx. 129. t. 229. *R.* xxi. 101.—L. 2—3-pinnate; lts. with linear-lanceolate mucronate serrulate lobes. No general invol., partial of many linear acuminate scales.—Broughton wood, Lincolnshire. Between Fordham and Chippenham, Cambridgeshire. P. VII. VIII. E.  
 188, l. 20, dele *hollow*.  
 200, l. 33, insert *J. of B.* xx. t. 226.  
 1. 36, insert South part of Mickel Fell range, Yorkshire. *Mr. J. Backhouse!*  
 203, l. 17, for 708. r. 704.  
 240, l. 34, add *Caps. included*.  
 241, l. 18,  $\beta$ . *capitata* (Koch); pan. compact, plant dwarf. *E. latifolia* E. B. S. 2719.  
 241, l. 34, add *Ripe caps.  $\frac{1}{2}$  protruded, unripe slightly so.*  
 1. 41, add *Cor.-tube does not lengthen after flowering*.  
 285, l. 28, for 2946 r. 2964.  
 303, l. 34, for *nearly r. mealy*.  
 310, l. 19, insert  $\beta$ . before *Sands*.  
 345, l. 34, for *latter r. lateral*.  
 351, l. 26, insert *Ringwood Chase near Ludlow. Misses Peele and Lloyd*.  
 396, l. 3, for 2 r. 3.  
 411, l. 12, for *only r. many*.  
 413, l. 2, for *uase r. base*.  
 421, insert before sp. 2.  
 1\*. *S. Townsendii* (Groves); l. jointed to their sheaths falling short of spike, spikelets 4—9, *rachis produced beyond the spikelets* and flexuose, outer glumes slightly downy.—*J. of B.* xx. l. t. 225.—St.  $1\frac{1}{2}$  to 4 ft. high. L. broadest at the base. Spikelets rather spreading.—Forms dense patches on mud-flats at Hythe, Hants.



Page

421, l. 5, for *spikes many* r. *spikelets many*.l. 9, for *pressed close together* r. *loosely pressed together*.

427, insert in place of sp. 3.

3. *A. nigra* (With.); *pan.-branches* constantly erect-patent simple below rigid strongly hispid, gl. nearly equal strongly toothed in upper half of keel, *ligule* long truncate.—*J. of B.* xx. t. 227.—Soboliferous. St. erect. Sheaths rather rough. Ped. toothed. Fl. awnless (?). Ripe anthers pale,  $\frac{1}{4}$  as broad as long.—Borders of fields.  
P. VII. E. S.

3\*. *A. vulgaris* (With.); *pan.-branches* constantly patent branched below, gl. nearly equal slightly toothed near top of keel, *ligules* short truncate.—*E. B.* 1671.—St. long, ascending or decumbent below and rooting at the knots, sometimes with very long prostrate stoles. Sheaths smooth. Pedicels softly hispid. Fl. rarely awned. Ripe anth. pale,  $\frac{1}{2}$  as broad as long.—Rather dry places.  
P. VII. E. S. I.

436, l. 9, add after 18:—*G. pedicellata* (Town.) “is nearer to *plicata* than *fluitans* by its rough furrowed sheaths, lower pale 3-toothed and never exceeding the upper.”—And *G. declinata* (Bréb., Town.) is a dwarf plant with smooth sheaths, upper exceeding the obtuse-angled 3-toothed lower pale, anth. twice as long as broad and purple. It also is near *plicata*.

441. Add to *B. asper*—branches 2 together with a semilunar strongly ciliate much decurrent scale at their base, upper pale glabrous, lower pilose on the veins below, anth. violet.—*B. Benekenii* (Syme); *pan.* slightly drooping with rather short suberect branches 3—6 together with a semilunar very blunt not ciliate subdecurrent scale at their base, upper gl. ciliate to the top, lower pale pilose throughout.

467, add 3\*. *L. complanatum* (L.); l. in 4 rows, lateral l. triangular compressed keeled acute decurrent, surface l. linear-lanceolate adpressed, spikes usually stalked terminal dichotomously branched, scales ovate acuminate serrulate, caps. reniform.—*J. of B.* xxi. t. 233.—St. ascending, dichotomous.—Hants, Glouc., Worc., Ross., Skye. E. S.

471, sp. 5, dele the Frensham station: and add Hants, Camb., Norf., Warw., Salop, Forfar. E. S.





M A N U A L  
OF  
B R I T I S H   B O T A N Y .

---

FLOWERING PLANTS.

Substance composed of cellular tissue, woody fibre, and spiral vessels. Epidermis with stomates. Flowers with stamens and pistils. Embryo with cotyledons.

Class I. DICOTYLEDONES.

Stems formed of bark, wood, and pith. The wood furnished with medullary rays and increasing by the addition of concentric layers externally. Leaves mostly with netted veins. Cotyledons 2 or more, opposite or whorled.—Each floral whorl composed of 5 or 4 parts.

Division I. THALAMIFLORÆ.

(Orders I.—XXI.)

Petals distinct (rarely 0), and as well as the stamens growing separately from the sepals on the top of the peduncle below the ovary (hypogynous).

Order I. RANUNCULACEÆ.

Sep. 3—6, often petal-like. Pet. 5 or more, rarely 0. Stam. usually many; anth. adnate, opening lengthwise. Carp. many, distinct, or forming a single pistil. Seeds erect or pendulous, albuminous.—Stip. 0, or adnate to petiole.

A. *Anthers extrorse.*

† Fruits (achenes) many, 1-seeded, short.

‡ *Sepals valvate in the bud.*

Tribe I. *CLEMATIDÆ*. Achenes with feathery persistent styles. Seed pendulous. Leaves opposite.

1. *CLEMATIS*. Cal. of 4 or 5 sepals. Pet. 0. Carp. not bursting, awned. Stam. and styles many. (St. woody.)

‡‡ *Sepals imbricate in the bud.*

Tr. II. *ANEMONEÆ*. Seed pendulous. Leaves radical or alternate.

2. *THALICTRUM*. Cal. of 4 or 5 sepals. Pet. 0. Carp. not bursting, without awns. Stam. and styles many.
3. *ANEMONE*. Cal. petal-like, sep. 5—9. Pet. 0. Carp. not bursting, *tipped with the persistent* sometimes feathery styles, placed upon a thickened hemispherical or conical receptacle. Stam. and styles many. (*Fl. involucrate.*)
4. *ADONIS*. Cal. of 5. sepals. Pet. 5—10, without a honey-bearing pore. Carp. not bursting, *without awns*. Stam. and styles many.

Tr. III. *RANUNCULÆ*. Seed erect (except in *Myosurus*). Pet. with a honey-bearing pore at their base.

5. *MYOSURUS*. Cal. of 5 sepals, prolonged into a spur at the base. Pet. 5, with a filiform tubular claw. Stam. 5. Styles many. Carp. not bursting, *closely imbricate upon a long filiform receptacle*. Seed pendulous.
6. *RANUNCULUS*. Cal. of 5, rarely 3, sepals. Pet. 5, rarely many, with a honey-bearing pore naked or covered by a scale. Carp. not bursting, *collected into a globular or oblong mass*. Stam. and styles many.

†† Fruits (follicles) many-seeded, bursting, long.

Tr. IV. *HELLEBOREÆ*. Stam. many. (Pet. small, often abnormal or wanting.)

\* Flowers regular.

7. *CALTHA*. Cal. of 5 petal-like deciduous sepals. Pet. 0. Follicles 5—10.

8. TROLLIUS. Cal. of 5 or many petal-like deciduous sepals. *Pet. small, linear, flat, clawed.* Follicles many, sessile.
- [9. ERANTHIS. Cal. of 5—8 petal-like deciduous sepals. *Pet. small, tubular, with a long claw, 2-lipped; inner lip very short.* Follicles many, *stalked.*]
10. HELLEBORUS. Cal. of 5 petal-like persistent sepals. *Pet. small, tubular, 2-lipped, clawed.* Follicles 3—10, *sessile.*
11. AQUILEGIA. Cal. of 5 petal-like deciduous sepals. *Pet. 5, funnel-shaped, with a long horn-like spur.* Follicles 5.

\*\* Flowers irregular.

12. DELPHINIUM. Cal. of 5 petal-like deciduous sepals, *upper sep. with a long spur at its base.* *Pet. 4; 2 upper ones with spurs included in the spurred sepal, or all combined into one spurred petal.* Follicles 1, 3, or 5.
13. ACONITUM. Cal. of 5 petal-like sepals, *upper sep. helmet-shaped. Two upper pet. tubular, on long stalks, concealed in the helmet-shaped sepal.* Follicles 3—5.

B. *Anthers introrse.* (*Stam. arising from a glandular disk.*)

Tr. V. PÆONIÆ or spurious Ranunculacæ.

14. ACTÆA. Cal. of 4 petal-like deciduous sepals. *Pet. 4, very small.* *Carp. 1, baccate, not bursting, many-seeded.*
- [15. PÆONIA. Cal. of 5 persistent sepals. *Pet. 5 or more.* *Follicles 2—5, many-seeded, bursting inwards.*]

### Tribe I. *Clematideæ.*

#### 1. CLEM'ATIS *Linn.* Traveller's Joy.

1. *C. Vital'ba* (L.); st. climbing, l. pinnate, leaflets ovate acuminate entire coarsely serrate or incise-lobate rounded or cordate below, petioles twining, sep. oblong downy on both sides, fr. with long feathery awns.—*E. B.* 612. *R.* iv. 64.—St. woody, angular, branched, very long. Petioles acting as tendrils.—Hedges and thickets on a calcareous soil. Sh. VI. E.

### Tribe II. *Anemoneæ.*

#### 2. THALIC'TRUM *Linn.* Meadow-Rue.

1. *T. alpinum* (L.); st. perfectly simple and nearly leafless, raceme terminal simple, fruitstalks reflexed, carp. shortly stalked

tipped with the hooked style.—*E. B.* 262. *R.* iii. 26.—St. 3—6 in. high, quite smooth. L. mostly radical, upon long stalks, twice ternate.—Higher parts of mountains. P. VI. VII. E. S. I.

2. *T. minus* (L.); st. striate branched *leafless* but sheathed at the base, stip. with spreading auricles, l. 2—3-pinnate, lts. ternate 3-cleft glaucous, petioles with angular ascending branches, fl. drooping, anth. apiculate, carp. fusiform 8 ribbed subcompressed ventricose below externally.—St. 1—1½ ft. high, usually solid; sheaths at its base rather lax. Fl. greenish yellow (as are those of Sp. 3 and 4).—*a. T. dunense* (Dum.); fr.-branches often horizontal or declining, pan. usually broad and short.—*β. T. montanum* (Wallr.); fr.-branches erect-patent, pan. much narrower than in var. *a.*—Sand-hills. *β.* Stony pastures. P. VI. VII. E. S. I.

3. *T. május* (Sm.); st. leafy to the base branched, l. 2—3-pinnate, lts. 3—5-cleft, fl. drooping, anth. apiculate, pan. with patent or reclinate branches.—*a. T. collinum* (Wallr.); petioles with divaricate branches, stip. with reflexed auricles, carp. narrowly elliptical. St. often 3—4 feet high, solid, striate. (My *T. saxatile* appears to be a smaller form. *T. flexuosum* R. not Bernh.)—*β. T. Kochii* (Fr.); petioles with patent branches, stip. with horizontal auricles, carp. ovoid. *E. B.* 611. St. often 4 ft. high, hollow, striate only below the joinings.—*a.* Damp bushy and stony places. *β.* Lake District. P. VII. VIII. E. S. I.

4. *T. flárum* (L.); st. erect furrowed, l. bipinnate, lts. broadly obovate or wedgeshaped trifid, panicle compact corymbose, fl. erect, anth. not apiculate, carp. ovoid.—L. rather paler beneath. Root-stock creeping.—*a. T. sphærocarpum* (Lej.); pan. rather close, carp. ovoid. *R.* iii. 4639.—*β. T. riparium* (Jord.); pan. usually lax with ascending branches, carp. ovoid.—*γ. T. Morisonii* (Gm.); pan. interrupted with erect-patent branches, carp. elliptic. *R.* iii. 4640.—In wet fields. P. VII. VIII. Common Meadow-Rue. E. S. I.

### 3. ANEMÓNE Linn.

1. *A. Pulsatilla* (L.); fl. solitary erect, involucre sessile in deep linear segments, l. doubly pinnate, leaflets pinnatifid with linear lobes, carp. with feathery tails.—*E. B.* 51.—Fl. bell-shaped, violet-purple, externally silky; stalk 5—8 in. high. Inv. silky, close to the flower, but distant from the fruit.—Open calcareous pastures, rare. P. IV. V. Pasque flower. E.

2. *A. nemorósa* (L.); fl. solitary, sep. 6 oblong spreading, inv. of 3 ternate or quinate stalked leaves with lobed and cut leaflets, l. similar, carp. pubescent keeled not tailed.—*E. B.* 355. *R.* iv. 47.—Fl. white or purplish; stalk 4—8 in. high. Beak

about as long as the carp. Rootstock horizontal. Sep. glabrous on both sides.—Groves and thickets, common. P. III. —V. *Wind-flower*. E. S. I.

[*A. apennina* (L.) ; fl. solitary, *sep. many lanceolate*, involucre of 3 ternate stalked deeply cut leaves, l. similar, *carp. pointed not tailed*.—*E. B.* 1062. *R.* iv. 47.—Fl. bright blue. Rootstock tuberous and roundish.—Scarcely naturalized. P. IV.] E.

[*A. ranunculoïdes* (L.) ; fl. solitary or in pairs, *sep. 5 elliptic*, involucre of 3 nearly sessile ternate deeply cut leaves, l. similar often quinate, *carp. pointed downy not tailed*.—*E. B.* 1484. *R.* iv. 47.—*Fl. bright yellow*. Sep. externally pubescent. Rootstock horizontal.—Not native. P. IV.] E.

#### 4. ADO'NIS Linn. Pheasant's Eye.

† 1. *A. autumnalis* (L.) ; cal. glabrous patent, pet. connivent, carp. without teeth collected into an ovate head and tipped with a straight beak.—*E. B.* 308. *R.* iii. 24.—Pet. scarlet, black at the base, scarcely exceeding the sepals. L. triply and copiously pinnatifid, segments linear. St. about 8 in. high.—Corn-fields, rare. A. VII. E.

### Tribe III. *Ranunculeæ*.

#### 5. MYOSU'RUS Linn. Mousetail.

1. *M. minimus* (L.).—*E. B.* 435. *R.* iii. 1.—Scapes many, single-flowered, 2—5 in. high. L. linear, fleshy. Receptacle becoming very long, with many oblong carpels. Seed attached to the upper part of the carp. and pendulous, the radicle pointing upwards.—In damp places in fields. A. V. VI. E.

#### 6. RANUN'CULUS Linn. Crowfoot.

A. *Fr.-st. arching ; carp. transversely wrinkled laterally attached, pet. white (with a yellow claw in all our plants), nectary naked.*  
BATRACHIUM Fries. Water Crowfoot.<sup>1</sup>

\* Submersed leaves twice or thrice 3-furcate with filiform segments spreading in the segment of a sphere, rarely wanting. Receptacle hispid.—(The submersed leaves become stiff when old. The younger ones should be examined.)

<sup>1</sup> See Mr. Hiern's paper on *Batrachium* in *J. of B.* ix. 43.

1. *R. trichophyllus* (Chaix); *submersed l.* (blackish green) closely trifurcate *not collapsing* into a pencil (tassel-like) when taken from the water, segments short rigid, ped. not tapering equalling or slightly exceeding the l., fl. small, *pet.* obovate 5—7-veined not contiguous *evanescent*, stig. oblong, receptacle globular, *carp.*  $\frac{1}{2}$ -obovate laterally apiculate *compressed*.—*E. B. S.* 2968. *St.* 67. 11.—*St.* not rising out of the water. Upper l. sessile, all dense. Floating l. very rare (when it is *R. Godronii* Gren. ! and also *R. radians* Rev. !) *Stip.*  $\frac{1}{2}$ -adnate, large, round, auricled. Fr.-ped. short, thick, arching, most curved near their base. *Buds globose*. Fl. starlike. *Stam.* exceeding the pistils. *Stig.* short but lengthening. Receptacle as thick as peduncle. *Carp.* a little narrowed at the end.—*R. confervoides* from Rescobie appears to be a small wholly submersed form of this species. *J. of B.* xviii. 344.—How is *R. aspergillifolius* (Dum.) to be separated from this?—Ponds and ditches. P. V. VI. *Water Fennel.* E. I.

2. *R. Drouet'ii* (F. Schultz); *submersed l.* (light green) rather loosely trifurcate, tassel-like segments flaccid, floating l. (rare) tripartite with subsessile or stalked wedgeshaped bifid segm., ped. not tapering about equalling the l., fl. small, *pet.* obovate 5—7-veined not contiguous *evanescent*, stig. oblong, receptacle oblong, *carp.*  $\frac{1}{2}$ -obovate sublaterally apiculate *inflated at the end*.—*E. B. S.* 2967.—Bright green. *St.* not rising out of the water. Upper l. nearly or quite sessile. Floating l. rare, *evanescent*; lateral segm. stalked and in a different plane from the usually sessile deflexed middle segm. *Stip.* much adnate, large, auricled. Fr.-ped. short, slender, bent at the base, nearly straight above. *Buds oblong*. Fl. starlike. *Stam.* exceeding the pistils, *few*. Fr.-receptacle as thick as peduncle. *Carp.* with the edge flattish at the end, base of style small often subcentral.—Ponds and ditches. P. V. VI. E.

3. *R. heterophyllus* (Fries); *submersed l.* loosely trifurcate, segments long tassel-like, floating l. subpeltate tripartite with sessile or stalked wedgeshaped 3—5-lobed segm., ped. not tapering but narrowed scarcely exceeding the l., fl. large, *pet.* broadly cuneate-obovate 7—9-veined not contiguous *persistent*, stig. oblong, *receptacle conical*, *carp.*  $\frac{1}{2}$ -obovate laterally pointed.—*R. aquatilis* Sm. *E. B.* 101.—*St.* not rising out of the water. Fr.-ped. slender, curved near their base, nearly straight above. Floating l. nearly circular. *Stip.* much adnate. *Buds slightly depressed* and rather pentagonal. Fl. becoming starlike. *Stam.* many, exceeding pistils. Style hooked. *Carp.* blunt, inner edge straight.— $\beta$ . *submersus*, floating l. 0, submersed l. more dense and often rather less flaccid, fl. large, *stam.* many.—Ponds and streams. P. V. VI. E. I.

4. *R. confusus* (Godr.); *submersed l.* loosely trifurcate not tassel-like, segments long rather rigid, *floating l.* long-stalked subpeltate *subtripartite with sessile obovate 3—5-lobed segments*, ped. slender gradually tapering exceeding the l., fl. large, pet. cuneate-obovate 7—9-veined not contiguous persistent, stig. tongueshaped, receptacle ovoid-conic, *carp.*  $\frac{1}{2}$ -ovate compressed and narrowed upwards.—*St.* 82. 2.—Floating l. semicircular, flat, nearly tripartite, outer base of lateral segments rounded. Stip. oblong, much adnate. Buds globular. Fl. starlike. Stam. many, exceeding pistils. Style recurved. Is *R. salsuginosus* (Hiern) when floating l. are wanting.—Ponds and ditches, especially near the sea; often in brackish water. P. VI.—IX.  
E.

5. [*R. triphyll'los* (Wallr.); *submersed l.* loosely trifurcate not tassel-like, segm. long rather rigid, *floating l.* long-stalked *tripartite with subsessile wedgeshaped 3—5-lobed segments*, ped. slender tapering exceeding the l., fl. moderately large, pet. cuneate-obovate 5—7-veined contiguous persistent, stig. . . . ., receptacle spherical, *carp.*  $\frac{1}{2}$ -ovate inflated.—Floating l. nearly or quite tripartite, acute, or blunt, or with deep linear lobes at the end.—Ponds and ditches. Guernsey. P. V.—VII.]

6. *R. Baudóttii* (Godr.); *submersed l.* closely trifurcate, segments rather rigid not tassel-like, *floating l.* long-stalked tripartite with sessile or stalked *wedgeshaped 3—4-lobed segments*, ped. thick narrowed at the top exceeding l., pet. 7-veined not contiguous persistent, *stam. not exceeding pistils*, stigma tongue-shaped, receptacle long-conic, *carp.*  $\frac{1}{2}$ -obovate inflated at the end.—*E. B. S.* 2966.—Floating l. nearly or quite tripartite, the base of all the segments wedgeshaped; or often of many linear blunt segments. Stip. much adnate. Buds globular but a little flattened at the top. Fl. starlike. Stam. 15—20. Style straight, beaklike and persistent below, recurved above. Receptacle thicker than ped., very tall. *Carp.* very many.—*R. marinus* (Fr.) is a form which wants the floating leaves.—Slightly brackish water or near the sea. P. V.—VIII.  
E. S. I.

7. *R. floribun'dus* (Bab.); *submersed l.* closely trifurcate, segments rather rigid divaricate not tassel-like, *floating l.* long-stalked subpeltate  $\frac{1}{2}$ -trifid or tripartite with obovate 3—5-lobed segments, *ped. not tapering* scarcely exceeding the l., fl. large, *pet.* obovate-cuneate 9—many-veined not contiguous persistent, *stam.* many exceeding pistils, *stig. tongueshaped*, receptacle spherical, *carp.*  $\frac{1}{2}$ -obovate very blunt.—*A. N. H.* ser. 2. xvi. 397. *E. B. S.* 2969.—Floating l. convex, divided more than

halfway down, more than semicircular; outer base of lateral segments much rounded; rarely with stalked segments. Stip. broad, with a free rounded end. Fl. starlike. Stam. 20—30. Style recurved. Receptacle as thick as peduncle. Inner edge of carp. nearly straight.—Ponds. P. V.—IX. E.

8. *R. penicillatus* (Hiern); submersed l. loosely trifurcate tassel- or whip-shaped subsessile, segments very long, floating l. long-stalked subpeltate  $\frac{1}{2}$ -trifid or tripartite with, obovate segments, each having two or three notches, ped. very long exceeding the leaves, fl. large, pet. broad becoming obovate-cuneate 9-veined contiguous persistent, stam. many exceeding the pistils, stigma . . . , receptacle spherical, carp.  $\frac{1}{2}$ -obovate very blunt.—*B. penicillatus* (Dum.). *R. pseudo-fruitans* (ed. 6).—St. wholly submersed. Floating l. semicircular or broader than long, outer base of lateral segments much rounded. Often there are no floating leaves, when it seems to be the *B. aquatile*,  $\beta$ . *rivulare* (Schur), *R. pseudo-fruitans* (Hiern). Submersed l. often 3—4 in. long with the segm. lying close together almost as in *R. fruitans*, flaccid and whip-shaped. In water, especially streams. P. V.—VIII. E. I.

9. *R. peltatus* (Fr.); submersed l. loosely trifurcate, segments rather rigid divaricate not tassel-like, floating l. long-stalked subpeltate nearly half 3—5-fid with obovate segments having 2 or 3 notches, ped. tapering exceeding the leaves, fl. large, pet. round becoming obovate-cuneate 9-veined contiguous persistent, stam. many exceeding pistils, stigma clubshaped, receptacle ovoid, carp.  $\frac{1}{2}$ -obovate very blunt.—*E. B. S.* 2965. *St.* 67. 7. *B. truncatum* Dum.—Floating l.  $\frac{2}{3}$ -circular, convex, outer base of lateral segments much rounded. Stip. adnate nearly throughout. *Fl. sweet-scented*, very large. Stam. about 30. Style curved. Receptacle small. Inner edge of carp. nearly straight.—*R. elongatus* (Hiern!), *B. elongatum* (Schultz) has very much longer ped. but otherwise does not seem to differ, except that the l.-segm. are rather less rigid.—In water and wet places. P. V.—IX. E. S.

10. *R. intermedius* (Knaf, Hiern); divided l. rare rather rigid not collapsing, floating or aerial, floating l. subpeltate deeply trifid with cuneate-obovate 2—4-fid lobes, pet. scarcely exceeding the cal., style subulate terminal with a slender base, carp. unequally obovate much inflated with a nearly terminal point.—*R. tripartitus* (DC.) *E. B. S.* 2946.—St. usually sub-erect, aerial. L.  $\frac{2}{3}$ -circular, lateral lobes with 3, middle with 2—4 crenatures. Upper stip. free. Pet. pinkish. Stam. 5—10. Style deciduous. Inner edge of carpels much rounded.—Damp ground and wet ditches, rare. A. V.—VIII. E.



\*\* Submersed l. not as in Section \*.—† Receptacle hispid.

11. *R. circinatus* (Sibth.); l. all submersed and sessile trifurcate with repeatedly and closely forked *rigid segments all placed in one roundish plane* not tassel-like, ped. tapering exceeding l., fl. large, pet. obovate many-veined nearly contiguous persistent, stam. exceeding pistils, stig. cylindrical, receptacle oblong, carp.  $\frac{1}{2}$ -ovate compressed rather acute.—*E. B. S.* 2869.—St. submersed. L. sheathing, not auricled, forming a flat rigid disk. Buds flattened at the top. Stam. 15—20. Receptacle narrower than ped. Inner edge of carp. nearly straight.—Streams and ponds, but not common. P. VI.—VIII. E. S. I.

†† Receptacle not hispid.

12. *R. fluitans* (Lam.); l. all submersed about twice trifurcate with very long linear twice or thrice forked nearly parallel segments, ped. tapering, fl. large, pet. broadly obovate many-veined contiguous persistent, stam. falling short of pistils, stig. cylindrical, receptacle conical, carp. obovate inflated much rounded at the end laterally apiculate.—*E. B. S.* 2870.—St. submersed, usually very long. Petioles and stout segments of l. often very long, together a foot in length. Stip. broadly lanceolate. Buds shortly pyramidal, pentagonal. Pet. often more than five. Stam. very short, many. Inner edge of carpels slightly rounded.— $\beta$ . *R. Bachii* (Wirtg.); slender, l. short almost sessile finely divided, pet. narrowly obovate.—Rivers.  $\beta$  rare. P. VI. VII. E. S. I.

\*\*\* Usually no divided submersed leaves; receptacle not hispid.

13. *R. Lenorman'di* (Schultz); l. all roundish cordate with 3—5 rather deeply divided lobes which widen from their base, pet. exceeding cal., style terminal upon the ovate-conical ovary, carp. unequally obovate with a terminal point.—*E. B. S.* 2930. *R. cœnósus* Bab. (not Guss.).—St. floating or creeping upon mud. L. not spotted, often opposite before the plant flowers; lobes very broad at the top, mostly with 2—3 notches. Upper stip. very broad,  $\frac{1}{2}$ -adnate. Pet. narrow, obovate, 5-veined. Stam. 8—10. Style nearly central on the ovary and usually so on the carpel. Inner edge of carp. much rounded towards the top.—Shallow ponds or mud. P. VI.—VIII. E. S. I.

14. *R. hederáceus* (L.); l. roundish cordate with 3—5 shallow rounded lobes widening to their base, pet. scarcely exceeding cal., style prolonging the inner edge of the ovary, carp.

$\frac{1}{2}$ -oval or  $\frac{1}{2}$ -obovate with a lateral point.—*E. B.* 2003.—St. floating or creeping upon mud. L. usually spotted; lobes usually entire or with a central notch, often rather triangular. Stip. narrow, much adnate. Pet. narrow, 3-veined. Stam. 6—8. Style lateral upon both ovary and carpel. Inner edge of carpel nearly straight.—When floating it is *R. homiophyllus* (Ten.), *R. cœnosus* (Guss.).—Shallow ponds or mud. P. VI.—IX. E. S. I.

*B. Fr.-st. straight; carp. transversely wrinkled on the middle of each side basally attached, nectary naked.*

15. *R. scelerátus* (L.); root fibrous, lower l. stalked tripartite, segments blunt crenate, upper l. trifid linear entire or incise-dentate, calyx reflexed, *head of fr. oblong*, carp. minute.—*E. B.* 681. *R.* iii. 11.—Fl. very small, pale yellow. Nect. round, open, bordered all round, rarely not bordered above. Lower l. broad, glabrous, shining. Stem 1—2 feet high, thick.—By and in ditches and ponds. A. VI.—IX. *Celery-leaved Crow-foot.* E. S. I.

*C. Fr.-st. straight; carp. not transversely wrinkled, basally attached.*

[*R. alpestris* and *R. gramineus* were probably recorded through mistakes.]

† L. undivided, fl. yellow, nectary nearly or quite naked bordered.—L. nearly parallel-veined.

16. *R. Flam'mula* (L.); l. ovate- or linear-lanceolate nearly entire stalked, stem reclining at the base and rooting, ped. furrowed, carp. obovate minutely pitted pointed.—*E. B.* 387. *R.* iii. 10.—Stem 6—18 in. high; sometimes procumbent and rooting. L. sometimes serrate, hairy or glabrous.—Small forms are often mistaken for *R. reptans*.—In wet places. P. VI.—VIII. *Lesser Spearwort.* E. S. I.

17. *R. reptans* (L.); l. linear entire, st. procumbent filiform creeping, carp. ovate beaked.—*Sy. E. B.* 30. *St.* 82. 14.—St. rooting at every joining. Fl. very small.—Wet places. West end of Loch Leven, Kinross-shire. P. VI.—VIII. S.

[*R. ophioglossifolius* (Vill.); lower l. cordate-ovate stalked, upper l. oblong sessile amplexicaul, stem erect hollow, carp. obliquely ovate margined tubercled with a short terminal point.—*E. B. S.* 2833. *R.* iii. 21.—Glabrous. Fl. small. St. about 1 foot high, branched, many-flowered, tapering below.—St. Peter's Marsh, Jersey. A. VI.]

18. *R. Lin'gua* (L.); l. long-lanceolate acute somewhat serrate sessile amplexicaul, stem erect, ped. not furrowed, carp. margined minutely pitted with a broad swordshaped beak.—*E. B.* 100. *R.* iii. 10.—Fl. large. St. 2—3 feet high. St. and l. glabrous or with adpressed hairs. Early submersed l. ovate-oblong blunt cordate at base.—Marshes and ditches, rather rare. P. VI. VII. *Great Spearwort.* E. S. I.

†† L. undivided, fl. yellow, nectary with a scale.

19. *R. Ficaria* (L.); root with fasciculate knobs, l. cordate stalked angular or crenate, st. with 1—3 l., single-flowered, sep. usually 3, carp. smooth blunt.—*E. B.* 584. *R.* iii. 1.—Fl. about 1 in. across; pet. usually 8; rarely apetalous. St. 3—8 in. long, weak, often producing axillary bulbs.—*a. divergens*; lobes of lowest l. separate at base, lowest sheaths narrow.—*β. incumbens*; lobes of lowest l. overlapping at the base, lowest sheaths broad clasping.—Damp rather shady places, common. P. IV. V. *Pilewort.* E. S. I.

††† L. divided, fl. yellow.

‡ Nectary without a scale, carp. smooth.

20. *R. auricomus* (L.); root fibrous, radical l. reniform 3—7-partite with crenate or cut lobes stalked, stem-l. sessile digitate, with linear or lanceolate more or less toothed segments, peduncles round, calyx pubescent, carp. downy ventricose, beak slender hooked.—*E. B.* 624. *R.* iii. 12.—Pet. often wanting. Sep. yellow. Receptacle covered with cylindrical tubercles upon which the carp. are seated. St. about 1 foot high.—Woods and thickets, common. P. IV. V. *Goldilocks.* E. S. I.

‡‡ Nectary with a scale, carp. smooth.

21. *R. ácris* (L.); st. not bulblike, root fibrous, radical l. palmately tripartite, segments trifid and deeply cut, uppermost stem-l. tripartite with linear segments, ped. terete, calyx pubescent erect-patent, carp. oval glabrous margined, beak short marginal recurved, receptacle glabrous.—*E. B.* 652. *R.* iii. 17.—Hairy. No stoles. St. 2—3 feet high. Beak about  $\frac{1}{4}$  the length of the carpel.—Sometimes dwarf and 1-flowered on mountains.—The variations *R. vulgatus* (Jord.) and *R. tomophyllus* (Jord.) have been found. *J. of B.* viii. 257, x. 238. The former has usually an oblique or horizontal rhizome, the latter a præmorse rootstock.—Meadows and pastures, common; mountains. P. VI. VII. *Upright Crowfoot.* E. S. I.

22. *R. répens* (L.); l. with three-lobed segments, lobes 3-fid and cut, *ped. furrowed*, *calyx* pubescent *erect-patent*, carp. oval glabrous margined minutely pitted, beak longish slightly curved, *receptacle hairy*.—*E. B.* 516.—Root fibrous. Stoles strong, leafy. Primary stem usually erect, 10—12 in. high.—Meadows and pastures, common. P. V.—VIII. *Creeping Crowfoot*. E. S. I.

23. *R. bulbósus* (L.); *stem bulblike* at the base, radical l. with 3 segments each tripartite trifid and cut, *ped. glabrous furrowed*, *calyx* hairy *reflexed*, carp. round margined smooth, beak short, *receptacle hairy*.—*E. B.* 515. R. iii. 20.—St. about 1 foot high. Upper l. cut into narrow segments.—Meadows and pastures. P. V. *Bulbous Crowfoot*. E. S. I.

††† Nectary with a scale, carp. rugose or tubercular.

[*R. chærophyllus* (L.); *root of short ovoid knobs* and fibres, *stoles very slender* with minute scales and ending in a young plant, rt.-l. with 3 segments each 3—7-parted or 3-fid, st.-l. 1—2 with linear segments, *ped. terete*, *cal. hairy spreading*, carp. many roundish, beak acute, *receptacle oblong glabrous*.—*J. of B.* x. t. 125.—St. 6—12 in. high, and, as well as the leaves, hairy.—Dry places, St. Aubin's, Jersey. P. V.]

24. *R. hirsútus* (Curt.); root fibrous, radical l. with 3 stalked trifid and cut leaflets, peduncles furrowed, *calyx reflexed*, carp. round margined *with a series of tubercles near the margin*, beak short curved, *receptacle hairy*.—*E. B.* 1504. R. iii. 23. *R. Philonotis* (Ehrh.) Koch.—St. 4—18 in. high; the smaller specimens are *R. parvulus* L. Upper l. in narrow acute segments. Fl. pale yellow.—Waste land and corn-fields, rare. A. VI.—X. E. S.

25. *R. parviflórus* (L.); root fibrous, *stems spreading*, l. roundish-cordate 3—5-lobed cut, upper l. oblong undivided or 3-lobed, *calyx* at first erect afterwards reflexed, carp. orbicular muricate.—*E. B.* 120. R. iii. 22.—Peduncles opposite the leaves. Pet. narrow.—Corn-fields and dry banks, rare. A. V. VI. E. I.

†26. *R. arven'sis* (L.); root fibrous, radical l. 3-cleft dentate, stem-l. once or twice ternate with linear-lanceolate segments, *calyx erect-patent*, carp. margined *beaked and spinous*.—*E. B.* 135. R. iii. 21.—St. 6—18 in. high. Fl. pale yellow.—Corn-fields. A. VI. *Corn Crowfoot*. E. S. I.

Tribe IV. *Helleboreæ*.

## 7. CAL'THA Linn. Marsh Marigold.

1. *C. palustris* (L.); *st. ascending, l. cordate rounded crenate*.—*E. B.* 506. *R.* iv. 101.—About a foot high. Fl. large. Sep. bright yellow.— $\alpha$ ; sep. roundish-ovate contiguous, carp. spreading their beak very short.— $\beta$ . *C. Guerangerii* (Bor.); sep. oblong-ovate not contiguous, carp. spreading their beak long.— $\gamma$ . *minor*; *st. decumbent mostly 1-flowered, fl. small, sep. not contiguous, carp. erect their beak very short*.—Marshy places, common.  $\gamma$ . Mountains. P. III.—V. E. S. I.

2. ? *C. radicans* (Forst.); *st. creeping, l. triangular serrate-crenate small, sep. not contiguous narrow*.—*E. B.* 2175.—Base of the l. almost at right angles with petiole.—“In a ditch that runs from the farmhouse called Haltoun, on the estate of C. Gray, Esq., of Carse” (2 miles north of Forfar near Rescobie). *Mr. G. Don.* P. V. VI. S.

## 8. TROL'LIVUS Linn. Globe Flower.

1. *T. europæus* (L.); sep. 10—15 concave converging into a globe, pet. 10 about equalling the stam., l. palmately 5-parted, segments rhomboidal 3-partite incise-serrate.—*E. B.* 28. *R.* iv. 101.—Fl. bright yellow. Pet. ligulate. St. 1—2½ feet high.—Damp mountain pastures. P. VI. VII. E. S. I.

## 9. ERAN'THIS Salisb. Winter Aconite.

[\* *E. hyemalis* (Salisb.); sep. 6—8 oblong.—*Sy. E. B.* 43. *R.* iv. 101.—Radical l. on long stalks, 5—7-parted, deeply cut into linear-oblong segments. St. 4—6 in. high; invol. of 3 sessile leaflets just below the solitary yellow fl. Rhizome tuberous.—Naturalized in thickets. P. II. III.] E. S. ?

## 10. HELLEBORUS Linn. Hellebore.

1. *H. viridis* (L.); radical l. *digitate stalked, stem-l. sessile at the ramifications, st. few-flowered, calyx spreading*.—*E. B.* 200.—Veins of the l. prominent beneath. Stigma erect. St. 1 foot high, annual. Fl. greenish-yellow.—Thickets on a calcareous soil. P. III. IV. *Green Hellebore. Bear'sfoot.* E.

2. *H. foetidus* (L.); l. *pedate stalked, st. leafy many-flowered, calyx converging*.—*E. B.* 613.—L. successively contracting upwards into bracts. St. 2 feet high, perennial. Fl. globose,

drooping, greenish tipped with purple.—Thickets in chalky districts. P. III. IV. *Stinking Hellebore. Setterwort.* E.

### 11. AQUILE'GIA Linn. Columbine.

1. *A. vulgaris* (L.); spur of the pet. incurved, limb blunt falling short of the stamens, l. biternate, leaflets 3-lobed crenate.—*E. B.* 297. *R.* iv. 114.—St. 2—3 feet high, slightly leafy. Caps. hairy. Inner stam. frequently imperfect.—Woods and thickets and heaths, not common. P. V. VI. E. I.

### 12. DELPHIN'IUM Linn. Larkspur.

\*1. *D. Ajacis* (Gay not R.); st. erect with spreading branches, racemes 4—16-fl., pet. combined, ovary abruptly narrowed into a subulate style, follicle downy obliquely acuminate, seeds with transverse contiguous wavy ridges.—*D. Consolida* Sm. *Sy. E. B.* 47.—St. loosely and sparingly branched, about a foot high. Fl. of a vivid and permanent blue, rarely red pink or white. L. deeply multifid. Style equal to about  $\frac{1}{6}$  of carpel.—Sandy or chalky corn-fields. A. VI. VII. E.

[*D. Consolida* (L.); st. erect with patent branches, racemes few-flowered, pet. combined, ovary narrowed into style, follicle glabrous truncate short, seeds with transverse interrupted ridges.—*R.* iv. 116.—Much like *D. Ajacis*. Style lateral.—Found once in Jersey and in Cornwall. A. VI. VII.]

### 13. ACONI'TUM Linn. Monk's-hood.

†1. *A. Napellus* (L.); pet. horizontal upon curved stalks, spurs bent down, fl. racemose, young carpels diverging.—*E. B.* S. 2730. *R.* iv. 92.—Rootstock black, of 2 oblong knobs. St. 1—2 ft. high. Fl. purple. Filaments slightly hairy, with cuspidate wings. Pet. inflated above; lip broad. Helmet open, hemispherical. Pedicels erect, downy.—Banks of rivers and brooks, rare. P. VI. VII. E.

## Tribe V. *Pæoniæ*.

### 14. ACTÆ'A Linn. Bane-berry.

1. *A. spicata* (L.); raceme simple elongate, pet. as long as the stamens, berries oval.—*E. B.* 918. *R.* iv. 121.—L. stalked, biternate; lts. ovate, trifid, deeply cut. St. 1—2 feet high. Fl. white.—Mountainous limestone tracts in the North. P. V. E. S.

15. PÆO'NIA *Linn.* Pæony.

[\* *P. Coral'ina* (Retz.); l. biternate, leaflets ovate entire glaucous beneath, caps. downy recurved from the base.—*E. B.* 1513. *R.* iv. 128.—Roots fleshy, knobbed. Herb 2 feet high. Fl. large, crimson with yellow anthers.—On the Steep Holmes Island in the Severn. P. V. VI.] E.

## Order II. BERBERIDACEÆ.

Sep. 3 or 4 or 6 in a double row. Pet. the same or double that number. Stam. opposite to the petals. Anth. opening by valves attached at the top and turning upwards. Carpel 1, 1-celled; seeds attached to the bottom or on a lateral 'placenta, albuminous.—Stipules usually wanting.

1. BERBERIS. Sep. 6, deciduous. Pet. 6, each with 2 glands at the base within. Stam. 6. Berry 2—3-seeded.

[2. EPIMEDIUM. Sep. 4, deciduous. Pet. 4. Nectaries 4, cupshaped. Stam. 4. Caps. podlike, many-seeded.]

1. BER'BERIS *Linn.* Barberry.

1. *B. vulgaris* (L.); spines 3-parted, l. obovate ciliate-serrate, racemes pendulous many-flowered, petals entire.—*E. B.* 49. *R.* iii. f. 4486.—Height 6—8 feet. Fl. yellow. Berries red, oblong, slightly curved. Filaments curiously elastic.—Hedges and thickets. Sh. V. VI. E. S. P I.

2. EPIME'DIUM *Linn.* Barrenwort.

[*E. alpinum* (L.); rhizome producing leaves and stems, stem-l. twice ternate.—*E. B.* 438. *R.* iii. f. 4485.—Lts. ovate-heart-shaped, serrate. Pan. shorter than the l. and appearing to grow from the petiole. Fl. red with yellow nectaries.—Subalpine woods. Not a native. P. V.] E. S.

## Order III. NYMPHÆACEÆ.

Sep. 4—6. Petals many, seated with the many stamens upon a fleshy disk more or less completely surrounding the ovary. Stigma peltate. Fruit many-celled; seeds many, in a gelatinous aril. Embryo in a bag on the outside of the base of the albumen.—St. prostrate, submersed. L. floating, falsely peltate.

1. NYMPHÆA. Cal. of 4 sepals. Pet. many, seated together with the stamens upon a fleshy disk enveloping the germen and passing gradually into them. Berry many-celled, many-seeded. Stigma sessile, of many rays.
2. NUPHAR. Cal. of 5 sepals. Pet. many, seated together with the stamens upon the receptacle. Berry many-celled, many-seeded. Stigma sessile, of many rays.

1. NYMPHÆ'A Linn. White Water-Lily.

1. *N. alba* (L.); l. roundish deeply cordate entire with approximate lobes, stigma of 12—20 rays.—*E. B.* 160. *R.* vii. 67.—Fl. large, white, floating. (There is a small-flowered form.) Stig. yellow. Ovary covered with the stam. almost to its top. Caps. dissolving into pulp. Notch in the leaves with nearly parallel sides.—Slow rivers, lakes and clear ditches. P. VII.  
E. S. I.

2. NU'PHAR Sm. Yellow Water-Lily.

1. *N. lutea* (Sm.); l. oblong-cordate, sep. 5, stigma with 10—20 rays not extending to the entire margin, anthers linear.—*E. B.* 159. *R.* vii. 63.—Fl.  $2\frac{1}{2}$  in. across, yellow, smelling like brandy. Caps. bursting irregularly. Submersed l. thin, transparent, wavy, oblong-cordate.— $\beta$ . minor; fl.  $1\frac{1}{2}$  in. across, margin of stigma wavy. *Sy. E. B.* 55. *N. spenneriana* Gaud. ?—Lakes and ditches. P. VII. *Brandy-bottle.*  
E. S. I.

2. *N. pímila* (DC.); l. oblong deeply cordate with distant lobes, sep. 5, stigma with 8—10 rays extending to the margin and forming acute teeth, anth. subquadrate.—*E. B.* 2292. *R.* vii. 65.—Fl. yellow, small. Caps. furrowed upwards. Anth. not twice as long as broad. Much smaller than *N. lutea*. Submersed l. thin, transparent, wavy, reniform.—In small Highland lakes, and at Ellesmere, Salop. P. VII. VIII.  
E. S.

Order IV. PAPAVERACEÆ.

Sep. 2, deciduous. Cor. regular, of 4 petals. Stam. generally many, free. Ovary free; placentas parietal, usually projecting; seeds many. Stigma as many as the placentas, simple or lobed; the lobes of adjoining stigmas combining, thus appearing (falsely) to be opposite to the placentas. Seeds albuminous.—Stip. 0.



- \* *Fruit globular, oblong or clavate, opening by pores; stigmas radiant.*
1. PAPAVER. Pet. 4. Stam. many. *Style 0. Stigmas 4—20, radiant on a flattish disk.* Caps. opening by pores beneath the stigma. Placentas like dissepiments.
  2. MECONOPSIS. Pet. 4. Stam. many. *Style short. Stigmas 4—6, radiant, free.* Caps. obovate, opening by pores beneath the style. Placentas filiform.
- \*\* *Fruit linear, opening by valves; stigmas 2—4.*
3. ROEMERIA. Pet. 4. Stam. many. *Stigmas 2—4, sessile, radiant.* Caps. 2—4-valved, 1-celled; *placentas distinct.* Seeds not crested.
  4. GLAUCIUM. Pet. 4. Stam. many. *Stigmas 2, sessile. Caps. 2-valved; placentas connected by a spongy dissepiment.* Seeds not crested.—A maritime plant with yellow flowers.
  5. CHELIDONIUM. Pet. 4. Stam. many. *Stigmas 2. Caps. 2-valved, 1-celled; placentas distinct.* Seeds crested.

### 1. PAPA'VER Linn. Poppy.

†1. *P. Argemone* (L.); filaments dilated upwards, *caps. clavate* hispid with erect bristles, stem leafy many-flowered, l. bipinnatifid.—*E. B.* 643. *R.* iii. f. 4475.—St. usually about 1 ft. high. Fl. small. Pet. distinct, fugacious, pale red, black at the base. Bristles sometimes spreading.—In corn-fields. A. VI. VII. E. S. I.

†2. *P. hybridum* (L.); filaments dilated upwards, *caps. roundly ovoid* hispid with spreading bristles, stem leafy many-flowered, l. bipinnatifid.—*E. B.* 43. *R.* iii. f. 4476.—Sap milky. Fl. small. Pet. purplish, often with a dark spot at the base.—Sandy fields, rare. A. VI. VII. E. I.

†3. *P. Rhœas* (L.); *filaments subulate, caps. roundly obovate without bristles*, stem bristly many-flowered, l. pinnatifid cut.—*E. B.* 645. *R.* iii. f. 4479.—Fl. large. Pet. deep scarlet, often nearly black at the base. Peduncles with spreading hairs, or (*β. strigosum* Boeningh.) with adpressed hairs.—In arable fields, common. A. VI. VII. *Corn Rose.* E. S. I.

†4. *P. dubium* (L.); filaments subulate, *caps. clavate narrowing gradually* from the top throughout without bristles, *stig.-*

*disk with patent edge*, l. pinnatifid with distant broad entire bluntish lobes, sap milky.—*Curt. Lond.* ii. 104. *P. Lamottei* Bor.—St. 1—2 ft. high. Sap not turning yellow. Fl. large. Pet. transversely oval, pale red. Ped. with adpressed hairs. *Stig.-disk of nearly ripe caps. like eaves*, obscurely lobed.—Sides of fields. A. VI. VII. E. S. I.

†5. *P. Lecoc'ii* (Lamot. ?); filaments subulate, *caps. clavate-oblong* suddenly narrowed near the base without bristles, *stig.-disk folded over the edge of the caps.*, l. bipinnatifid with distant narrow entire acute lobes. *Sap becoming dark yellow* in the air.—*Sy. E. B.* 60.—St. 1—2 ft. high. Fl. large. Pet. usually distinct, obovate-wedgedshaped, red. Ped. with adpressed hairs. *Stig.-disk with bluntly triangular lobes.*—Sides of fields chiefly on a calcareous soil. A. VI. VII. E. S. I.

†6. *P. somniferum* (L.); *filaments dilated* upwards, *caps. truly globular without bristles*, l. oblong unequally toothed amplicaul.—*E. B.* 2145. *R.* iii. f. 4481.—Fl. large. Pet. bluish white with a violet spot at the base. Whole herb smooth, except sometimes a few rigid spreading bristles on the ped. and one tipping many of the teeth of the leaves.—On sandy ground near the sea, and in the Fens, rare. A. VII. E.

## 2. MECONOP'SIS *Vig.* Welsh Poppy.

1. *M. cam'brica* (*Vig.*); caps. smooth, l. stalked.—*E. B.* 66.—Caps. elliptic-oblong, beaked. St. many-flowered, about 1 foot high. L. pinnate, cut, glaucose beneath. Fl. large, yellow, on long stalks.—Damp rocky and shady places. P. VI. E. I.

## 3. ROEME'RIA *Cand.*

†1. *R. hybrida* (DC.); pod 3-valved erect with a few rigid hairs at its top.—*E. B.* 201.—Caps. linear, 2—3 in long. L. 2—3-pinnatifid, with linear nearly smooth bristle-pointed lobes. St. about 1 foot high, usually slightly hairy. Fl. violet-blue. Pet. falling before noon.—Chalky corn-fields in Cambridgeshire and Norfolk, very rare. A. V. VI. E.

## 4. GLAU'CIUM *Tourn.* Horned Poppy.

1. *G. huteum* (Scop.); st. smooth, stem-l.  $\frac{1}{2}$ -clasping sinuate, caps. minutely tubercular-asperous.—*E. B.* 8.—Glaucous. St. 1—3 feet high, stout, much branched, glabrous or slightly hairy. Root-l. stalked, lyrate, lobed and cut, hairy. Pet. large, golden

yellow. Caps. 6—12 in. long, curved.—Sandy sea-shores. B. VI.—VIII. E. S. I.

[*G. phœniceum* (Crantz); st. pilose, stem-l. pinnatifid cut, caps. hispid.—*E. B.* 1433.—Pet. scarlet with a black spot at the base.—Rarely found, on cultivated ground. A. VI. VII.] E.

### 5. CHELIDO'NIUM *Linn.* Celandine.

†1. *C. május* (L.); ped. umbellate, l. deeply pinnatifid, segments rounded and bluntly lobed.—*E. B.* 1581.—Fl. yellow, small. Caps. linear. St. 1—2 feet high. All parts full of orange juice.—[ $\beta$ . *C. laciniatum* (DC.); segments of the leaves deeply pinnatifid, lobes incise-serrate.]—Waste places and old walls.  $\beta$  a very doubtful native. P. V.—VIII. E. S. I.

### Order V. FUMARIACEÆ.

Sep. 2 or 0. Cor. irregular, of 4 parallel petals, one or both of the two outer pet. gibbous or spurred at the base. Stam. 6, in 2 bundles opposite to the outer petals; lateral stam. in each bundle 1-celled. Ovary free, 1-celled. Style filiform. Stigma with 2 or more points. Seeds albuminous.—Stipules 0.

1. CORYDALIS. Cal. of 2 sepals or wanting. Pet. 4, the upper one spurred at the base. Stam. diadelphous. *Pod 2-valved many-seeded*, compressed.
2. FUMARIA. Cal. of 2 sepals. Pet. 4, the upper one spurred at the base. Stam. diadelphous. *Fruit a nut*, indehiscent, 1-seeded.

#### 1. CORYD'ALIS *Cand.*

[*C. sol'ida* (Hook.); root tuberous solid, l. biternate cut, lowest petiole a leafless scale, bracts palmate.—*E. B.* 1471.—Lobes of the l. blunt. Fl. purplish. St. a span high.—Not a native, scarcely even naturalized. P. IV. V.] E.

\*1. *C. lítea* (DC.); root fibrous, l. triternate, bracts minute oblong cuspidate, seeds shining granular-rugose with a patent denticulate crest.—*E. B.* 588.—Lts. obovate, trifid. Bracts shorter than pedicels. Fl. yellow. St. 1 ft. high, brittle, spreading.—Naturalized on old walls. P. V.—VIII. *Yellow Fumitory*. E. S.

2. *C. claviculáta* (DC.); root fibrous, leaves pinnate, pinnæ ternate, footstalks ending in tendrils, bracts oblong acuminate.—

*E. B.* 103.—Leaflets entire, elliptic. Bracts rather longer than the pedicels. Fl. small, yellow or nearly white. St. slender, climbing, 1—4 feet long.—Bushy places in hilly districts. P. VI. VII. *White Climbing Fumitory.* E. S. I.

## 2. FUMA'RIA, *Linn.* Fumitory.

\* *Capreolata*. Lower pet. narrowing gradually and slightly from its middle upwards. Fr. not retuse.—*F. capreolata* L.—St. erect, climbing or diffuse. Fr. subcompressed, smooth. Raceme lax, short, few-flowered. Sep. as broad as cor-tube. L. flat.

1. *F. pallidiflora* (Jord.); sep. ovate toothed at least  $\frac{1}{2}$  as long as the cor., fr. blunt its neck short and narrow, fr.-st. patent or reflexed.—Sep. soon falling. Cor. large. Fr. not regularly rounded vertically but squarish; apical pits small and deep.—*a. Jordani*; fr. longer than broad, its neck narrower than the tip of ped., fr.-st. reflexed. Cor. cream-coloured, tipped with red or pink. *Sy. E. B.* 71.— $\beta$ . *F. Boræi* (Jord.); fr. broader than long, its neck very narrow, fr.-st. patent. Cor. purplish, tipped with dark purple. *Sy. E. B.* 72. *Curt.* ii. 145.—Borders of fields. A. VI.—IX. E. I.

2. *F. confusa* (Jord.); sep. ovate toothed not  $\frac{1}{2}$  as long as cor., fr. rounded at the top its neck very broad, fr.-st. patent.—*E. B. S.* 2976. *Sy. E. B.* 73.—Sep. often persistent. Cor. large, dull white or pinkish, tipped with dark purple. Fr. regularly rounded vertically; apical pits broad and shallow; neck nearly as broad as the fruit.—Borders of fields, chiefly in the West. A. VI.—IX. E. S. I.

3. *F. muralis* (Sond.); sep. ovate toothed not  $\frac{1}{2}$  as long as the cor., fr. obovate rounded at the top its neck obconic narrow, fr.-st. erect-patent.—*Sy. E. B.* 74. *Fl. Dan.* 2473.—More lax than Sp. 1 and 2, cor. smaller and nearly black at the tip. Fr. uniformly rounded at the sides and top, nearly pyriform if taken with its neck; apical pits very faint.—Borders of fields. A. VI.—IX. E. S.

\*\* *Officinales.* Lower petal spatulate.

4. *F. officinalis* (L.); sep. ovate-lanceolate acute toothed narrower than the cor.-tube, fr. obovate retuse, bracts much shorter than the fruitstalks, lts. flat.—*Curt.* i. 14.—St. erect or diffuse. Raceme long, many-flowered. Sep. broader than the pedicel,

about  $\frac{1}{3}$  the length of the corolla. Pet. rose-coloured. Fr. rugose, broader than long.—Common. A. V.—IX. E. S. I.

†5. *F. densiflora* (DC.); sep. large roundish dentate broader than the cor.-tube, fr. subglobose subapiculate, bracts longer than the fr.-stalks, lts. linear channelled.—*F. micrantha* Lag. *E. B. S.* 2876.—St. branched, diffuse. Pet. pale purple. Fr. rather longer than broad.—Fields. A. VI.—IX. E. S.

6. *F. parviflora* (Lam.); sep. minute ovate acute cut, fruit-stalk shorter than the obovate pointed fr. and equalling the bract, lts. linear channelled.—*E. B.* 590. *R.* iii. 1.—Foliage yellowish green glaucous. Lts. ascending. Fl. whitish, afterwards pale purple.—Fields in Kent, Essex, Camb. A. VI.—IX. E.

7. *F. Vaillantii* (Lois.); sep. very minute triangular, fr.-st. longer than the obovate pointed fr. and twice as long as the bract, lts. narrow flat.—*E. B. S.* 2877. *R.* iii. 1.—Foliage greyish green glaucous. Lts. spreading. Fl. purplish, afterwards whitish.—Fields in the South and East, rare. A. VI.—IX. E.

## Order VI. CRUCIFERÆ.

Sep. 4. Cor. cruciform, of 4 petals. Stam. 6, tetradynamous; 2 shorter opposite the lateral sepals. Ovary free, with marginal placentas connected by a false septum. Stigmas 2, opposite to the placentas (or, rather, alternate with them, lobed and combining). Fruit a 2-celled and 2-valved caps. or pod, the valves opposite the shorter stamens, deciduous from the placentas.—Stipules 0.

### Suborder I. SILIQUOSÆ.

Pod (silique) long, not dividing transversely, linear or linear-lanceolate, opening by 2 valves throughout; dissepiment narrow, but in the broadest diameter of the pod.

Tribe I. *ARABIDEÆ*. Cotyledons accumbent (radicle lateral), seed compressed.

\* *Stigma 2-lobed.*

1. *MATTHIOLA*. Pod round or compressed. Lobes of stig. erect gibbous or horned at the back.
2. *CHEIRANTHUS*. Pod compressed or 2-edged, with an ele-

*vated longitudinal rib upon each valve. Lobes of stig. patent.*  
Seeds in a single row in each cell.

\*\* *Stigma a disk or head.*

3. NASTURTIUM. Pod terete; *valves convex, veinless.* Seeds irregularly in 2 rows.
4. BARBAREA. Pod terete; *valves convex, with a prominent longitudinal rib.* Seeds in a single row.
5. ARABIS. Pod compressed; valves nearly flat, with a prominent longitudinal rib or with numerous longitudinal veins. *Seeds in a single or double row.*
6. CARDAMINE. Pod compressed; *valves flat, veinless.* Seeds in 1 row; seed-stalk simple, filiform.
7. DENTARIA. Pod lanceolate, compressed; valves flat, veinless. Seeds in a single row; seed-stalk *dilated, winged.*

Tr. II. *SISYMBRIÆ.* Cotyledons incumbent (radicle dorsal), seed compressed. Seeds in one row.

\* *Stigma of two closely converging erect ovate lobes.*

8. HESPERIS. Pod quadrangular or subcompressed; valves keeled, somewhat 3-veined. Seeds in a single row.

\*\* *Stigma a disk with a thickened edge.*

9. SISYMBRIUM. Pod terete or rarely 4-edged; valves convex, with 3 longitudinal veins. Seeds in a single row, smooth; seed-stalk filiform.—*S. thalianum* has a tetragonous pod with 1 strong conspicuous rib on the valves.
10. ALLIARIA. Pod terete; valves convex, with 3 longitudinal veins, the middle one prominent and strong, the 2 lateral slender and branching. *Seeds in a single row, striate; seed-stalk flattened, winged.*
11. ERYSIMUM. Pod 4-edged; *valves prominently keeled, with 1 longitudinal vein.* Seeds in a single row; seed-stalk filiform.

Tr. III. *BRASSICÆ.* Cotyledons conduplicate (longitudinally folded in the middle); radicle dorsal, within the fold.

12. BRASSICA. Cal. erect. Pod terete or angular. Seeds globose, in a single row.

13. *SINAPIS*. Cal. spreading (in the flower). Pod terete or angular. Seeds in a single row.
14. *DIPLOTAXIS*. Cal. patent. Pod compressed. *Seeds* oval or oblong, *in 2 rows*.

## Suborder II. LATISEPTÆ.

Pouch (silicle) short, not dividing transversely, opening with two valves; dissepiment in its broadest diameter.

Tr. IV. *ALYSSINEÆ*. Cotyledons accumbent.

15. *ALYSSUM*. Pouch roundish or oval, compressed. Seeds 2—4 in each cell. Filaments simple or toothed or the shorter ones with a gland or subulate process on each side at the base.
16. *DRABA*. Pouch oval or oblong, slightly convex. *Seeds many* in each cell, in 2 rows, not margined. Filaments simple.
17. *COCHLEARIA*. Pouch globose; *valves very convex*, dorsal vein prominent. Seeds many. Filaments simple.
18. *ARMORACIA*. Pouch oblong or globose; *valves very convex, no vein*. Seeds many. Filaments simple.

[Tr. V. *CAMELINEÆ*. Cotyledons incumbent.

19. *CAMELINA*. Pouch subovate; valves ventricose, with a linear prolongation at the end which is confluent with the persistent style.]

[Tr. VI. *VELLEÆ*. Cotyledons conduplicate.

20. *VELLA*. Pouch ovate, with a dilated winged flat style longer than the convex valves.]

## Suborder III. ANGUSTISEPTÆ.

Pouch (silicle) short, laterally compressed, opening with 2 boatshaped valves keeled or winged on the back; dissepiment narrow, linear or lanceolate.

Tr. VII. *THLASPIDEÆ*. Cotyledons straight.

\* *Cotyledons accumbent or rarely incumbent.*

21. *THLASPI*. Pouch roundish, notched; valves boatshaped, winged at the back. *Seeds more than 2 in each cell*. Pet. equal, filaments simple.

22. HUTCHINSIA. Pouch elliptic, entire; valves boatshaped, keeled *not winged* at the back. Seeds 2 in each cell. Pet. equal. Filaments simple. Cotyledons rarely incumbent.
23. TEESDALIA. Pouch roundish, notched; valves boatshaped, their back keeled *below narrowly winged above*. Seeds 2 in each cell. Pet. equal or two outer ones larger. Filaments with a little scale at the base of each within.
24. IBERIS. Pouch ovate or roundish, notched; valves boatshaped, winged at the back. Seeds 1 in each cell. Pet. unequal, 2 outer ones much larger. Filaments simple.

\*\* Cotyledons usually incumbent

25. LEPIDIUM. Pouch roundish or oblong, entire or notched; valves compressed, keeled or winged at the back. Seeds 1 in each cell. Pet. equal. Cotyledons rarely accumbent.
26. CAPSELLA. Pouch triangular-obcordate; valves compressed, keeled *but not winged*. Seeds many.

Tr. VIII. SUBULARIÆ. Pouch with a rather broad dissepiment. Cells many-seeded. Cotyledons incumbent, long, linear, curved back above their base, thus appearing like 4 in transverse section.

27. SUBULARIA. Pouch oval-oblong, laterally subcompressed; valves boatshaped.

Tr. IX. SENEBIERÆ. Valves not separating. Cells one-seeded. Cotyledons incumbent, long, linear, curved back above their base, as in Tr. VIII.

28. SENEBIERA. Pouch broader than long, somewhat kidney-shaped, entire at the end, or notched above and below and almost 2-lobed.

#### Suborder IV. NUCUMENTACEÆ.

Pouch (silicle) scarcely dehiscent, often 1-celled owing to the absence of the septum.

Tr. X. ISATIDÆ.

29. ISATIS. Pouch laterally compressed, 1-celled, 1-seeded; valves keeled, eventually separating. Cotyledons incumbent.



## Suborder V. LOMENTACEÆ.

Silicle or silique dividing transversely into 1-seeded cells; the true silique often barren, all the seeds being in the beak.

Tr. XI. *CAKILINEÆ*. Silicle 2-jointed.

30. *CAKILE*. Silicle angular, of two 1-seeded indehiscent joints, upper joint deciduous with an erect seed, lower persistent seedless or with a pendent seed.

31. *CRAMBE*. Silicle 2-jointed, upper joint globose with 1 seed pendent from a long curved seed-stalk springing from the bottom of the cell, lower joint barren stalklike.

Tr. XII. *RAPHANEÆ*. Silique linear or oblong, terete or moniliform.

32. *RAPHANUS*. Silique linear or oblong, tapering upwards, smooth and indehiscent, or moniliform and dividing transversely into 1-seeded cells, lowermost cell barren imperfectly 2-valved stalklike.

Suborder I. *Siliquosæ*. Tribe I. *Arabideæ*.1. *MATTHIOLA* R. Br. Stock.

†1. *M. incana* (R. Br.); st. shrubby upright branched, l. lanceolate entire hoary, pods "cylindrical without glands."—*Sy. E. B.* 1935. *R.* ii. 45.—Fl. of a full purple.—Cliffs in the Isle of Wight. P. V. VI. *Hoary Stock*. E.

2. *M. sinuata* (R. Br.); st. herbaceous diffuse, l. oblong downy, lower l. *sinuate*, pods compressed *muricated* with glands.—*E. B.* 462. *R.* ii. 45.—Fl. purple.—South and South-west coasts. B. VI.—VIII. *Sea Stock*. E. I.

2. *CHEIRANTHUS* Linn. Wallflower.

\*1. *C. Cheiri* (L.); st. shrubby, l. lanceolate acute entire with bipartite adpressed hairs, pods tetragonal.—*E. B.* 1934. *R.* ii. 45.—Fl. yellow or tinged with red.—Old walls. P. IV. V. E. S. I.

3. *NASTURTIUM* R. Br. Water-Cress.

1. *N. officinale* (R. Br.); l. pinnate, lts. ovate or oblong subcordate sinuate-dentate, pet. twice as long as calyx (*white*), pods linear.—*E. B.* 855. *R.* ii. 50.—Pods patent. When growing out of water it is slender with small leaves and is *N. microphyllum* (R.); when remarkably luxuriant, many feet in length, the stem often nearly an inch thick, and the leaves very large and resembling those of a *Sium*, it is *N. siifolium* (R.).—Running water. P. VI. VII. *Water-Cress*. E. S. I.

2. *N. sylvestre* (R. Br.); creeping, l. deeply pinnatifid, lts. oblong or lanceolate cut, uppermost l. often nearly entire, *pet. twice as long as the calyx* (yellow), pods linear.—*E. B.* 2324.—Fr.-stalks patent; pods patent or ascending, variable in length, usually as long as their stalks, sometimes shorter.—River banks and wet places. P. VI.—VIII. E. S. I.

3. *N. palustre* (DC.); root fibrous, lower l. lyrate, upper l. deeply pinnatifid, lts. oblong toothed, *pet. not longer than the calyx*, pods oblong thick.—*N. terrestre* Sm. *E. B.* 1747.—Fl. small. Fr.-stalks patent or deflexed; pods ascending, short, about as long as their stalks.—Wet places. P. VI.—IX. E. S. I.

#### 4. BARBARE'A R. Br. Rocket.

1. *B. vulgaris* (R. Br.); lower l. lyrate, upper pair of lobes equalling the breadth of the large roundish subcordate terminal lobe, uppermost l. undivided toothed, pods adpressed obliquely erect or patent with a subulate point.—*E. B.* 443. *R.* ii. 47.—*Pet.* twice as long as the calyx. Flowering raceme lax. Pods short.—I cannot separate *B. arcuata* (*Sy. E. B.* 121).—In damp places. B. ? V.—VIII. Yellow Rocket. E. S. I.

2. *B. stricta* (Fries); lower l. lyrate, upper pair of lobes small much shorter than the breadth of the large oblong-ovate terminal lobe, uppermost l. undivided toothed, pods adpressed with a subulate point.—*Sy. E. B.* 122. *R.* ii. 47. *B. parviflora* Fries.—*Pet.* half as long again as the calyx. Flowering raceme close. Fl. much smaller than in *B. vulgaris*. Pods short. Lateral lobes of the lowermost l. very small, often obsolete.—Yorkshire, By Thames near Kew. B. ? V.—VIII. E.

†3. *B. intermedia* (Bor.); lower l. lyrate, upper pair of lobes equalling the breadth of the cordate-ovate term. lobe, upper l. all pinnatifid, pods erect with a short conical point.—*Sy. E. B.* 123.—St. with 2 angles,  $1\frac{1}{2}$ —2 ft. high. Raceme close. *Pet.* exceeding calyx. Pods thick, short, angular.—Near Manchester and Armagh. B. V.—VIII. E. I.

‡4. *B. præcox* (R. Br.); lower l. lyrate gradually larger upwards, upper pair of lobes equalling the breadth of the subcordate single terminal lobe, uppermost l. pinnatifid with linear-oblong entire lobes, pods patent with a short thick point.—*E. B.* 1129. *B. patula* Fries.—Fl. moderate. Raceme close. Pods long, torulose. Lower l. usually interruptedly pinnate.—South of England. B. V.—VII. E. I.

## 5. AR'ABIS Linn. Rock-Cress.

\* *Seeds in one row.*

1. *A. hirsúta* (R. Br.); l. hispid dentate, *stem-l. truncate-auricled or cordate at the base*, auricles patent, pods erect narrow straight.—*E. B.* 587.—St. 1 foot high, clothed with spreading mostly simple hairs and many erect leaves. Sometimes the hairs on the stem are adpressed and branched. Root-l. narrowed into a footstalk.—Walls and banks. B. VI.—VIII. E. S. I.

2. *A. ciliáta* (R. Br.); l. glabrous and ciliate or hispid somewhat toothed nearly sessile, *stem-l. sessile with a rounded base*, pods erect narrow straight.—*E. B.* 1746.—St. erect, glabrous. L. glabrous, ciliate. Seeds without wings. St. and l. sometimes hispid.—Rocks by the sea. Connemara, Ireland. Lidstep, Pembrokeshire. B. VII. VIII. E. I.

3. *A. stric'ta* (Huds.); l. hispid and ciliate deeply *sinuate-dentate*, narrowed into a footstalk, *stem-l. sessile*, pods few distant erect patent straight.—*E. B.* 614.—St. erect, glabrous, hispid below, 6—8 in. high. Fl. rather large. Seeds winged at the apex.—Limestone cliffs near Bristol. P. IV. V. E.

4. *A. petræ'a* (Lam.); l. glabrous or with forked hairs lyrate-pinnatifid or oblong-ovate nearly entire with long stalks, *stem-l. narrow nearly entire stalked*, pods spreading slender straight.—*E. B.* 469.—St. erect or decumbent, 3—8 in. long, glabrous. Fl. large, white tinged with purple. Seeds oblong with a straight wing at the end.—Alpine rocks in E. and S. Glenade Mountain, Co. Leitrim, Ireland. P. VII. VIII. E. S. I.

[\**A. Turríta* (L.); l. clothed with short forked hairs dentate elliptic narrowed into a stalk, *stem-l. deeply cordate-amplexicaul.* long, pods flat with a thickened margin recurved from an erect stalk, *seeds with a membranous margin.*—*E. B.* 178.—St. 1 foot high, erect hairy. Fl. yellowish. Pods 3—4 in. long, without any central vein but with many prominent longitudinal anastomosing veins.—On walls at Oxford, Cambridge, and Cleish Castle, Kinross. Become very rare. B. V.] E. S.

\*\* *Seeds in two rows.*

5. *A. perfoliáta* (Lam.); radical l. stalked toothed hairy, st.-l. glabrous entire clasping, pods slender erect.—*Turritis glabra* (L.) *E. B.* 777.—Very erect and straight, 1—3 ft. high. Fl. pale yellow.—Banks and cliffs, local. B. V.—VII. E. S. I.

## 6. CARDAMINE Linn. Bitter Cress.

1. *C. impatiens* (L.); l. pinnate, leaflets of the lower l. ovate 3-fid, of the upper l. oblong lanceolate toothed or entire, petioles of the stem-l. with slender sagittate auricles, pet. linear or wanting.—*E. B.* 80. *R.* ii. 26.—Pet. erect, white. Distinguished by having auricles at the base of its petioles.—Hilly districts, preferring limestone. A. VII. VIII. E. S. ?

2. *C. flexuosa* (With.); l. pinnate, leaflets of the lower l. roundish angled or toothed, of the upper l. narrower, pet. twice as long as the calyx, pods erect upon patent pedicels, *stam.* 6, *style long.*—*E. B.* 492. *R.* ii. 26. *C. sylvatica* Link.—Fl. small. Pet. erect, white. *St. wavy*, more leafy than in *C. hirsuta*. Root oblique, covered with fibres. In shady places. A. or B. IV.—IX. E. S. I.

3. *C. hirsuta* (L.); l. pinnate, leaflets of the lower l. roundish angled or toothed, of the upper l. narrower, pet. twice as long as the calyx, pods and pedicels erect, *stam.* 4, *style equal in length to about ½ the breadth of pod.*—*Sy. E. B.* 110. *R.* ii. 26.—Fl. small. Pet. erect, white. *St. nearly straight*, rather leafy.—In damp places. A. IV.—VIII. E. S. I.

4. *C. pratensis* (L.); l. pinnate, leaflets of the lower l. roundish slightly angled, of the upper l. linear-lanceolate entire, pet. 3 times as long as the calyx spreading, *stam.* half the length of the petals, *st.* terete.—*E. B.* 776. *R.* ii. 28.—Fl. large, lilac. Anth. yellow. Style short. Lts. sometimes stalked and occasionally acutely angular (*C. dentata* Schultz). *C. Hayneana* R. closely resembles Sp. 3, but belongs here.—Moist meadows, common' P. IV. *Lady's Smock.* E. S. I.

5. *C. amara* (L.); l. pinnate, leaflets of the lower l. roundish-ovate, of the upper l. oblong, all angular, pet. 3 times as long as the calyx erect, *stam.* nearly as long as the petals, *st.* angular.—*E. B.* 1000.—Fl. large, white. *Anthers purple.* Style long, slender. Stigma small. Stoloniferous.—Moist meadows near streams, rare. P. V. VI. *Common Bitter Cress.* E. S. I.

## 7. DENTA'RIA Linn. Coralwort.

1. *D. bulbifera* (L.); *st.* simple, l. alternate, lower l. pinnate, upper l. simple, axils of the l. producing bulbs.—*E. B.* 309.—Rhizome thick, with fleshy toothlike knobs. *St.* 1—1½ foot high. Lts. and l. lanceolate, serrate or entire. Fl. large, rose-coloured or purple.—Woods and shady places, rare. P. V. VI. E. S.

Tribe II. *Sisymbriæ*.

## 8. HES'PERIS Linn. Dames' Violet.

\*1. *H. matronalis* (L.); st. erect branched above, l. ovate-lanceolate acuminate toothed, pedicels about as long as the calyx, pet. obovate blunt apiculate, pods erect from a patent pedicel terete.—*E. B.* 731.—Fl. lilac, "fragrant," large and handsome.—Pastures. B. V. VI. E. S. I.

[*Malco'lmia marit'ima* (R. Br.) has been found in Kent, but is not a native.]

## 9. SISYMBRIUM Linn. Hedge-Mustard.

1. *S. officinale* (Scop.); pods subulate adpressed to the stem, l. runcinate-pinnatifid with 2 or 3 pairs of oblong dentate lobes and a large hastate terminal lobe.—*E. B.* 735. *R.* ii. 72.—St. 1—2 feet high, with divaricate branches, upper part leafless. Fl. small, pale yellow. Pods on exceedingly short stalks, downy, sometimes glabrous.—Common. A. VI. VII. *Hedge Mustard*. E. S. I.

[\**S. polycerátium* (L.); pods subulate spreading sessile axillary about 3 together, l. lanceolate repand-dentate or subhastate.—*Sy. E. B.* 97. *R.* ii. 73.—St. leafy throughout, branched, "prostrate." Fl. small.—Bury St. Edmunds, naturalized. A. VII. VIII.] E.

2. *S. Irio* (L.); pods terete 4 times as long as their pedicels erect-patent, the young pods exceeding the fl., seeds oblong, l. runcinate-pinnatifid, lobes dentate oblong the terminal lobe angular, lobes of the upper l. lanceolate with the terminal lobe hastate.—*E. B.* 1631. *R.* ii. 75.—Erect, branched, st. and l. glabrous. Fl. yellow. Pods narrow, linear. Pedicels slender.—Near old towns, rare. A. VII. VIII. *London Rocket*. E. S. I.

3. *S. Sophía* (L.); pods terete 3 times as long as the pedicels erect-patent, seeds oblong, l. doubly or trebly pinnatifid, segments linear or linear-lanceolate.—*E. B.* 963. *R.* ii. 84.—St. erect, branched, and as well as the l. slightly downy. Fl. yellow. Pet. short. Pods linear, narrow. Pedicels slender.—Waste places, not common. A. VI.—VIII. *Flixweed*. E. S. I.

4. *S. thabianum* (Gaud.); pods 4-angular linear ascending twice as long as their patent pedicels, seeds oblong not striate, l. oblong-lanceolate undivided toothed.—*Arabis Sm. E. B.* 901. *Coringia R.* ii. 60.—St. erect, slender, much branched, with

few leaves which are nearly all radical. Fl. small, white. *Pods angular*, not convex on the back as in the other species nor with the lateral longitudinal veins so strongly marked.—On walls and banks. A. IV. V. and IX. X. E. S. I.

[*S. pannonicum* (Jacq.) is quite established near Crosley, Lanc. *J. of B.* x. 239.]

#### 10. ALLIARIA Adans. Sauce-alone.

1. *A. officinalis* (Andrzj.); l. heartshaped the lower ones reniform sinuate-dentate all stalked, pods erect-patent much longer than their stalks, seeds oblong subcylindrical striate.—*R.* ii. 60. *Sisymbrium Alliaria* Sm. *E. B.* 796.—St. erect, 1—3 feet high, slightly branched. L. large, thin, veined, smelling like garlic when bruised. Fl. white.—Hedge-banks. B. V. VI. *Jack-by-the-Hedge.* E. S. I.

#### 11. ERYSIMUM Linn. Worm-seed.

1. *E. cheiranthoides* (L.); l. oblong-lanceolate slightly toothed with stellate-tripartite hairs, all narrowed into a slight foot-stalk, pedicels longer than the calyx 2- or 3-fold shorter than the pods, ped. patent, pods suberect, seeds small many.—*E. B.* 942. *R.* ii. 83.—Seeds very small, so many in the pod as to be nearly 2-rowed. Fl. small.—Cultivated ground, rare. Wild in the Fen country. B. VI.—VIII. E.

[*E. orientale* (R. Br.); l. oval-heartshaped blunt clasping the stem, radical l. obovate, all smooth glaucous undivided entire.—*E. B.* 1804. *R.* ii. 61.—Fl. white or cream-coloured.—Fields. Introduced. A. V.—VII.] E.

### Tribe III. Brassicæ.

#### 12. BRAS'SICA.

\* *Valves of pod 1-ribbed.*

1. *B. oleracea* (L.); l. glabrous glaucous waved and lobed, lower l. lyrate, upper l. oblong sessile.—*E. B.* 637. *R.* ii. 97.—Rootstock stout, branched. St. thick, persistent, usually decumbent. L. very large, thick, somewhat fleshy. Fl. large, cream-coloured. Raceme elongated before the fl. expand. Cal. erect, adpressed. The wild state of the garden cabbage.—Sea-cliffs in South and West. B. VI.—VIII. *Wild Cabbage.* E. S. ? I.

2. *B. campestris* (L.); radical l. glaucous hispid lyrate-dentate, st.-l. glabrous ovate-lanceolate auricled clasping, fl. subcorymbose.—*a*; fl. pale orange (*Rape*).—*β*. *B. Rapa* (L.); not glaucous, fl. bright yellow (*Turnip*).—*γ*. *B. Napus* (L.?) ; glaucous, fl. yellow (*Rape*) or buff (*Sweed*). Root-l. of all hispid.—See Watson and Dyer in *J. of B.* vii. viii. and ix.—Fields and riverbanks. A. or B. VII. VIII. E.

\*\* *Valves of pod 3-ribbed; beak 1—3-seeded.*

3. *B. monensis* (R. Br.); l. stalked all deeply pinnatifid, lobes oblong unequally toothed those of the upper l. linear.—*E. B.* 962.—Fl. yellow. St. usually prostrate, glabrous. L. glabrous.—*β*. *B. Cheiranthus* (Vill.); st. 1—3 ft. high erect leafy hispid below, l. hispid. *E. B. S.* 2821.—On the western coasts, rare. B. or P. VI.—VIII. E. S.

### 13. SINA'PIS Linn. Mustard.

1. *S. nigra* (L.); pods quadrangular adpressed, beak short sterile subulate, valves 1-veined, lower l. lyrate, terminal lobe large and lobed, upper l. lanceolate entire.—*E. B.* 969. *R.* ii. 88.—Fl. yellow. Lower l. large, rough.—Willowy riverbanks, not common. A. VI.—VIII. *Black Mustard.* E. S. I.

2. *S. arvensis* (L.); pods subcylindrical knotty longer than the conical compressed beak, valves 3-veined, l. ovate the lowermost sublyrate stalked, upper l. sessile.—*E. B.* 1748. *R.* ii. 86.—Scabrous. Fl. large, yellow. Pods glabrous or rough with deflexed bristles; beak with one seed; valves with faint intermediate veins. Stem 1—1½ foot high.—Corn-fields. A. VI.—VIII. *Charlock.* E. S. I.

†3. *S. alba* (L.); pods cylindrical knotty shorter than the swordshaped beak, valves 5-veined, l. lyrate pinnatifid irregularly lobed.—*E. B.* 1677. *R.* ii. 85.—Fl. large, yellow. Pods hispid. St. 1—2 feet high.—Cultivated and waste calcareous land. A. VII. *White Mustard.* E. S. I.

[*S. incana* (L.); pods adpressed thick prominently veined with a short 1-seeded beak, l. lyrate hispid, stem-l. linear-lanceolate, st. much branched.—*E. B. S.* 2843.—St. 1—3 feet high, branches divaricate with few very small leaves. Pods very short, glabrous or hairy, often scarcely longer than their glabrous beak.—Sandy places in Jersey and Alderney. B. VII. VIII.]

14. DIPLTAX'IS *Cand.* Wall-Mustard.

1. *D. tenuifolia* (DC.); *st. woody below branched subglabrous leafy*, l. glaucous linear-lanceolate very acute sinuate-dentate or pinnatifid, segments linear remotely dentate, ped. very long, pet. roundish-obovate with a short claw.—*Sinapis* Sm. *E. B.* 525.—Fl. large; pet. blunt but slightly acuminate. St. 1—1½ foot high. Plant fœtid.—Old walls. P. VII.—IX. E. S. I.

2. *D. muralis* (DC.); *st. herbaceous simple hispid and leafy at the base*, l. almost glabrous ovate-lanceolate sinuate-dentate or pinnatifid, ped. as long as expanded flower, pet. roundish-ovate with a short claw.—*Sinapis* Sm. *E. B.* 1090.—Pedicels as long as the flowers. Pet. abrupt or emarginate. L. often blunt, never very acute.—We have two forms: (1) with the base of st. leafy with axillary branches; (2) l. all radical in a rosette and st. simple.—Waste ground. A. VIII. IX. E. S. ? I.

Suborder II. *Latiseptæ.* Tribe IV. *Alyssineæ.*15. ALYS'SUM *Linn.*

\*1. *A. calycinum* (L.); herbaceous hoary with starry pubescence, l. obovate-lanceolate attenuate below, pods orbicular stellate-pubescent, calyx persistent, filaments all toothless, shorter ones from between 2 setaceous processes.—*E. B. S.* 2853. *R. ii.* 18.—Cells of the pod 2-seeded. Pet. yellow, becoming at length white.—Grassy commons, S.; ploughed land, E. A. V. VI. E. S.

[*A. maritimum* (L.); procumbent, hairs bipartite, l. linear-lanceolate acute, cal. deciduous, pods oval pointed glabrous.—*E. B.* 1729. *R. ii.* 18.—St. rather woody below. Fl. white, sweet-scented.—Naturalized near the sea. P. VIII. IX.] E.

16. DRA'BA *Linn.* Whitlow-grass.

1. *D. aizoides* (L.); st. leafless glabrous, l. linear rigid acute keeled glabrous ciliate, stam. equalling the slightly notched petals, style as long as the breadth of the pouch.—*E. B.* 1271. *R. ii.* 15.—Fl. bright yellow. L. fringed with rigid hairs, densely collected into cushion-like tufts.—On rocks and walls at and near Pennard Castle near Swansea. P. III. IV. E.

2. *D. rupes'tris* (R. Br.); st. leafless or with 1 or 2 leaves pubescent, l. lanceolate flat stellately pubescent, stam. shorter than



the slightly notched petals, style short.—*D. hirta* Sm. *E. B.* 1338.—Pouch long-oval, slightly hairy. St. very short, branched, each branch bearing a dense tuft of leaves and 1—3 short scapes. Fl. small. L. mostly entire.—Tops of the Scottish mountains, rare. Ben Bulbin, I. P. VII. S. I.

3. *D. incána* (L.); stem-l. several, l. lanceolate stellately pubescent toothed, pet. twice as long as the calyx, pouch longer than its pedicel twisted, style very short.—*E. B.* 388. *R.* ii. 14.—Pouch nearly or quite glabrous, erect, lanceolate-oblong. St. 4—12 in. high, simple or branched. Fl. white.—Extreme forms are often taken for species, as *D. contorta* (Ehrh.), st. simple or branched only at top densely leafy, pouch glabrous; and the more alpine state *D. confusa* (Ehrh.), st. branched only near its base, l. mostly rosulate, pods with stellate down.—Mountains and sand-hills by the sea. P. VI. VII. E. S. I.

4. *D. murális* (L.); st. leafy branched, l. ovate amplexicaul toothed hairy, pet. "entire," pedicels spreading horizontally rather longer than the glabrous pouch.—*E. B.* 912. *R.* ii. 12.—Pouch elliptic. St. 5—12 in. high. Fl. white. Pubescence branched. Root-l. narrowed below. Limestone hills. A. IV. V. E. S. I.

5. *D. ver'na* (L.); st. leafless glabrous above, l. lanceolate acute narrowed below hairy, pet. deeply cloven, pouch oblong shorter than its pedicel.—*E. B.* 586. *R.* ii. 12.—Pouch compressed, acute or rounded at the end, often rather inflated. Fl. white. L. entire or toothed. Scape sometimes rather pilose.—Very variable in minute points.—Very common on walls, banks, &c. A. III.—V. Common Whitlow-grass. E. S. I.

### 17. COCHLEA'RIA Linn. Scurvy-grass.

1. *C. officinális* (L.); radical l. cordate-reniform stalked, stem-l. sessile oblong sinuate clasping, pouch globose or obovoid, style short.—*E. B.* 551.—Petioles long. Lower l. entire or sinuate. Seeds large. St. occasionally rooting and proliferous—*C. alpina* (Wats.); pouch obovoid narrowed at both ends, leaves smaller. *E. B.* 2403.—Sea-coast, mostly in muddy places.  $\beta$  on the higher parts of mountains. B. ? VI.—VIII. Common Scurvy-grass. E. S. I.

2. *C. dan'ica* (L.); l. stalked, radical l. cordate somewhat lobed, stem-l. 3—5-lobed rather triangular uppermost sessile, pouch ovoid, style short—*E. B.* 696. *R.* ii. 16.—Petioles of the root-l. very long, gradually shortening as they become

more distant from the root. *Seeds small*.—Sea-coast. B. V.—VIII. E. S. I.

3. *C. ang'lica* (L.); radical l. stalked *ovate-oblong* entire, stem-l. oblong entire or toothed mostly sessile the *upper ones clasping, pouch ovate-oblong*, style slender.—*Sy. E. B.* 133.—Pouch twice as large as that of *C. officinalis*, much compressed laterally, usually deeply furrowed on each side, dissepiment very narrow. Seeds large. Lower l. rounded below or narrowed into a footstalk. Fl. large.—Muddy sea-shores. A. V. *English Scurvy-grass*. E. S.

### 18. ARMORA' CIA Rupp.

[*A. rustica'na* (Rupp.); rootstock long and thick, radical l. oblong crenate-serrate on long stalks, stem-l. long lanceolate incise-serrate or entire subsessile, pet. (white) twice as long as cal., pouch ovoid 4-seeded, stigma peltate.—*Coch. Armoracia* L. *E. B.* 2323. *R.* ii. 17.—St. 2—3 feet high. Rootstock cylindrical.—Waste ground, not native. P. V. *Horse-radish*.] E. S. I.

1. *A. amphib'ia* (Koch); roots fibrous, l. oblong narrowed at both ends serrate or pinnatifid, pet. (yellow) twice as long as the cal., pouch ovoid, "stigma capitate."—*Nasturtium* R. Br. *E. B.* 1840.—St. 2—3 feet high. Submersed l. deeply pinnatifid. Ped. usually deflexed.—Watery places. P. VI. VIII. E. S. ? I.

### Tribe V. *Camelineæ*.

#### 19. CAMELI'NA Crantz. Gold-of-pleasure.

[*C. sativa* (L.); pouches obovate inflated.—*E. B.* 1254.— $\alpha$ ; pouches flaccid truncate.— $\beta$ ; pouches rounded at the end brittle.—In fields of flax and corn, introduced, but not naturalized.]

### Tribe VI. *Velleæ*.

#### 20. VEL'LA Linn.

[*V. an'nuua* (L.); "l. doubly pinnatifid, pouches deflexed."—*E. B.* 1442.—Salisbury Plain. An error? A. VI.] E.

Suborder III. *Angustiseptæ*. Tribe VII. *Thlaspideæ*.

21. THLAS'PI *Linn.* Penny Cress.

1. *T. arven'se* (L.); fruitbearing raceme elongate, *pouch orbicular with a broad dorsal wing*, seeds concentrically rugose and striate 5—6 in each cell, stem-l. oblong sagittate toothed.—*E. B.* 1659. *R.* ii. 5.—Pouch very large ( $\frac{1}{2}$ — $\frac{3}{4}$  in.), stigma sessile. St. often a foot high.—Fields and roadsides. A. V.—VII. *Penny Cress.* E. S. I.

2. *T. perfoliatum* (L.); fruitbearing raceme elongate, *pouch obcordate* broadly winged above, *style very short* included within the notch, seeds 3—4 in each cell smooth, *stem-l. cordate-oblong*.—*E. B.* 2354. *R.* ii. 5.—About 6 in. high.—Limestone in Oxfordshire and Gloucestershire. A. V. E.

3. *T. alpes'tre* (L.); fruitbearing raceme elongate, *pouch oblong-obovate* narrowed below, *style equalling or exceeding the notch*, seeds 4—8 in each cell, stem-l. oblong cordate.—*R.* ii. 5.—Fl. white, often tinged with rose, small. Pet. about twice as long as the calyx.—*a. T. sylvestre* (Jord.), terminal lobes of pouch rounded. L. usually entire. *Sy. E. B.* 146.—*β. T. occitanum* (Jord.); fruitbearing raceme rather shorter, pouch triangular-obcordate with almost divaricate lobes, style much exserted. L. usually slightly toothed. *Sy. E. B.* 147.—Mountain pastures. Teesdale; Thornhaugh, Northumb.; Glen Isla, Forfar. *β.* Settle, York; Llanrwst, N. Wales. P. VI.—VIII. E. S.

4. *T. virens* (Jord.); fruitbearing *raceme oval or oblong*, pouch obovate scarcely notched, style much projecting, seeds 4—5 in each cell, stem-l. oblong cordate.—*Sy. E. B.* 148. *Jord. Obs. Pl. de Fr.* iii. t. 1. *T. alpestre* Sm.—Fl. white, small. Pet. about thrice as long as the calyx. Pouch with a very broad and shallow notch often almost truncate. L. usually entire.—Limestone rocks at Matlock. P. VI.—VIII. E.

22. HUTCHINS'IA *R. Br.*

1. *H. petræ'a* (R. Br.); l. pinnate, st. branched leafy, fr.-raceme lax long, pouch blunt at both ends.—*E. B.* 111. *R.* ii. 6. *St.* 65. 10.—St. 2—4 in. high.—Fl. small; pet. scarcely exceeding the calyx. Cotyledons accumbent.—Limestone rocks, rare. A. III.—V. E.

[*H. alpina* (R. Br.); l. pinnate, st. simple leafless, fr.-raceme lax long, pouch acute at both ends.—*R.* ii. 11. *St.* 20. 11.—

St. 2—5 in. high.—Said to have been found on Ingleborrow.  
*J. of B.* i. 359, and iii. 92 and xviii. 375.]

### 23. TEESDA'LIA R. Br.

1. *T. nudicaulis* (R. Br.); petals unequal.—*E. B.* 327. *R.* ii. 6.—L. many, spreading on the ground, lyrate-pinnatifid, rarely orbicular-spathulate and entire. St. 2—4 in. high, solitary or several from the crown of the root, sometimes bearing 1 or 2 small leaves. Stam. with remarkable scales. Pouch emarginate.—Sandy and gravelly places. A. V. VI. E. S.

### 24. IBE'RIS Linn. Candytuft.

1. *I. amara* (L.); herbaceous, l. lanceolate somewhat toothed, pouches racemose orbicular notched with triangular pinnate lobes.—*E. B.* 52. *R.* ii. 7.—L. usually with 1—3 blunt teeth on each side. St. often 1 foot high, diffuse, branched. Fl. at first corymbose, afterwards in lengthened clusters. Outer pet. radiant.—Chalky fields in South and East. A. VII. *Bitter Candytuft.* E.

### 25. LEPID'IUM Linn. Pepperwort.

\* *Pouch cordate with turgid valves, style filiform.*

[*L. Drába* (L.); l. oblong entire or toothed lower ones narrowed into a footstalk, stem-l. sagittate and amplexicaul, style as long as the dissepiment.—*E. B. S.* 2683. *R.* ii. 9.—One foot or more in height, branched. Fl. many, small, white, upon long pedicels in a subumbellate corymb.—Established in many places, but not naturalized. P. V. VI.] E.

\*\* *Pouch ovate or roundish winged notched.* † Style manifest.

1. *L. campestre* (R. Br.); l. downy toothed lower ones oblong narrowed into a footstalk, stem-l. lanceolate sagittate and clasping, pouch ovate scaly notched and rounded at the end, style scarcely longer than the notch.—*E. B.* 1385. *R.* ii. 9.—Anth. yellow. Scales on the pouch only minute globular blisters when fresh. St. one, upright, about a foot high, branched in the upper part.—Dry gravelly soil. B. VI.—VIII. E. S. I.

2. *L. Smith'ii* (Hook.); l. hairy toothed lower ones obcordate stalked, stem-l. lanceolate-sagittate clasping, pouch ovate smooth notched and rounded at the end, style twice as long as the notch.—*L. hirtum*, Sm. *E. B.* 1803.—Seeds  $\frac{1}{2}$  as long in *L. campestre*. Pouch sometimes with a few scales, never hairy. St. several, 6—12 in. long; central st. erect, others diffuse.—Hedge-banks. P. VI.—VIII. E. S. I.

†† Style minute.

[*L. sativum* (L.); lower l. lobed pinnate or bipinnate, upper l. sessile linear entire, pouch roundish oval blunt.—*R.* ii. 9.—Escaped from cultivation in many places.] E.

\*\*\* *Pouch oval or roundish notched, style minute.*

3. *L. ruderale* (L.); lower l. pinnatifid, upper l. linear entire, pouch oval, diandrous, petals 0.—*E. B.* 1595. *R.* ii. 10.—End of pouch narrowed, winged. St. branched, often a foot high. Radicle dorsal.—Waste places near the sea, rare. A. V. VI. E. I.

\*\*\*\* *Pouch oval or roundish scarcely notched wingless, style minute.*

4. *L. latifolium* (L.); l. ovate-lanceolate serrate or entire undivided, pouch oval entire downy.—*E. B.* 182. *R.* ii. 10.—Fl. many, small, in compound leafy paniced clusters. St. 3 feet high, erect, branched. L. large, the lower ones upon long stalks, the upper nearly sessile and narrower.—In salt marshes. P. VII. VIII. *Dittander.* E. S. ? I.

## 26. CAPSEL'LA Vent. Shepherd's Purse.

1. *C. Bursa-pastoris* (DC.); radical l. lanceolate pinnatifid or undivided toothed, upper l. clasping auricled, pouch triangular-obcordate.—*Thlaspi* Sm. *E. B.* 1485.—Varying greatly in size and the division of its leaves. Known by its peculiar pouches.—A common weed. A. III.—X. E. S. I.

## Tribe VIII. *Subulariæ*.

### 27. SUBULA'RIA Linn. Axlwort.

1. *S. aquatica* (L.); *E. B.* 732. *R.* ii. 12.—Plant small, submerged. Root of many long white fibres. St. 1—3 in. high. L. linear-subulate, radical. Fl. small, white, often perfected under water.—Margins of alpine lakes. P. VII. E. S. I.

## Tribe IX. *Senebieræ*.

### 28. SENEBIE'RA Pers. Wart-Cress.

1. *S. Corónopus* (Poiret); *pouch undivided reniform* wrinkled and crested with little sharp points, style prominent, l. pinnatifid.—*E. B.* 1660. *R.* ii. 9.—St. much branched, prostrate.—Fl.

small, white, in lateral clusters opposite to the leaves. Pouches large, in dense clusters.—Waste ground, common. A. VI.—IX. *Swine's Cress.* E. S. I.

2. *S. did'yma* (Pers.); *pouch notched of two wrinkled lobes*, style very short, l. pinnatifid.—*E. B.* 248. *R.* ii. 9.—St. spreading, prostrate, a foot or more in length. Fl. small, white, in long slender lax clusters. Pet. very short or none. Often only 2 stamens.—Waste ground near the sea in the South and South-west. A. VII.—IX. E. I.

#### Suborder IV. *Nucumentaceæ.* Tribe X. *Isatideæ.*

##### 29. I<sup>S</sup>ATIS *Linn.* Woad.

†1. *I. tinctoria* (L.); “radical leaves oblong crenate,” pouch abrupt smooth thrice as long as broad.—*E. B.* 97. *R.* ii. 4.—St. 1—4 ft. high branched. Fl. yellow. Pouches  $\frac{1}{2}$  in. long, pendent.—Waste land, rare. B. VII. E.

#### Suborder V. *Lomentaceæ.* Tribe XI. *Cakilineæ.*

The fruit consists of a very small 2-celled, stalklike, usually sterile pod, with a long moniliform beak bearing the seeds and dividing transversely into as many indehiscent cells as there are seeds.

##### 30. CAKI'LE *Gaert.* Sea-Rocket.

1. *C. marit'ima* (Scop.); joints of the pouch 2-edged, upper with 2 teeth at the base, l. fleshy pinnatifid or somewhat toothed.—*E. B.* 231. *R.* ii. 1.—Fl. purplish. Pouches an inch long, erect, with 4 sharp angles, swordshaped in the upper part.—Sandy sea-shores. A. VI. VII. *Purple Sea-Rocket.* E. S. I.\*

##### 31. CRAM'BE *Linn.* Sea-Kale.

1. *C. marit'ima* (L.); pouch without a style, l. roundish sinuate wavy toothed glaucous and as well as the st. glabrous.—*E. B.* 924. *R.* ii. 2.—Root thick, fleshy. St. 2 feet high. Fl. white. Longer filaments forked at the end. Pouches roundly ovoid, large.—Sandy sea-shores. P. VI. E. S. I.

#### Tribe XII. *Raphaneæ.*

##### 32. RAPH'ANUS *Linn.* Radish.

†1. *R. Raphanistrum* (L.); pods moniliform shorter than

the very long beak with slender ribs, *l. simply lyrate*.—*E. B.* 856.—Root slender. Lobes of the leaves quite distinct. Pet. veined, yellow, white, or lilac.—Corn-fields. A. VI. VII. *Jointed Charlock*. E. S. I.

2. *R. maritimus* (Sm.); pods moniliform longer than the short beak with thick ribs, *radical l. interruptedly pinnate*.—*Sy. E. B.* 82.—Root thick. Lobes of *l.* usually so close as to overlap each other. Pet. yellow.—Sea-coast in South and West, rare. B. ? VI.—VIII. *Sea-Radish*. E. S. I.

### Order VII. RESEDACEÆ.

Sep. 4—8, persistent. Cor. irregular, pet. 4—8 entire or deeply cut. Stam. many, filaments variously united, inserted on a glandular irregular 1-sided hypogynous disk. Ovary 3- or 6-lobed, 1-celled, with 3 or 6 parietal many-seeded placentas; or of several 1-celled carpels. Fruit opening nearly at the end.—Stip. 0?

1. RESEDA. Cal. many-parted. Pet. entire or variously cut, unequal. Stam. many. Fruit of one cell opening at the top. Styles 3—6.

#### 1. RESE'DA Linn. Mignonette.

1. *R. hutea* (L.); sep. 6 linear, *pet. very unequal*, ped. longer than the cal., *l. 3-cleft or pinnatifid*.—*E. B.* 321. *R. ii.* 100.—St. 2 feet high, branched, smooth. L. very variable. Two upper pet. with two wing-like lobes, lateral pet. with a single wing, lower ones nearly entire. Fl. yellow. Fr. oblong, wrinkled.—Waste chalky and limestone places. B. VI.—VIII. E. S. I.

‡2. *R. suffruticulosa* (L.); sep. 5 linear-lanceolate, *pet. 5 nearly equal 3-fid* longer than the calyx, ped. shorter than the cal., *l. all pinnatifid*, segments linear acute sometimes wavy.—*E. B. S.* 2628. *R. ii.* 101.—St. 1½—2 feet high, rather shrubby below. Fl. white. Fr. oblong, wrinkled. Sometimes there are 6 sep. and pet., when it is *R. alba* (L.).—Waste sandy places near the sea, rare. B. or P. VII. VIII. E. S. I.

3. *R. Lutéola* (L.); sep. 4, pet. 4 or 5 very unequal longer than the calyx, *l. long-lanceolate undivided*.—*E. B.* 320. *R. ii.* 99.—St. 2 feet high. Pet. usually 4, upper one 3-, 4-, or 5-cleft, 2 lateral 3-cleft, segments linear, lower one (or two) linear entire. Fr. broad, depressed.—Waste places, particularly on chalk or limestone. B. VII. VIII. *Weld*. E. S. I.

## Order VIII. CISTACEÆ.

Sep. 5, two outer smaller sometimes wanting, 3 inner twisted in the bud. Pet. 5, crumpled and twisted in the bud the contrary way to the sepals. Stam. many. Ovary 1, 1- or many-celled. Style simple. Stigmas 3. Capsule 3-, 5-, or 10-valved. Embryo spiral or curved, in the albumen.—Stip. small or 0.

1. HELIANTHEMUM. Cal. of 5 sepals, 2 outer smaller. Pet. 5, deciduous. Stam. many. Caps. 3-valved.

## 1. HELIANTHEMUM Gaert. Rock-rose.

1. *H. guttatum* (Mill.); erect herbaceous, l. oblong-lanceolate or linear, lower l. opposite without stipules, upper l. alternate stipulate, *stigma subsessile*.—St. mostly simple or branching from the base. Fruitstalks patent. Pubescence of long simple, and short stellate hairs. Fl. yellow, usually with a deep-red spot at the base of each petal.— $\alpha$ ; racemes without bracts. *E. B.* 544.— $\beta$ . *H. Breweri* (Planch.); racemes with or without bracts on the same plant. *Curt.* ii. 102. *J. of B.* iii. 21. *Sy. E. B.* 166.—Very rare.  $\alpha$ . Three-Castle Head, Co. Cork, Jersey.  $\beta$ . Holyhead Mountain and Amlwch, Anglesea. A. V.—VIII.

E. I.

2. *H. canum* (Dun.); shrubby, without stipules, l. opposite ovate or oblong stalked flat hoary beneath, *racemes* terminal with bracts, “style twisted at the base reflexed but at the apex inflexed.”—*Cistus marifolius* Sm. *E. B.* 396.—St. decumbent. L. hoary beneath, hairy above. Fl. yellow, small. “Anth. emarginate at both ends. Style longer than the stigma.” The Teesdale plant, *H. vineule* (Pers.), has rather fewer hairs on the upperside of the leaves—On limestone rocks, rare. P. V.—VII.

E. I.

[*H. ledifolium* (Willd.); herbaceous, with stipules, ped. solitary, styles straight.—*E. B.* 2414.—Brean Down, Somerset. Probably an error. A. VI. VII.]

3. *H. vulgare* (Gaert.); procumbent, shrubby, with stipules, l. oval or linear-oblong opposite nearly flat green above hoary beneath, *racemes* with bracts, style longer than the germen bent at the base, sep. subglabrous inner ones blunt apiculate.—*Cistus* L. *E. B.* 1321.—Fruitstalks contorted and deflexed. Varying much in the size and shape of its leaves and the amount of hoariness and pubescence. Fl. yellow.—*C. tomentosus* E. B. 2208. scarcely differs. *C. surrejanus* E. B. 2207. is a garden form. *H. vulg. pet. fl. perangustis* (Dill. H. Elth. 145) was



again found near Croydon by the late Mr. Christy and is a monstrosity of *H. vulgare*.—Common on dry hilly places. P. VII.—IX. *Common Rock-rose*. E. S.

4. *H. polifolium* (Pers.); shrubby, procumbent, hoary, with stipules, l. opposite ovate-oblong or oblong-linear more or less revolute at the edges, racemes with bracts, style bent at the base longer than the germen, *sep. tomentose inner ones blunt*.—*E. B.* 1822.—*Fl. white*.—Very rare. Brean Down, Som.; Torquay, Devon. P. VI. VII. E.

### Order IX. VIOLACEÆ.

Sep. 5, imbricate. Pet. 5, regular or irregular. Stam. 5, filaments dilated, connective extended beyond the anthers as a flat membrane. Ovary 1-celled with 3 parietal placentas. Style with a hooded stigma. Caps. with 3 valves. Embryo straight, in fleshy albumen.—Stip. persistent.—Minute but fertile fl. which do not open are found on many species.

1. VIOLA. Sep. 5, extended at the base. Pet. 5, unequal, the lower one extended into a hollow spur behind. Stam. 5. Anth. connate, 2 lower ones spurred behind.

#### 1. VIOLA Linn. Violet.

A. *Two interm. pet. patent laterally. Style from a slender base.*

\* St. creeping. Stigma flat above. Fr.-st. erect. Caps. nodding.

1. *V. palustris* (L.); anth.-spurs short thick rounded, cor.-spur very short blunt, l. *reniform-cordate* glabrous.—*E. B.* 444. *R.* iii.—Anth.-cells nearly parallel. Fl. pale lilac with purple streaks. Scentless.—Bogs in mountainous districts. P. IV.—VI. *Marsh-Violet*. E. S. I.

\*\* Rhizome short. Stigma hooked; beak deflexed. Fruit-stalks prostrate. Caps. globular, downy.

a. *Stoles from axils of terminal rosettes.*

2. *V. odorata* (L.); anth.-spurs *lancet-shaped* decurved blunt, cor.-spur blunt straight, lateral pet. entire lower one emarginate, l. *cordate, with stoles*.—*E. B.* 619. *R.* iii.—Anth.-cells diverging below. Spurs of the pet. inflated towards the end, slightly channelled above. Fl. purple, often white, sweet-scented. Fr.-sep. triangular, acute, twice as long as broad, not ciliate, glabrous. *Bracts above the middle of the flowerstalk. Petioles with*

*deflexed hairs*.—Several species (?) allied to this are nearly or quite scentless: *V. permixta* (Jord.) has large glabrous glandular-denticulate stip. ciliate towards the tip, short robust stoles usually not rooting, fl. pale blue: *V. sepincola* (Jord.) has large glandular-denticulate stip., longer somewhat rooting stoles, and much darker flowers. These plants and their allies seem quite different when growing, but are very difficult to define. Some authors join them to *V. hirta*.—Common. P. III. IV. *Sweet Violet*. E. S. I.

b. *Stoles wanting or very short*.

3. *V. hirta* (L.); *anth.-spurs* nearly linear blunt, spur of the cor. blunt hooked at the end, pet. entire or slightly emarginate, l. cordate-ovate, *stoles wanting*.—*E. B.* 894. *R.* iii.—*Anth.-cells* diverging below. Spur of the petals compressed, not channelled. Fl. pale blue sometimes white, scentless. Fr.-sep. roundly triangular, bluntish, as long as broad, more or less ciliate, mostly downy. *Bracts* below the middle of the flowerstalk. Stip. not hispid at the margin. *Petioles* with spreading hairs.— $\beta$ . *calcareæ* (Bab.); fl. smaller, ped. much longer than the leaves, sep. oblong-ovate.—*V. sciaphita* (Koch?), found by Newbould at Harlton near Cambridge, differs by having larger l. with the basal lobes more spreading and *ovoid glabrous capsules*.—Common on limestone. B. Gogmagog Hills. Portland. P. IV. *Hairy Violet*. E. S. I.

\*\*\* Stigma hooked; beak horizontal. Having a stem.  
Fruitstalks erect.

† *Without a true sobole*.

4. *V. sylvatica* (Fries); *anth.-spurs* narrowly lancet-shaped (5 times as long as broad), cor.-spur blunt, caps. glabrous, l. broadly cordate, *flowering branches axillary from a short flowerless central rosette of leaves*.—Fl. scentless.—It is probable that the following are distinct species:— $\alpha$ . *V. Reichenbachiana* (Bor.); l. cordate-prolonged, calycine appendages small becoming indistinct, pet. oblong narrow (*black*), *lower with few parallel nearly simple veins* at the base, spur compressed entire (usually darker than the petal). *E. B. S.* 2986.— $\beta$ . *V. Riviniana* (R.); l. broad cordate-acute, calycine appendages broad and squarish persistent with fruit, pet. broadly obovate (*blue*), *lower with many branched dark veins at the base*, spur thick (usually yellowish white). *E. B.* 620. *Curt.* i. 182. *V. flavicornis* (Forst.) is a dwarf form having small l. and large flowers.—Common on hedge-banks or in thickets. P. IV. V. *Wood-Violet*. E. S. I.

5. *V. arenaria* (DC.); anth.-spur very narrowly lancet-shaped, cor.-spur blunt, l. roundly cordate, flowering branches axillary from a short flowerless central rosette of leaves, ped. young l. and acute caps. downy, pet. broadly obovate, lower pet. with many branched veins throughout, cal.-appendages broad squarish persistent.—*Sy. E. B.* 174 b. *R.* iii. 9.—A small compact plant with large flowers.—Elevated pastures. Upper Teesdale. B. V. VI. E.

6. *V. canina* (L.); anth.-spur lancet-shaped (3 times as long as broad), cor.-spur blunt, l. cordate-ovate roundly acute, primary and lateral stems flowering and lengthening.—*R.* iii. 10. *E. B. S.* 2984.—Rather caespitose. L. always roundedly acute, longer than those of Sp. 4. Fl. bluish purple, scentless. Cor.-spur yellow, 1—3 times as long as cal.-appendages. Lower pet. spatulate.—*V. flavicornis* (Sm.) is a small form with cordate leaves. When the l. are cordate-oblong it is perhaps *V. montana* (Linn.).— $\beta$ . *V. lancifolia* (Thore); l. ovate-lanceolate rounded below, stip. lanceolate incise-serrate. *V. lactea* Sm. *E. B.* 445. *V. pumila* Fries (not Vill. which is *V. pratensis* Fr. and has a sobole). L. narrowing gradually from near their base, to a narrow but rounded point.—Sandy and peaty places.  $\beta$  in turf-bogs, rare. P. IV. V. Dog-Violet. E. S. I.

†† Rhizome or rather sobole slender.

7. *V. stagnina* (Kit.); anth.-spur short broadly lancet-shaped acute (not twice as long as broad), cor.-spur very short blunt, l. ovate-lanceolate subcordate below, petioles winged at the top, stip. linear-lanceolate incise-serrate shorter than the petioles, primary and lateral st. flowering and elongated.—*E. B. S.* 2985. *V. lactea* R. iii. 16. not Sm.—Sobole threadlike. St. erect. L. narrowing gradually from the base which in the lowest is sometimes rounded not cordate. Fl. pale blue, nearly white. Cor.-spur scarcely longer than the cal.-appendages.—Rare. Turf-bogs. P. V. VI. E. I.

B. Four upper pet. directed upwards and imbricate. Style clavate. Stigma inflated.

8. *V. hutea* (Huds.); anth.-cells nearly parallel, anth.-spurs long filiform, spur of the cor. as long as or longer than the calycine appendages, sep. acute, l. crenate-serrate lower one ovate-cordate, upper l. ovate or lanceolate, stip. palmate-pinnatifid, terminal lobe linear or linear-lanceolate, st. ascending diffuse and filiform underground.—*Sy. E. B.* 181.—Fl. wholly yellow, yellow with the 2 upper petals purple, or wholly purple, varying greatly in size. Caps. globose. All the lobes of the stip. of nearly

equal size, lateral ones (usually 3 on one side and 1 on the other) all springing from near the base of the stip., the terminal lobe narrow and very nearly always quite entire but sometimes considerably larger than the others.— $\beta$ . *V. Curtisii* (Forst.); stems angular rough, lower part of the stip. somewhat lengthened so as slightly to separate the lateral lobes. *E. B. S.* 2693. *V. sabulosa* Bor.—Mountainous pastures.  $\beta$ . Sands near the west coast. P. VI. VII. E. S. I.

9. *V. tricolor* (L.); *anth.-cells diverging below, anth.-spurs long subclavate-filiform*, spur of the corolla about equalling the calycine appendages, l. crenate-serrate, lower ones ovate-cordate, upper l. ovate or ovate-lanceolate, *stip. lyrate-pinnatifid*, terminal lobe spathulate crenate, st. ascending.—*E. B.* 1287. *R.* iii. 21. *V. Curtisii* (Mack.) from Portmarnock.—Root simple. Fl. with the upper pet. purple, lateral ones bluish, lower one yellow. Caps. ovoid. Terminal lobe of the stip. often having only one tooth on each side.— $\beta$ . *V. arvensis* (Murr.); pet. shorter than the calyx whitish, caps. nearly globular. *E. B. S.* 2712.—Small form from Scilly is very like *V. parvula* Tin.—Common. A. V.—IX. *Heartsease. Pansy.* E. S. I.

### Order X. DROSERACEÆ.

Sep. 5, imbricate. Pet. 5, regular. Stam. 5 or 10, free. Styles 3 or 5. Ovary free. Caps. 3—5-valved; valves bearing the seeds along their middle. Seeds without an aril.—L. with a circinate veneration.—Joined to *Saxifragaceæ* by some authors.

1. DROSERÆ. Cal. deeply 5-cleft. Pet. 5. *Stam.* 5, *hypogynous.* *Styles* 3—5, *deeply bifid.* Caps. 1-celled, with 3—5 valves, many-seeded.

#### 1. DRO'SERÆ Linn. Sundew.

1. *D. rotundifolia* (L.); *l. orbicular* spreading, petioles hairy, fl.-stalks erect from centre of rosette of leaves, seeds with a loose chaffy coat.—*E. B.* 868. *R.* iii. 24.—Flower-stalks 2—6 in. high. Stigmas white, clubbed, entire. Anth. white. L. covered, as in all other species, with hairs terminating in large glands secreting a viscid fluid which retains insects that settle upon them. Rachis recurved parallel to itself.—Common in boggy places. P. VII. VIII. *Round-leaved Sundew.* E. S. I.

2. *D. intermedia* (Hayne); *l. spathulate* blunt erect, petioles glabrous, *fl.-stalks arcuate* or decumbent at the base *from base*

of rosette of leaves, seeds with a close rough not chaffy coat.—*R.* iii. 24. *D. longifolia* Sm.—Stig. pink, bifid. Anth. yellow. Rachis not closely recurved. A variety with shorter leaves and the flower-stalks shorter than the leaves is sometimes found.—Common in boggy places. P. VII. VIII. E. S. I.

3. *D. anglica* (Huds.); l. obovate-lanceolate blunt erect, petioles glabrous, fl.-stalks erect from centre of rosette of leaves, seeds with a loose chaffy coat.—*E. B.* 868. *D. longifolia* *R.* iii. 24, *Koch, Fries.*—Much larger and taller than the last.—β. *D. obovata* (M. and K.); has broader leaves and the styles often, though not always, emarginate. *D. longifolia* Sm. *E. B.* 867.—In bogs, rather rare, common in Ireland. P. VII. VIII. E. S. I.

### Order XI. FRANKENIACEÆ.

Sep. 4 or 5, in a furrowed tube below. Pet. 4 or 5, clawed, with appedages at the base of the limb. Stam. 4 or 5 or more, free, 2-celled, opening by 2 terminal pores or longitudinally. Caps. 1-celled, 2—4-valved; placentas 3, parietal. Style slender, simple or trifid. Seeds many, minute. Embryo in the albumen.—Stip. 0.

1. FRANKENIA. Style 3-fid; lobes oblong with the stigma on their inner side. Caps. 1-celled, 3—4-valved.

#### 1. FRANKENIA *Linn.* Sea-heath.

1. *F. lævis* (L.); l. linear (or rather oblong with reflexed edges) glabrous ciliate at the base.—*E. B.* 205.—St. slightly downy, prostrate, wiry. Cal. slightly hispid between its prominent angles. Fl. terminal or from the forks of the stem, sessile, rose-coloured. L. sometimes pulverulent.—Salt marshes on the East and South coast. P. VIII. E.

[*F. pulverulenta* (L.); l. obovate retuse glabrous above pulverulent beneath, petiole ciliate.—*E. B.* 2222.—Formerly on the Sussex coast, now lost. A. VII.] E.

### Order XII. POLYGALACEÆ.

Sep. 5, imbricate, irregular, 2 interior much larger petal-like. Pet. unequal, usually 3, 1 anterior and larger than the rest. Stam. subdiadelphous, in 2 equal opposite bundles. Anth. 1-celled, opening by a pore at their apex. Caps. 1—2-celled, with placentas in the axis. Seeds pendulous, usually with an aril at the base.—Stip. 0.

1. POLYGALA. Sep. 5, persistent; 2 inner (wings) broader and often petal-like. Cor. irregular. Pet. 3—5, connected together; the lower one keelshaped. Caps. compressed. Seeds solitary, with a 3-pointed basal aril.—Fl. crested.

1. POLYGALA *Linn.* Milkwort.

1. *P. vulgaris* (L.); *l.* scattered, lower *l.* smaller oblong, upper *l.* linear-lanceolate, cal.-wings elliptic mucronate their veins branched the lateral joining a branch of the central vein, caps. obcordate, lateral bracts shorter than the pedicels.—*E. B.* 76.—St. prostrate, ascending. *L.* scattered. Racemes terminal. Fl. blue, pink, or white. Central vein of wings nearly simple ending in a mucro; lateral only branched externally; branches joining in loops and also with the upper ones of the central vein. Lobes of aril unequal, blunt, lateral ones  $\frac{1}{2}$  as long as the seed which is a little stalked within the aril. Upper part of st. and ped. sometimes pubescent (*P. ciliata* Lebel). *P. oxyptera* (R.) is a state with smaller fl., and fr. broader than the wings. *E. B. S.* 2827.— $\beta$ . *P. depressa* (Wend.); lower *l.* mostly opposite and crowded, st. long, prostrate, wiry, racemes ultimately lateral. *Sy. E. B.* 187. *P. serpillacea* Weihe.— $\gamma$ . *grandiflora* (Bab.); upper *l.* large lanceolate, cal.-wings oval apiculate, their lateral veins rejoin the mostly simple central vein near its tip and have many netlike veins externally. Fl. deep blue.—Dry pastures and peaty fens.  $\gamma$ . Ben Bulbin, I. P. VI.—IX. E. S. I.

2. *P. calcærea* (F. Sch.); *l.* chiefly in an irregular terminal rosette large obovate blunt those of fl.-st. short smaller lanceolate, cal.-wings oblong their veins branched the lateral looping with a branch from near the middle of the central vein, caps. oblong-obcordate lateral bracts shorter than the pedicels.—*P. amara* Don, *E. B. S.* 2764. *P. amarella* Coss. et Germ. Atl. t. 7.—St. weak, procumbent or ascending, nearly naked below. Fl.-st. several from the axils of the term. rosette, simple, short; racemes terminal. Fl. blue. Lobes of aril unequal, lateral  $\frac{1}{2}$  as long as seed.—Chalk hills, rare. P. V. E.

3. *P. austriaca* (Cr.); lower *l.* larger obovate blunt in a basal rosette, upper *l.* oblong-lanceolate, cal.-wings oblong or obovate blunt their veins simple or slightly branched free, caps. obcordate broader than the wings, lateral bracts shorter than the pedicels.—Rosette at crown of rootstock, its *l.* much the largest broad rounded at the end. Fl.-st. from the axils of the rosette, simple, short; upper *l.* acute; racemes terminal. Cal.-wings longer than caps. Lobes of aril nearly equal, blunt,  $\frac{1}{2}$  as long as seed.—*a*; caps. rounded at the base, fl. very small. *Sy. E.*

*B.* 189.— $\beta$ . *P. uliginosa* (R.); caps. rather wedgeshaped below, fl. larger.— $\alpha$ . Wye Downs, Kent. *Mr. J. F. Duthie.*  $\beta$ . Cronkley Fell, Yorkshire. *Mr. James Backhouse.* P. VI. VII. E.

### Order XIII. ELATINACEÆ.

Sep. 3—5, distinct, or slightly connate. Pet. 3—5. Stam. as many or twice as many as the pet., free. Caps. 3—5-celled, 3—5-valved, loculicidal, dissepiments adhering to central axis. Styles 3—5; stigmas capitate. Seeds many, albumen 0, embryo curved with the seed.—L. opposite. Stip. minute or inconspicuous.

1. ELATINE. Cal. 3—4-parted. Pet. 3—4. Stam. 3—4 or 6—8. Styles 3—4. Caps. 3—4-celled, many-seeded. Seeds cylindrical, straight or bent.

#### 1. ELA'TINE *Linn.* Waterwort.

1. *E. hexan'dra* (DC.); l. opposite longer than their petioles, fl. slightly stalked with 6 stam. and 3 obovate pet., caps. turbinate concave at the summit 3-celled, seeds nearly straight ascending 8—12 in each cell.—*E. B.* 955. *R. I.* f. 599. *E. Hydro-piper* and *E. tripetala* (Sm.).—Plant minute, creeping. Fl. alternate, axillary. Cal. 3-fid.—Forming small matted tufts under water, rare. A. VIII. E. S. I.

2. *L. Hydro-piper* (L.); l. opposite shorter than their petioles, fl. sessile with 8 stam. and 4 ovate pet., caps. roundish depressed 4-celled, seeds bent almost double pendulous 4 in each cell.—*E. B. S.* 2670. *E. siphosperma Dum.*—Plant minute, creeping. Cal. 4-fid.—Very rare, growing under water. A. VIII. E. I.

### Order XIV. CARYOPHYLLACEÆ.

Sep. 5 or 4, distinct or connected into a tube. Pet. 5 or 4, clawed (rarely 0). Stam. usually twice as many, sometimes as many as the petals, free or connected at the base. Anth. opening longitudinally. Ovary one, often stalked. Stigmas 2—5, sessile, filiform. Caps. 1- or imperfectly 2—5-celled, opening by twice as many teeth as stigmas, sometimes valvular. Placenta central. Embryo generally curved round mealy albumen.—L. opposite, without or rarely with scarious stipules.

#### Suborder I. SILENEÆ.

Sep. connate, forming a tube. Stam. 10. Filaments con-

nate into a tube below and adnate to the stalk of the ovary. Caps. usually stalked.—Pet. 5, clawed in all our plants. No stipules.

\* *Two or more imbricated opposite scales at the base of the calyx.*

1. DIANTHUS. Cal. 5-toothed. Styles 2. Caps. 1-celled, many-seeded, 4-valved at top. Seeds peltate, convex above, concave beneath and more or less keeled.

\*\* *No scales at the base of the calyx.*

2. SAPONARIA. Cal. 5-toothed, terete. Styles 2. Caps. 1-celled, 4-valved at top. Seeds globular or reniform.
- [3. CUCUBALUS. Cal. 5-toothed. Styles 3. Caps. a globose 1-celled berry. Seeds reniform.]
4. SILENE. Cal. 5-toothed. Styles 3. Caps. more or less completely 3-celled, 6-valved at top. Seeds reniform.—(Rarely 5 styles and 5-valved caps. in *S. maritima*.)
5. LYCHNIS. Cal. 5-toothed. Styles 5. Caps. 1- or half 5-celled, opening at the top with 5 or 10 teeth.

#### Suborder II. ALSINEÆ.

Sep. distinct. Stam. free, inserted into a more or less evident hypogynous ring. Caps. sessile. No stipules.

\* *Valves of the caps. the same in number as the styles.*

- [6. BUFFONIA. Sep. 4. Pet. 4, entire. Stam. 4. Styles 2.]
7. SAGINA. Sep., entire pet. (or none), styles and valves of caps. each 4—5. Seeds reniform, wingless. Stam. 4—10.
8. HONKENEJA. Sep. 5. Pet. 5, large. Stam. 10, alternating with glands. Styles and valves 3. Seeds few, large.
9. ALSINE. Sep. and pet. 5 or 4. Styles and valves 3 (or 4). Seeds many, with a naked hile.—L. linear.
10. CHERLERIA. Sep. 5. Pet. 0 or 5, minute. Stam. 10, outer ones opposite to the sep. and springing from an oblong emarginate glandular base. Styles and valves 3. Seeds few, small.



\*\* *Valves of the caps. bifid or twice as many as the styles.*

11. ARENARIA. Sep. 5. *Pet. 5, entire or slightly emarginate.* Stam. 10 or rarely 5. Styles 3. Caps. 6-valved. Seeds many.—L. broad.
12. HOLOSTEUM. Sep. 5. *Pet. 5, jagged at the end.* Stam. 5 or 3 or 4. Styles 3. Caps. subcylindrical, many-seeded, opening at the end with 6 teeth.
13. STELLARIA. Sep. 5. *Pet. 5, bifid.* Stam. 10 (or fewer). Styles 3. Caps. opening with 6 valves or teeth, many-seeded.
14. MALACHIUM. Sep. 5. *Pet. 5, bifid or entire.* Stam. 10. Styles 5. Caps. opening with 5 bifid valves.
15. CERASTIUM. Sep. 5. *Pet. 5, bifid.* Stam. 10 or 5 or 4. Styles 5 or 4. Caps. tubular, opening at the end with 10 teeth.—In *C. trigynum* the styles are mostly 3.
16. MOENCHIA. Sep. 4, erect. *Pet. 4, entire.* Stam. 4. Caps. many-seeded, opening at the end with 8 or 10 teeth.

#### Suborder III. POLYCARPEÆ.

Sep. distinct. Stam. free. Caps. sessile. *Stipules scarious.*

17. POLYCARPON. Sep. keeled at the back, hooded at the end. *Pet. 5, emarginate.* Stam. 3—5. Styles 3, short. Fr. 1-celled, many-seeded.
18. LEPIGONUM. Sep. 5. *Pet. 5, entire, usually as long as the calyx.* Stam. 5—10. Styles 3 or 5. Fr. 3—5-valved, many-seeded; valves fewer than or alternate with the sepals.
19. SPERGULA. Sep. 5. *Pet. 5, entire, as long as the calyx.* Stam. 5—10. Styles 5. Fr. 5-valved, many-seeded; the valves opposite to the sepals.

#### Suborder IV. SCLERANTHEÆ.

Sep. connate, forming a hardened tube enclosing the 1-seeded capsule. No stipules.

20. SCLERANTHUS. Tube of calyx vase-shaped, contracted at the mouth by a glandular ring; limb 5-fid. Pot. 0. Stam. 10 or rarely 5, inserted in the throat of the calyx. Styles 2. Fr. membranous.

Suborder I. *Sileneæ*.

## 1. DIAN'THUS Linn. Pink.

\* *Fl. clustered.*

1. *D. prolifer* (L.); *fl.* in a dense cluster enveloped in membranous bracts, *cal.-scales membranous pellucid* the two outer ones shorter mucronate, inner ones blunt about equalling the calyx, *st.* glabrous, l. all linear, seeds boatshaped with a longitudinal membrane in the hollow rough pointed at one end.—*E. B.* 956. *R.* vi. 247.—*St.* 1—1½ ft. high, erect, usually simple. *Fl.* expanding one at a time, small. *Pet.* rose-coloured, obovate. Cluster quite inclosed by brown dry scales.—Sandy and gravelly places, very rare. A. VII. E.

2. *D. Arméria* (L.); *fl.* close together, *cal.-scales and bracts lanceolate-subulate* herbaceous downy ribbed equalling the tube, *st.* and linear l. downy, seeds nearly flat on one side slightly hollowed and with a longitudinal keel in its middle rough pointed at one end.—*E. B.* 317. *R.* vi. 249.—*St.* 1—2 feet high, erect, branched. *Pet.* rose-coloured, speckled with white dots, toothed.—Waste places, rare. A. VII. VIII. *Deptford Pink.* E.

\*\* *Fl. solitary, one or more on the stem.*

\*3. *D. plumárius* (L.); *st.* 2—5-flowered, *fl. solitary*, *cal.-scales* roundish-ovate shortly mucronate 4-fold shorter than the tube, *l. rough at the edge* linear-subulate, *pet. digitate multifid as far as the middle* with the central entire part obovate downy, barren *st.* procumbent rooting much branched, *seeds flat orbicular* with a point on one side.—*R.* vi. 257. *E. B. S.* 2979.—Flowering-stems 6—12 in. high. Calyx-teeth ciliate at the margin, slightly shorter than the capsule. *Fl.* pale pink, sometimes white, fragrant.—Old walls. P. VI. *Common Pink.* E.

\*4. *D. Caryophyll'lus* (L.); *fl.* solitary, *cal.-scales* broadly obovate pointed three fourths shorter than the tube, *l. with smooth edges* linear, *pet. crenate-dentate* ovate glabrous, barren *st.* elongate procumbent branching, seeds pyriform nearly flat.—*E. B.* 214. *R.* vi. 268.—*Fl.-stems* 12—18 in. high. Calyx-teeth not ciliate, longer than the capsule. *Fl.* pink, fragrant.—Old walls, Kent. P. VII. VIII. *Clove Pink.* E. S. ?

5. *D. cæ'sius* (Sm.); *st.* mostly single-flowered, *cal.-scales* adpressed roundish shortly pointed three fourths shorter than the tube, *l. with rough edges* linear, *pet.* obovate crenately-cut bearded, barren stems long procumbent branching, seeds ovate

pointed at one end.—*E. B.* 62. *R.* vi. 265.—Flowering-stem 6—8 in. high. Fl. pale rose-colour, fragrant.—Limestone cliffs at Cheddar, Som. P. VI. VII. *Cheddar Pink.* E.

6. *D. deltoïdes* (L.); fl. solitary, *cal.-scales* usually 2 ovate with a subulate point  $\frac{1}{2}$  as long as the tube, l. linear-lanceolate the lower blunt rough at the edges and keel, *stem-l.* acute and as well as the st. *pubescent-asperous*, pet. obovate-dentate, barren st. short procumbent simple (?), seeds obovate flat netted-rugose.—*E. B.* 61. *R.* vi. 263.—Flowering-stems 6—12 in. high, branched. Calyx-teeth lanceolate, minutely ciliate. Fl. rose-coloured, with a darker circle round the mouth, scentless.—Hilly pastures, rare. P. VI.—IX. *Maiden Pink.* E. S.

## 2. SAPONA'RIA Linn. Soapwort.

\*1. *S. officinâlis* (L.); fl. in corymbose cymes, cal. cylindrical, pet. retuse crowned, l. elliptic-lanceolate ribbed glaucous, st. erect.—*E. B.* 1060. *R.* vi. 245. *St.* 6. 10.—St. 1—3 feet high, stout, leafy. Fl. flesh-coloured or pale pink, large. Upper l. connate and sheathing.—Hedges near villages, but probably introduced. Banks of streams on the borders of Wales, perhaps indigenous. P. VIII. E.

## 3. CUCU'BALUS Linn.

[*C. bacciferus* (L.); st. branched spreading, l. ovate acute, cal. bellshaped, pet. distant.—*E. B.* 1577.—Fruit fleshy.—Isle of Dogs near London, not native.—P. VIII.]

## 4. SILE'NE Linn. Catchfly.

1. *S. anglica* (L.); *racemes terminal*, fl. alternate, cal. hairy with setaceous teeth ovate when in fruit, pet. slightly cloven or entire obovate, l. lanceolate lower ones spathulate.—*E. B.* 1178.—Hairy and viscid. St. 6—12 in. high, simple or branched, erect. Fl. solitary, secund, axillary, white or reddish. Fr-st. often reflexed.— $\beta$ . *S. quinquevulnera* (L.); fl. white with a large crimson spot on disk of each petal. *E. B.* 86.—Sandy and gravelly fields. About forms of this, see *J. of B.* xviii. 146. A. VI.—X. *English Catchfly.* E. S. I.

2. *S. nitans* (L.); pubescent, glandular-viscid above, *panicle* secund with drooping trichotomous opposite 3—7-flowered branches, *cal. ventricose* with acute teeth, pet. bifid crowned, segments linear, lower l. lanceolate-spathulate, *stem-l.* sessile lanceolate, carpophore scarcely half as long as the capsule, teeth of the caps. reflexed.—*E. B.* 465.—St.  $1\frac{1}{2}$  foot high. Fl. white,

most expanded and sweetest in the evening.—β. *S. paradoxa* (Sm.); rt.-l. roundly spatulate mucronate with long hafts.—On limestone and chalky places. β. Dover Cliffs. P. VI. VII. *Nottingham Catchfly*. E. S.

[*S. italica* (Pers.); pubescent, panicle nearly erect with opposite trichotomous viscid branches, cal. clavate with blunt teeth, pet. bifid not crowned, segments broad, lower l. lanceolate spatulate, stem-l. linear-lanceolate, carpophore as long as capsule.—*S. patens* E. B. S. 2748.—St. 2 feet high.—Not a native. P. VI. VII.] E. S.

3. *S. Otites* (Sm.); panicle long with opposite tufted whorled-racemose branches, whorls many-flowered, ped. glabrous, cal. faintly veined smooth with blunt teeth, pet. linear undivided not crowned, l. lanceolate-spatulate, stem-l. small linear erect.—E. B. 85.—Fl. subdioecious, small, yellowish. Caps. sessile. St. viscid at about the middle, 1 foot high. L. mostly radical.—Sandy and gravelly places in Suff., Norf. and Cambridgeshire. P. VI. E.

4. *S. inflata* (Sm.); panicle terminal, fl. many drooping, cal. inflated netted, pet. deeply cloven scarcely ever crowned, segments narrow, l. elliptic-lanceolate, stem erect.—E. B. 164.—Glabrous, smooth. St. 2—3 feet high. Inflorescence between corymbose and paniced. Pet. white. Cal. inflated especially below, its mouth narrower than its base. Bracts scarious. Branches of panicle unequal. Scented at night. Sometimes *S. puberula* (Jord.); the st. and l. are rough with hairs, and cal. downy.—Fields and roadsides. P. VI.—VIII. *Bladder Campion*. *White-bottle*. E. S. I.

5. *S. maritima* (With.); panicle terminal, fl. few usually solitary erect, cal. inflated netted, pet. crowned, segments broad, l. lanceolate or ovate-lanceolate, st. spreading decumbent, fl.-shoots ascending.—E. B. 957.—Barren procumbent shoots forming a cushion. Fl. larger than those of Sp. 4. Cal. elliptic, its mouth broader than its base, most inflated above the middle. Bracts herbaceous.—Near the sea, also by alpine rills. P. VI.—VIII. E. S. I.

6. *S. conica* (L.); st. erect forked, fl. from the forks or terminal, cal. with 30 furrows conical in fruit, teeth subulate acute, pet. obcordate crowned, l. linear subulate downy, caps. oblong-ovate.—E. B. 922. *S. conoidea* R.—St. 3—12 in. high, simple or branched. Cal. of the flowers conical-tubular, rounded below; of the fruit very broad at the base. Carpophore very short. Fl. reddish.—Sandy fields, rare. A. V. VI. E.

7. *S. noctiflora* (L.); st. erect repeatedly forked, fl. from the forks or terminal, cal. veined and with 10 hairy glandular ribs in fruit elliptic-oblong, teeth long subulate, pet. deeply bifid crowned, l. lanceolate lower ones obovate, caps. ovate.—*E. B.* 291. *St.* 3. 10.—Resembling *Lychnis vespertina*. *St.* about 1 foot high, downy and glandular. Caps. opening with 6 patent teeth. Carpophore very short. Fl. reddish white, rather large, sweet-scented in the evening; pet. rolled up by day; peduncles glandular.—Sandy and gravelly fields, rare. A. VII. VIII. E. S. I.

[*S. Arméria* (L.); pan. forked many-fl. level-topped, pet. notched each with a double awlshaped scale, cal. and l. smooth, caps. clavate, st. viscid.—*E. B.* 1398.—Yalding, Kent. Between Par and Fowey Point, Cornwall. A. VII.] E.

8. *S. acutilis* (L.); st. densely tufted and much branched, fl. solitary, peduncles and cal. glabrous, cal. bellshaped with 10 striæ, teeth ovate blunt, pet. slightly notched crowned, l. linear ciliate below.—*E. B.* 1081. *R.* 5084.—Forming broad dense tufts 2—3 in. high. Fl. purple or white, upon longish solitary stalks, sometimes nearly sessile. Caps. twice as long as the calyx. Plants somewhat dioecious.—On the higher mountains. P. VII. VIII. Cushion Pink. E. S. I.

## 5. LYCHNIS Linn. Campion.

\* *Cal. tubular, not inflated; teeth short.*

1. *L. Viscária* (L.); pet. emarginate crowned, st. glabrous viscid below the joinings, l. lanceolate glabrous the margins woolly at the base, fl. racemose-panicled somewhat whorled, carpophore  $\frac{1}{2}$  the length of the capsule.—*E. B.* 788.—*St.* simple, 1 foot high. Fl. large, rose-coloured. Caps. 5-celled when young. Seeds reniform, minute, acutely tubercled.—Dry rocks, very rare. P. VI. E. S.

2. *L. alpina* (L.); pet. cloven scarcely crowned, st. glabrous not at all viscid, l. linear-lanceolate glabrous sometimes minutely ciliate at the base, fl. corymbose, carpophore  $\frac{1}{2}$  the length of the capsule.—*E. B.* 2254.—*St.* simple, 5—6 in. high. Fl. small, rose-coloured, crown scarcely more than 2 small tubercles upon each petal. Caps. 5-celled when young. Seeds reniform, minute, bluntly tubercled.—Mountains. Glen Isla, Forfar. (3200 ft.). Hobcarten Crag, Cumb. (2000 ft.). P. VI. VII. E. S.

3. *L. Flos-cuculi* (L.); pet. deeply 4-cleft crowned, segments linear palmately diverging, cal. with short teeth, st. with deflexed hairs, l. lanceolate the lower ones narrowed below, fl.

loosely paniced, caps. 1-celled, teeth 5, *carphophore very short*.—*E. B.* 573.—St. viscid and brownish olive, 1—2 feet high. Pet. rose-coloured, the crown bipartite; segm. subulate erect, usually with an acute tooth on the middle of the outer margin. Cal. 10-ribbed.—Moist places. P. V. VI. *Ragged Robin*.

E. S. I.

\*\* *Cal. inflated; teeth falling short of the petals.*

4. *L. vespertina* (Sibth.); *pet. half-bifid* crowned, st. villose, l. peduncles and cal. hairy, l. ovate-lanceolate, fl. dichotomously paniced dioecious, *calyx-teeth of the fertile fl. linear-lanceolate long, caps. conical, teeth 10 straight*.—*E. B.* 1580. St. 23. 9. *L. dioica*  $\beta$  L.—St. 1—2 ft. high. Fl. white, very rarely reddish. Calyx of barren fl. obovate-oblong; of fertile fl. ovate, teeth twice as long as those of *L. diurna*. Carphophore broad, short.—Fields. B. (?) VI.—IX. *White Campion*. E. S. I.

5. *L. diurna* (Sibth.); *pet. half-bifid* crowned, st. l. and cal. villose, l. ovate acute, fl. dichotomously paniced subdioecious, *calyx-teeth of the fertile fl. triangular, caps. nearly globular, teeth 10 recurved*.—*E. B.* 1579. St. 23. 8. *L. dioica*  $\alpha$  L.—Forms a turf of decumbent leafy barren shoots. St. 1—2 ft. high. Fl. red, very rarely nearly white. Carphophore narrow, short. Sp. 4 and 5 vary in colour from red to white and white to red.—Damp hedgebanks. P. V. VI. *Red Campion*. E. S. I.

\*\*\* *Cal. with long leaflike narrow lobes exceeding the petals.*

6. *L. Githágo* (Scop.); *pet. entire or emarginate* crownless, *calyx-teeth longer than the tube exceeding the petals*, fl. solitary upon long stalks.—*Agrostemma* L. *E. B.* 741. St. 5. 6.—Fl. large, purple. St. dichotomous, 2—3 feet high. Cal. coriaceous, ribbed, with 5 linear-lanceolate constantly erect-patent very long segments. Styles downy. Caps. 5-toothed.—Cornfields. A. VI.—VIII. *Corn-Cockle*. E. S. I.

## Suborder II. *Alsineæ*.

### 6. BUFFO'NIA Linn.

[*B. tenuifolia* (Sm.) is said to have been found near Boston, Linc. and on Hounslow Heath. Probably an error. A. VI.]

### 7. SAGI'NA Linn. Pearlwort.

\* *Sep., Stam., styles and valves of caps. 4; pet. very small or wanting.*

1. *S. procumbens* (L.); branches long procumbent from a central rosette, l. linear awned, sep. blunt slightly shorter than

the capsule, *apex of the ped. reflexed* after flowering *ultimately erect*.—*E. B.* 880. *R. v.* 201. *St.* 30. 3.—Glabrous. Central st. never lengthening nor flowering; branches axillary, often rooting, usually with fasciculate leaves. Pet. blunt, often wanting. A fifth part is occasionally added to the fl., in which case it is distinguished from *S. saxatilis* by its cal. spreading when in fruit and *styles reflexed* during flowering. A fleshy, maritime form is *S. maritima* (Gren.).— $\beta$ . *spinosa* (Gibs.); l. longer and narrower very minutely spinose-ciliate on the edges.—Waste ground. *P. V.*—IX. E. S. I.

2. *S. apet'ala* (L.); st. and branches ascending, l. *linear awned*, sep. blunt shorter than the capsule hooded ultimately spreading in the form of a cross, ped. always erect.—*E. B.* 881. *R. v.* 200.—Central st. lengthening flowering and together with the branches erect, never rooting; upper part of st., ped., and cal. often bearing glandular hairs. Pet. very minute, inversely wedge-shaped and truncate. Caps. conical-ovoid, subpeltate below, stalked.— $\beta$ . *prostrata*; *prostrate*, branching from a rosette which lengthens into a stem and flowers.—Walls and dry places.— $\beta$ . Common on gravel walks. *A. V.*—IX. E. S. I.

3. *S. ciliata* (Fr.); st. long, branches diffuse or spreading, l. linear awned, *outer sep. pointed scarcely shorter than and adpressed to the mature caps. their tips patent*.—*Sy. E. B.* 247. *R. v.* 200. *S. patula* Jord.—Glabrous. Central st. flowering and together with the branches erect or ascending, not rooting. Cal. and tips of ped. sometimes with gland-tipped hairs. Caps. ovate-attenuate, rounded below, stalked.—*S. ambigua* (Lloyd) is probably a maritime form.—Dry places and sandy heaths. *A. V.* VI. E. S. I.

4. *S. marit'ima* (Don); *central st. long forked*, branches ascending, l. *fleshy blunt* or apiculate rounded at the back glabrous, sep. blunt about equalling the capsule ultimately spreading slightly, *ped. erect*.—*E. B.* 2195. *S. stricta* Fries.—St. often purple, brittle. The central stem produces flowers and is erect, or in luxuriant plants more or less procumbent. Sep. concave with incurved tips. Caps. ovate, rounded below.—A much-branched prostrate form is *S. debilis* (Jord.); its calyx usually exceeds the caps., and all its stems are often prostrate and spring from a false rosette.—Another very much branched state forming dense tufts, with short joints and shorter linear plane-convex l., is *S. den'sa* (Jord.).—On the seashore. (Fries states that his plant sometimes occurs upon mountains in Norway; and G. Don seems to have found it on Ben Nevis.) *A. V.*—IX. *Sea Pearlwort*. E. S. I.

\*\* *Sep., pet., styles and valves usually 5. Stam. 10. SPERGELLA.*

5. *S. saxat'ilis* (Wimm.); central st. short and barren, l. linear mucronate glabrous, st. ped. and cal. glabrous, pet. shorter than caps. longer than the calyx.—*Spergula saginoides*, L. *E. B.* 2105.—St. prostrate, slightly rooting, many. Ped. long, their tips reflexed after flowering ultimately erect. Caps. rather longer than the calyx, sometimes twice the length. Closely resembling *S. procumbens*, but distinguished by the valves of its capsule being much more narrowed upwards, sep. adpressed and narrower, pet. longer, styles not reflexed.—Highland mountains. P. VI.—VIII. S.

6. *S. nivâlis* (Fries); central st. and branches ascending cæspitose, l. subulate mucronate glabrous, ped. short straight, sep. very blunt adpressed to the ripe capsules, pet. rather exceeding cal. but falling short of caps. entire.—*Sy. E. B.* 250 (bad).—St. and branches dividing repeatedly (no true rosette), not rooting, 1—1½ in. long, forming a dense tuft. Fl. divided in fours or fives. Ped. wholly straight. Sep. white with diaphanous edges.—Tops of Highland mountains, very rare. P. VIII. S.

7. *S. subulâta* (Wimm.); l. awned linear often ciliate, ped. and calyx glandular-hairy, pet. about as long as the caps. longer than the calyx.—*Spergula* Sw. *E. B.* 1082.—St. procumbent. Ped. very long, the tip slightly reflexed after flowering, ultimately erect. Caps. ovate-attenuate, rounded below, sessile.—Dry gravelly and sandy places. P. VI.—VIII. E. S. I.

8. *S. nodôsa* (E. Meyer); l. subulate glabrous upper l. shorter fasciculate, pet. much longer than the calyx, ped. always erect.—*Spergula* L. *E. B.* 694.—Primary stem short, not flowering; lateral stems procumbent at the base then ascending, 2—6 in. long. Fl. terminal, 1, 2 or 3 together, white, conspicuous. Whole plant often quite glabrous. Sometimes (*S. glandulosa* Bess.) the upper parts of the st., the connecting membrane of the l., and the base of the cal. are glandular-hairy.—Wet and sandy places. P. VII. VIII. Knotted *Spurrey*. E. S. I.

### 8. HONKENE'JA Ehrh.

1. *H. peploïdes* (Ehrh.); l. sessile ovate acute fleshy glabrous 1-veined, pet. obovate, sep. ovate blunt 1-veined shorter than the petals.—*Arenaria* L. *Ammodenia* Gm. (name only). *E. B.* 189.—St. forked, procumbent, rhizomatous. Fl. from the forks of the stem, frequently diœcious or polygamous. Caps. large, globose. Seeds few, large.—Sandy sea-coasts. P. VI.—IX. E. S. I.



9. ALSI'NE *Wahl.*

1. *A. stricta* (Wahl.); *l.* filiform *veinless*, fl.-shoots erect naked above, pet. equalling the cal. oblong-oval attenuate below, *sep.* ovate-lanceolate *acute* 3-veined (when dry), *ped. terminal* 1—3 *very long*.—Aren. uliginosa *E. B. S.* 2890.—St. prostrate, cæspitose.—Widdy-bank Fell, Teesdale. P. VI. E.

2. *A. ver'na* (Jacq.); *l.* linear-subulate *acute* 3-veined, *pet. exceeding the calyx* rounded-obovate attenuate below, *sep.* ovate-lanceolate *acute* 3-veined with a membranous margin, peduncles 1- or many-flowered.—*E. B.* 512.—St. 3—4 in. high. L. usually not adpressed and mostly with a minute point. Bracts *acute*.— $\beta$ . *A. Gerardi* (Willd.); *l.* subulate bluntish not apiculate, *pet.* elliptic shortly clawed scarcely longer than the calyx. *R.* v. 208. L. usually pressed close to the stem.—Rocky places in mountainous districts.  $\beta$ . On the hills above Kynance Cove near the Lizard Point, Cornwall. P. V.—IX. E. S. I.

3. *A. rubel'la* (Wahl.); *l.* linear-subulate *blunt* 3-veined, *pet.* obovate attenuate below *shorter than the calyx*, *sep.* ovate-lanceolate *acute* 3-veined with a membranous margin, peduncles 1-flowered.—*E. B. S.* 2638.—Like a *Sagina*. St. many, short. Bracts blunt. Flowering shoots terminal, downy, nearly always 1-fl., about 1 in. long, with 1—3 pairs of leaves. Ped. longer than calyx. Styles and valves of caps. 3—5.—Tops of Scottish mountains, very rare. P. VII. VIII. S.

4. *A. tenuifolia* (Wahl.); *l.* subulate-acute 3-veined, *pet. ovate attenuate below shorter than the calyx*, *sep.* lanceolate-subulate 3-veined with a membranous margin.—*E. B.* 219.—St. slender, 4—6 in. high, much branched, forked, with flowers in the forks.—Glabrous. Sometimes the upper parts (*A. laxa*, Jord.), or the cal. alone (*A. hybrida* Jord.), bear patent gland-tipped hairs.—Sandy and chalky places, rare. A. V. VI. E.

10. CHERLE'RIA *Linn.*

1. *C. sedoides* (L.).—*E. B.* 1212.—Pet. generally wanting. Fl. solitary, on short stalks. St. very many, forming a dense mass close to the ground. L. very many, linear-subulate, finely ciliate.—Summits of mountains. P. VI.—VIII. S.

11. ARENA'RIA *Linn.* Sandwort.

1. *A. Triner'via* (L.); *l.* ovate acute ciliate stalked 3—5-veined the upper ones sessile, *stam.* 10, *pet.* shorter than the calyx, *sep.* long-lanceolate acute 3-ribbed the intermediate rib strongest and rough, seeds smooth.—*E. B.* 1483. *R.* v. 216.—St.

about a foot high, weak, branched, downy Fl. solitary from the forks of the stem and axils of the upper leaves. Ped. ultimately spreading and curved just below the fruit. Lateral veins of sep. often very faint. Distinguished from our other species by the appendage to the hile of its seeds.—Damp shady places. A. V. VI. E. S. I.

2. *A. serpyllifolia* (L.); *l. ovate* acute roughish sessile, *pet. shorter than the calyx, sep. ovate-lanceolate* acute 3—5-veined hairy on the veins, *fr.-st. erect* or patent straight longer than the *ampullaceous caps.* which exceed the sepals.—*E. B.* 923.—St. much branched, 3—6 in. long. Fl. from the forks of st. or axils of leaves. Pet. ovate, narrowed below. Ripe capsule brittle.—I cannot distinguish *A. Lloydii*.—Dry places and walls. A. VI.—VIII. E. S. I.

3. *A. leptoclados* (Guss.); *l. small ovate* acute sessile, *pet. shorter than the cal., sep. lanceolate* acute 3-veined hairy on the veins, *fr.-st. patent curved at the top* or ultimately straight longer than the *ovoid-oblong caps.* which exceed the sep.—*E. B. S.* 2972.—St. much branched. Fl. as in Sp. 2. Much more slender than Sp. 2; caps. small and often nearly oblong flexible when ripe; seeds smaller.—Dry places and walls. A. VI.—VIII. E. I.

4. *A. ciliata* (L.); *l. spatulate ciliate, pet. exceeding the calyx,* *sep. ovate-lanceolate* with 3 prominent ribs.—*E. B.* 1745.—St. long, much branched, prostrate, rough, with very short deflexed hairs, angular when dry. Fl. 1—5, terminal, somewhat paniced. Pet. ovate, slightly clawed.—Limestone cliffs of King Mountain, co. Sligo. P. VI. VII. I.

5. *A. norvégica* (Gunn.); *l. spatulate obovate fleshy not ciliate,\** *pet. exceeding the calyx, sep. ovate-acute obscurely 3-ribbed* glabrous.—*E. B. S.* 2852.—St. short, much branched, procumbent, nearly smooth, angular when dry. Fl. 1—3, terminal. Ped. with very short deflexed hairs. Pet. ovate, slightly clawed. Seed dark brown, tuberculated.—Unst, Shetland. P. VII. VIII. S.

## 12. HOLOS'TEUM Linn.

1. *H. umbellatum* (L.); fl. umbellate, peduncles pubescent viscid, pedicels reflexed after flowering, l. elliptic or long and acute.—*E. B.* 27. *R. v.* 221.—About 6 in. high.—On old walls and dry places in Norfolk and Suffolk. A. IV. E.

## 13. STELLARIA Linn. Stitchwort.

\* *Seeds on a linear long columella.*

1. *S. nem'orum* (L.); st. ascending downy above, l. stalked heartshaped, upper l. ovate sessile, cyme lax paniced, pet. deeply bifid twice as long as the lanceolate sepals, caps. exceeding the calyx.—*E. B.* 92. *R. v.* 252.—St. 1—1½ foot high. L. large, rough on the upper surface, ciliate.—Sep. with narrow scarious margins.—Damp woods, chiefly in the North. P. V. VI. *Wood Stitchwort.* E. S.

\*\* *Caps. rounded below or scarcely if at all narrowed; columella very short.*

2. *S. média* (Vill.); st. procumbent and ascending with a hairy line, l. ovate shortly pointed stalked, upper l. sessile, fl. axillary and terminal, pet. deeply bifid not exceeding the ovate-lanceolate single-ribbed glandular-pilose sepals, caps. oblong longer than the calyx.—*E. B.* 537. *R. v.* 222.—Very variable in length of st. and joints, size of l., number of stam. (3—10), and length of styles. Sep. with a narrow scarious margin, glabrous or with long hairs. Fr.-st. reflexed, often scarcely exceeding the leaves. Seeds with round tubercles. L. glabrous with broad ciliate petioles.— $\beta$ . *Alsine pallida* (Dum.); pet. 0, styles 0, stig. short arcuate, stam. 3, seeds small bluntly and minutely tubercled.  $\mathcal{S}$ . *Borœana* (Jord.).— $\gamma$ . *S. neglecta* (Weihe); l. larger with longer stalks, upper l. sessile lower subcordate, stam. 10, seeds with prominent rounded tubercles.— $\delta$ . *S. umbrosa* (Opitz); l. narrowed gradually into long points, cal. more narrowed below, sep. lanceolate acute tubercular, valves of the caps. narrower, seeds with prominent acute tubercles. St. much branched, slender, with autumnal barren shoots. Fr.-st. much exceeding the leaves. *S. Elizabethæ*, Schultz.—Common. A. ( $\delta$  P.) III.—IX. *Common Chickweed.* E. S. I.

3. *S. Holos'tea* (L.); st. ascending angular with rough angles, l. lanceolate-attenuate acute with a rough margin and keel all sessile, cyme paniced, pet. half-bifid twice as long as the lanceolate very obscurely 3-veined sepals, caps. globose about as long as the calyx, bracts leaflike.—*E. B.* 511. *R. v.* 223.—St. 1—2 feet high, slender and procumbent at the base, thicker upwards. L. gradually narrowing from a little above the base to the very acute point. Fl. large, white, few, in a leafy cyme.—Woods and hedges. P. IV.—VI. *Greater Stitchwort.* E. S. I.

4. *S. glauca* (Wither.); st. erect weak angular smooth l. linear-lanceolate acute quite smooth sessile, lower l. broader, fl. solitary

or in a few-flowered lax paniced cyme, pet. bipartite exceeding the lanceolate 3-veined sepals, caps. oblong-ovate about as long as the calyx, *bracts with scarious and glabrous margins*.—*E. B.* 825. *R.* v. 223. *S. palustris* Retz.—Usually glaucous. St. 6—12 in. high, leafy. Fl. rarely solitary. Pet. white, sometimes much exceeding the cal.; segments linear.—Marshy places, rather rare. P. V.—VII. E. S. I.

5. *S. gramin'ea* (L.); st. diffuse angular smooth, l. linear-lanceolate acute quite *smooth* ciliate below *sessile*, cyme lax paniced, pet. bipartite equalling or exceeding the 3-veined sepals, caps. oblong longer than the calyx, *bracts scarious ciliate*.—*E. B.* 803.—St. 1—2 feet high. Fl. smaller than those of Sp. 3 or 4, white. Shorter or longer pet. accompany an imperfection of the stam. or germen.—[*S. longifolia* (Fr.), *S. Friesiana* (Koch), has the upper part of its stem and the edges and keel of its leaves rough.]—Dry heathy and bushy places. P. V.—VIII. *Lesser Stitchwort*. E. S. I.

\*\*\* *Caps. narrowed below; hence the cal. has a funnelshaped base.*

6. *S. uligin'osa* (Murr.); st. diffuse angular *glabrous*, l. oblong-lanceolate acute with a hard tip *glabrous slightly ciliate below*, fl. irregularly paniced lateral and terminal, pet. bipartite shorter than the lanceolate 3-veined sepals, caps. ovoid nearly equalling the calyx, *bracts scarious with glabrous edges*.—*E. B.* 1074.—Very variable in size, from about a foot long to 2 inches. Fl. in small cymes.—In wet places. P. V. VI. E. S. I.

#### 14. MALA'CHIUM *Fries.* Great Chickweed.

1. *M. aquat'icum* (Fr.); st. decumbent and ascending angular covered with glandular hairs, l. cordate-ovate acuminate, fl. scattered solitary in the forks of the stem, pet. bipartite rather exceeding the calyx, caps. exceeding the calyx.—*R.* vi. 237. *Cerastium* L. *E. B.* 538.—Close resembling *Stellaria nemorum*. Lowest l. and those of the barren branches stalked, others larger and sessile.—Usually in wet places. P. VII. VIII. E.

#### 15. CERAS'TIUM *Linn.* Mouse-ear.

\* *Root fibrous. Pet. not or but little exceeding the calyx.*

† *Caps. curved.*

1. *C. glomerátum* (Thuil.); l. ovate, *sep. lanceolate very acute with a narrow membranous margin and as well as the herbaceous bracts hairy throughout*, caps. cylindrical ascending twice as long as the calyx, fruitstalks about equalling the calyx.—*C. vulgatum* Sm. *E. B.* 789. *R.* v. 229. *C. viscosum Fries.*—St. erect,

grandular-hairy. Fl. in close cymes, longer than their stalks.— $\beta$ . *C. apetalum* (Dum.); pet. 0, whole plant usually much more slender.—Fields and banks. A. IV.—IX. E. S. I.

2. *C. triviale* (Link); l. oblong-lanceolate, sep. oblong-ovate bluntish and as well as the bracts membranous at their margins and glabrous tips, caps. cylindrical ascending twice as long as the calyx, fruitstalks at least as long as the calyx.—*R.* v. 229. *C. viscosum* Sm. *E. B.* 790. *C. vulgatum* Fries.—St. downy, mostly procumbent, some short and barren. Fl. larger than those of the last, in small terminal cymes the branches of which are often lengthened as the fr. ripens, sometimes 5-androus.— $\beta$ . *C. holosteoides* (Fries); glabrous, sides of st. alternately downy, l. dark smooth shining, fr. much larger. *St.* 63. 9.—In fields, and on mountains (*alpestre*) with larger flowers.  $\beta$ . Near Newcastle on Tyne; and Perth. B. or P. IV.—IX. E. S. I.

†† Caps. nearly straight.

3. *C. semidecan'drum* (L.); l. broadly ovate, sep. lanceolate broadly membranous at their margins and tips, bracts with their upper half membranous, caps. cylindrical slightly inflated longer than the calyx, fruitstalks longer than the calyx at first reflexed afterwards erect.—*E. B.* 1630. *R.* v. 228.—St. erect or decumbent, downy, sometimes viscid. *Pet.* with simple veins, not distinctly notched. Lower l. with long linear hafts. Known by its half-membranous bracts.—Common in dry places. A. IV. V. E. S. I.

4. *C. pumilum* (Curt. !); l. spatulate, upper l. oblong, sep. lanceolate acute with their tips and margins narrowly membranous, uppermost bracts with an extremely narrow membranous margin, caps. slightly curved upwards longer than the calyx, fruitstalks short curved at the top declining from their base, ultimately erect.—*Curt.* ii. 92. *C. glutinosum* Fr., *Fl. Dan.* 2537.—Viscid. St. branched at the root, afterwards nearly simple. *Pet.* with branched veins, notched. Lower l. with long linear hafts. Fl. in terminal forked cymes.—Dry banks in the south. A. IV. V. E.

5. *C. tetrandrum* (Curt.); l. oval or oblong, sep. lanceolate acute, their tips and margins narrowly membranous, bracts wholly herbaceous, caps. a little exceeding the cal. straight, fruitstalks 2—4 times as long as the caps. straight, ultimately erect.—*E. B.* 166. *Curt.* ii. 93. *C. pumilum* Gren., Bor. *C. atrovirens* and *C. tetrandrum* Bab.—Viscid. St. cymose from the base. Bracts very broad, oval, rather acute or apiculate, or nearly round, leaflike. *Pet.* with branched veins, notched.

Fl.-whorls of 4 or 5 parts.—Walls and sandy places near the sea. A. V.—VII. E. S. I.

\*\* *Root truly perennial, with prostrate leafy shoots. Pet. much longer than the calyx.*

6. *C. arven'se* (L.); st. ascending prostrate below, l. linear-lanceolate, fl. many, sep. and bracts lanceolate slightly acute with membranous margins and tips, caps. at last longer than the calyx, seeds small acutely tubercled.—*E. B.* 93.—St. long. Fl. 3—14, in forked panicles. Fruitstalks erect, bent just under the calyx. St. and l. hairy.—The var. *Andrewsii* (Syme), which has subglabrous rigid l. with a strong midrib and is often 1-flowered, I have not seen.—In sandy, gravelly and chalky places, rare. P. IV.—VIII. E. S. I.

7. *C. latifolium* (L.); pubescence short, st. prostrate cæspitose, l. elliptic or lanceolate, fl. 1—3, sep. blunt with membranous margins, bracts herbaceous, caps. slightly narrowing straight, seeds large rugose, fruitstalks oblique patent.—*E. B.* 473.—Pubescence short, rigid, yellowish. Barren shoots usually long. L. variable in shape.— $\beta$ . *compactum* (Syme); l. roundish-ovate light green, st. short not densely leafy below.— $\gamma$ . *Edmonstonii* (Wats.); l. roundish-ovate dark greenish purple, st. short densely leafy below.—Alpine parts of Wales and Scotland.  $\beta$ . Braemar.  $\gamma$ . Unst, Shetland. P. V. E. S.

8. *C. alpinum* (L.); hairy, st. ascending, l. ovate ovate-oblong or lanceolate, fl. few, sep. bluntish with membranous margins, bracts herbaceous their margins often narrowly membranous, caps. nearly cylindrical curved at the end, seeds small acutely tubercled, fruitstalks obliquely patent.—*E. B.* 472.—Pubescence long, simple. St. much branched below, then simple, elongated, prostrate or ascending. Fl. 1, 2 or 3 together, in a forked panicle, shorter than their stalks.—Alpine parts of Scotland and the North of England. P. VI.—VIII. E. S.

9. *C. trigynum* (Vill.); st. decumbent with an alternate hairy line, l. elliptic-oblong, ped. pubescent 1—3-flowered terminal, bracts herbaceous, styles mostly 3, caps. rather longer than the calyx.—*Stellaria cerastoides* L. *E. B.* 911.—St. 4—8 in. long, slender, leafless and much branched below. L. light green, glabrous (or hairy in *C. nivale* Don), subsecund and subfalcate. Fl. large, white. Teeth of caps. 6—10.—Highland mountains. P. VII. VIII. S. I.

#### 16. MOENCHIA Ehrh.

1. *M. erecta* (Sm.); stam. 4.—*E. B.* 609. *R.* v. 227.—

Glaucous. St. erect, glabrous, 1—4 in. high. L. opposite, linear-lanceolate, acute, rigid. Sep. with broad white membranous margins, acute.—Dry gravelly and sandy places. A. V. VI. E. I.

### Suborder III. *Polycarpeæ*.

#### 17. POLYCAR'PON *Linn.*

1. *P. tetraphyllum* (L.); triandrous; pet. emarginate, stem-l. in fours, l. on the branches opposite.—*E. B.* 1031.—In young plants the l. are often all opposite.—Coasts of the South-west of England, rare. A. VI. VII. E.

#### 18. LEPI'GONUM *Fries*<sup>1</sup>. Sand-Spurry.

1. *L. rubrum* (Fr.); st. nearly terete, l. flat linear pointed, stip. triangular-ovate-prolonged mostly cut, caps. about equaling the cal.  $\frac{1}{2}$ — $\frac{2}{3}$  shorter than the fr.-stalk, seeds cuneate-obovate  $\frac{2}{3}$  surrounded by a thickened border none winged.—*Sy. E. B.* 254.—St. procumbent (as in the other species). Pan. leafy. Pet. pink, about equalling the calyx. Seeds usually gibbous on one side. Smaller in all respects than the other species.—Sandy places. A. V.—IX. E. S. I.

2. *L. rupicola* (Kindb.); st. terete, l. flattish fleshy pointed, stip. broadly ovate-prolonged mostly entire, caps. large equaling or slightly exceeding the cal.  $\frac{1}{2}$ — $\frac{2}{3}$  shorter than the fr.-stalk, seeds compressed pyriform nearly surrounded by a thickened border none winged.—*E. B. S.* 2977.—Root thick, woody. *L. fascicled*, short; pan.-l. very short, inconspicuous. Pet. pale pink. Stam. 10.—Near the sea. P. VI.—IX. E. I.

3. *L. salinum* (Kindb.); st. compressed, l. flattish fleshy bluntish or slightly pointed, stip. broadly triangular-ovate entire, caps. exceeding the cal. about as long as the fr.-stalk, seeds compressed roundish nearly surrounded by a thickened border and some often with a broad scarious wing.—*E. B. S.* 2978.—L. long; pan.-l. sometimes long, sometimes inconspicuous. Ped. occasionally twice as long as the capsule. Pet. pink with a white base. Stam. less than 10.—*L. medium* (Fr.) has shortly ovate stip., caps. about equaling cal., seeds triquetrous smooth

<sup>1</sup> *Lepigonum* Fr. Fl. Hal. (1818). *Spergularia* Presl, Fl. Cech. (1819). *Spergularia* (as a section of *Arenaria*) Pers. Syn. (1805). Therefore *Lepigonum* is the oldest generic name.

with a thickened border. *L. neglectum* (Kindb.) is glandular.—Near the sea. A. VI.—IX. E. S. I.

4. *L. marimum* (Wahl.); st. compressed, l. fleshy bluntish, stip. broadly triangular usually entire, caps. very large often twice as long as the cal. but scarcely  $\frac{1}{2}$  as long as the fr.-stalk, seeds compressed roundish nearly surrounded by a thickened border within a broad scarious wing.—*E. B.* 958.—L. long,  $\frac{1}{2}$ -terete; pan.-l. very short, inconspicuous. Caps. larger than in the other species. Seeds reddish. *Pet.* pale pink, with a white base. Stam. 10.—Muddy salt marshes. P. VI.—IX. E. S. I.

### 19. SPERGULA Linn. Spurry.

1. *S. aven'sis* (L.); l. linear convex above furrowed beneath, fl. in forked panicles, fr.-stalks deflexed, seeds slightly compressed with a narrow margin tubercled or papillose.—St. 6—12 in. high. L. long, slender. Seeds black.—*a. S. sativa* (Boen.); seeds margined and covered with minute elevated points, l. grey-green very viscid. *E. B.* 1535.—*β. S. vulgaris* (Boen.); seeds obscurely margined not winged covered with clubshaped papillæ, l. grass-green scarcely viscid.—Cultivated land. A. VI.—VIII. E. S. I.

### Suborder IV. *Scleranthææ*.

#### 20. SCLERANTHUS Linn. Knapwell.

1. *S. an'nuus* (L.); subdecandrous, *segm. of fr.-cal. patent acute* with a very narrow membranous margin, as long as their tube.—*E. B.* 351.—Styles exceeding the stamens. St. repeatedly dichotomous, green. Fl. green, often solitary in the forks of the stem, or densely corymbose.—A biennial state (*S. biennis* Reut.) is often taken for Sp. 2.—Sandy fields. A. VI.—VIII. E. S. I.

2. *S. peren'nis* (L.); decandrous, *segm. of fr.-cal. connivent blunt rounded* with a broad blunt membranous margin.—*E. B.* 352.—Styles usually falling short of the stamens. St. nearly simple or irregularly branched, procumbent, glaucous, at length reddish. Fl. variegated with green and white. L. erect, directed to one side.—Sandy fields in Norfolk and Suffolk. Stanner rocks, Radnorshire. P. VI.—VIII. E.



## Order XV. MALVACEÆ.

Sep. 5 or 3 or 4, more or less connected below, often double, valvate in the bud. Pet. as many as the sepals, adnate to base of stam., twisted in the bud. Stam. many, connected at the base into a tube; anth. 1-celled, reniform, bursting transversely. Ovary formed by the union of several carpels round a common axis. Carp. 1- or many-seeded. Embryo curved with twisted or doubled cotyledons, albumen variable in quantity.—L. alternate, with deciduous stipules. Fl. axillary.

1. MALVA. Styles many. Cal. double, *outer 3-leaved*, inner 5-fid. Carpels in a ring round a thick axis, each 1-seeded.
2. ALTHÆA. Styles many. Cal. double, *outer 6—9-fid*, inner 5-fid. Carpels as in *Malva*.
3. LAVATERA. Styles many. Cal. double, *outer 3-lobed*, inner 5-fid. Carpels as in *Malva*.

## 1. MAL'VA Linn. Mallow.

1. *M. moschata* (L.); st. erect, l. palmate with 5—7 deep bipinnatifid lobes, lower l. incise-crenate, stipules lanceolate acute, fruitstalks erect, outer sep. linear-lanceolate, *fruit hairy*.—*E. B.* 754. *R.* v. 169.—Fl. large, rose-coloured, on axillary single-flowered peduncles, crowded at the extremity of the stem and branches. Cal. hairy. St. 1—2 feet high.—Gravelly places. P. VII. VIII. *Musk Mallow*. E. S. I.

[*M. Alcæa* (L.); outer sep. ovate acute, fr. glabrous; should be found in England.]

2. *M. sylvestris* (L.); st. erect, l. palmate with 5—7 deep crenate lobes, stipules lanceolate, fruitstalks erect, outer sep. lanceolate, *fruit reticulate-rugose* its axis an acute cone with concave sides.—*E. B.* 671. *R.* v. 168.—Fl. large, in axillary clusters. Pet. much longer than the hairy calyx, purple. St. 2—4 feet high.—Roadsides and waste places. P. VI.—IX. *Common Mallow*. E. S. I.

3. *M. rotundifolia* (L.); st. decumbent, l. roundish-heartshaped with 5—7 shallow acutely crenate lobes, stipules ovate acute, fruitstalks reflexed, *outer sepals* linear-lanceolate *shorter than the ovate-acuminate stellately hairy inner ones*, pet. 2 or 3 times as long as the calyx, fruit pubescent, carp. rounded on the edge smooth.—*E. B.* 1092. *M. vulgaris* Fries, *R.* v. 167.—Fl. small, purple. Carp. meeting each other with a straight line. Axis

of the fr. rather large.—Waste places. P. VI.—IX. *Dwarf Mallow*. E. S. I.

\*4. *M. borealis* (Wallm.); outer as long as the glabrous but ciliate inner sepals, pet. scarcely exceeding cal., carp. margined netted rugose.—*E. B.* 241. *M. rotundifolia* Fr.—Carp. meeting each other with a toothed edge. Axis of fr.  $\frac{1}{2}$  as long as in Sp. 3. Much resembling Sp. 3.—Hythe, Kent; now lost. Anthony, Corn. Near London. A. VII. ? E.

## 2. ALTHE'A Linn.

1. *A. officinalis* (L.); l. soft on both sides crenate or crenate-serrate cordate or ovate 3—5-lobed, ped. axillary many-fl. shorter than the leaves, st. downy.—*E. B.* 147. *R.* v. 173.—St. 2—3 feet high. Covered with soft velvety pubescence. Outer cal.-lobes acute spreading much exceeding inner adpressed calyx.—Marshes, particularly near the sea. P. VIII. IX. *Marsh-Mallow*. E. S. ? I.

†2. *A. hirsuta* (L.); hispid cordate, lower l. reniform bluntly 5-lobed, upper l. palmate with 5 or 3 acute lobes, ped. axillary 1-flowered longer than the leaves, st. hispid.—*E. B.* S. 2674. *R.* v. 172.—Stem and calyx very hispid.—Between Cobham and Cuxton, Kent. A. VI. VII. E.

## 3. LAVATE'RA Linn. Tree-Mallow.

1. *L. arborea* (L.); st. woody, l. 7-angled plaited velvety, ped. axillary clustered 1-flowered shorter than the petioles.—*E. B.* 1841.—Fl. large, purplish rose-coloured with darker veins. St. 3—8 feet high.—Maritime rocks, rare. B. VII.—IX. E. S. I.

\*2. *L. sylvestris* (Brot.); st. herbaceous erect or ascending hispid, lower l. orbicular-cordate, upper l. 5-lobed truncate below, l.-lobes triangular acute, ped. axillary 1-fl. shorter than petioles, fr. smooth its axis a cone with convex sides.—*J. of B.* xv. t. 191.—Fl. rose-purple. St. rarely prostrate. Much like *Malva sylvestris*.—Scilly Isles. A. or B. VI. VII. E.

## Order XVI. TILIACEÆ.

Sep. 4—5, valvate in the bud. Pet. 4—5. Stam. many, distinct, or slightly connected into bundles at the base; anth. 2-celled, bursting longitudinally. Glands 4—5 at the base of the petals. Fr. 4—10-celled, several seeds in each cell; or by abortion 1-celled 1-seeded. Embryo erect in the axis of fleshy albumen; cotyledons flat, leafy.—L. alternate, with stipules.

1. *TILIA*. Sep. 5, deciduous. Pet. 5, with or without a scale at the base. Stam. many, free or polyadelphous. Ovary globose, 5-celled, cells 2-seeded. Style 1. Fr. 1-celled, with 1 or 2 seeds.—No scale to the pet. in our plants.

1. *TIL'IA* *Linn.* Lime tree.

‡1. *T. intermèdia* (DC.); *l.* obliquely cordate *glabrous* except woolly tufts at the branching of the veins beneath, ped. many-flowered, *ripe fr.* woody *not ribbed downy*.—*E. B.* 610.—Fl. in a naked cyme springing from a lanceolate leaflike bract. *L.* thin membranous, light transparent green, *twice the length of their petioles*.—In many old plantations. T. VII. E. S.

2. *T. parvifolia* (Ehrh.); *l.* obliquely cordate *glabrous* except woolly tufts at the branching of the veins beneath, ped. many-flowered, *fr. oblique angular thin and brittle*.—*E. B.* 1705.—*L.* thick, coriaceous, opaque above, *usually scarcely longer than their petioles*, with stellate hairs beneath. Lobes of the stigma ultimately spreading horizontally.—In old woods. T. VIII. E.

†3. *T. grandifolia* (Ehrh.); *l.* obliquely cordate *downy beneath* with woolly tufts at the branching of the veins beneath, ped. mostly 3-flowered, *ripe fr. with 5 prominent angles* woody downy turbinate.—*E. B. S.* 2720.—Young shoots hairy. *L.* thin, membranous, bright transparent green, longer than their petioles, with solitary hairs beneath. Lobes of the stigma erect.—*T. rubra* (Lindl.) is stated to have smooth fruit and to be *T. corallina* Sm. I have not seen it.—Old and rocky woods by the Wye, Teme and Severn. T. VI. VII. E.

Order XVII. HYPERICACEÆ.

Sep. 4—5, distinct or cohering, persistent, with glandular dots, imbricate. Pet. 4—5, twisted in the bud. Stam. many, connected in 3 or 4 bundles at the base. Anthers versatile. Styles several, rarely connate. Fruit a dry or fleshy capsule of many cells and many valves, the valves curved inwards. Seeds small, many, on a central axis or the incurved margins of the valves, embryo straight with no albumen.—*L.* mostly opposite, with pellucid dots. Stip. 0. Fl. yellow.

1. *HYPERICUM*. Cal. 5-parted or of 5 sepals. Pet. 5. Styles 3 (in nearly all our plants) or 5. Caps. more or less perfectly 3-celled, many-seeded.—Fl. yellow.

1. *HYPERICUM* Linn. St. John's Wort.\* *Styles 5. Pet. unequal-sided.*

\*1. *H. calycinum* (L.); *st. shrubby* square, l. oblong, fl. solitary, sep. unequal obovate blunt.—*E. B.* 2017.—Rootstock creeping. St. 1 ft. high. Fl. 3 or 4 in. across, yellow, as in all of this genus.—Naturalized in bushy places. P. VII.—IX.  
E. S. I.

\*\* *Styles 3, stam. in 5 sets, pet. deciduous.* ANDROSÆMUM.

2. *H. Androsæmum* (L.); *st. shrubby* compressed, l. broadly subcordate-ovate blunt, cymes trichotomous few-flowered, sep. broad unequal, pet. oval blunt, *styles much falling short of stam., caps. pulpy* imperfectly 3-celled blunt.—*Sy. E. B.* 264. *Curt.* i. 164.—St. very little branched, 2 ft. high. L. large, with a strong aromatic smell when rubbed. Pet. short. Styles much shorter than the black capsule, finally hooked.—Thickets and hedges. P. VII. VIII. *Tutsan.* E. S. I.

[*H. elatum* (Ait.); *st. shrubby*, 2-edged, *ped. 2-winged*, l. subcordate-ovate subacute, cymes few-flowered, sep. broad unequal  $\frac{1}{2}$  as long as petals persistent, *styles exceeding the stam., caps. oval.*—*Wats. Dendr. Brit.* 85. *H. Androsæmum* Sm. *E. B.* 1225. *H. anglicum* Bert.—St. 3—4 ft. high, much branched. Fl. large, in terminal cymes. Sep. nearly as large as those of *H. Androsæmum*. Pet. about equalling stamens.—The remains of cultivation. P. VII.—IX.] E.

[*H. hircinum* (L.), *Wats. Dendr.* 86, *Sy. E. B.* 246, has lanceolate acute deciduous sepals, but otherwise much resembles *H. elatum*; it is established in some places.]

\*\*\* *Herbs. Styles 3, stam. shortly united in 3 sets, pet. persistent.*

3. *H. tetrapterum* (Fr.); *st. erect with 4 wings*, l. ovate with pellucid dots and veins, *sep. erect lanceolate acuminate entire*, pet. lanceolate, styles half as long as the capsule.—*R.* vi. 344. *H. quadrangulum* Sm. *E. B.* 370.—St. 1—2 ft. high. Fl. in terminal forked close many-flowered cymes, pale.—In wet places. P. VII. E. S. I.

4. *H. undulatum* (Schousb.); *st. erect branched 4-edged*, l. oblong wavy at the edge with many pellucid dots netted with pellucid veins and (as well as the sep. and edges of the stem) with black marginal dots beneath, *sep. erect ovate-lanceolate acute usually bluntly crenate*, styles not  $\frac{1}{2}$  as long as the capsule.—*J. of B.* ii. 97. t. 16. *H. bæticum* Boiss. *Voy.* t. 34. *Sy. E. B.* 270 a.—St. 2—3 ft. high. L. wavy at the edge, much and

uniformly dotted, declining. Fl. in very lax, much branched cymes. Pet. yellow, tinged externally on one longitudinal half with red. Anth. with a black spot. Styles divaricate.—Boggy places in Devon and Cornwall. P. VII. E.

5. *H. perforatum* (L.); st. erect 2-edged, l. oblong with pellucid dots, sep. erect lanceolate acute, pet. obliquely oblong, styles as long as the capsule.—*E. B.* 295. *R.* vi. 343.—St. 1—2 feet high. L. elliptic-oblong, they vary much in form, and in the number and size of the pellucid dots; chief veins pellucid, but not forming a network. Sep. usually denticulate near the tip.— $\beta$ . *angustifolium*; l. linear-oblong, sep. lanceolate acute finely denticulate.—Woods, hedgebanks, &c. P. VII. VIII.

E. S. I.

6. *H. dubium* (Leers); st. erect quadrangular, l. elliptic-ovate blunt with a few pellucid dots *netted with pellucid veins*, sep. reflexed with many black dots on the outside, pet. elliptic, styles half as long as the capsule.—*E. B.* 296. *H. quadrangulum* Fries.—St. 1—2 ft. high. Fl. in forked terminal cymes. Sep. ovate, blunt, nearly entire. Caps. longitudinally striate.— $\beta$ . *maculatum*; sep. oblong-lanceolate minutely denticulate, l. narrower.—Moist places by ditches, &c. P. VII. E. S. I.

7. *H. humifusum* (L.); st. prostrate somewhat 2-edged, l. oval-oblong blunt minutely pellucid-punctate the margins with black dots beneath, fl. subcymose, sep. unequal, 3 oblong blunt mucronate, 2 lanceolate, all entire or glandular-serrate and having a few black dots beneath, *stam.* 15—20, styles very short.—*E. B.* 1226. *R.* vi. 342.—St. slender, 3—6 in. long.—Gravelly and heathy places. P. VII. E. S. I.

8. *H. linariifolium* (Vahl); st. erect or ascending terete, l. linear blunt with revolute margins, fl. cymose, sep. rather unequal lanceolate-acute with glandular teeth and with many black dots beneath, *stam.* about 30, styles half as long as the capsule.—*E. B.* S. 2851.—St. 6—12 in. high. Fl. larger than in Sp. 7.—Channel Isles. Cape Cornwall. Banks of the Teign, Tamar and Tavy, Devon. P. VII. E.

9. *H. hirsutum* (L.); st. erect round hairy, l. ovate or oblong slightly stalked pellucid-punctate pubescent, sep. lanceolate acute fringed with shortly stalked glands, pet. linear oblong tipped with stalked glands, styles deciduous.—*E. B.* 1156. *R.* vi. 349.—St. about 2 ft. high, nearly simple. Fl. in axillary and terminal forked panicles.—Woods and thickets. P. VII. VIII. E. S. I.

10. *H. montanum* (L.); st. erect round glabrous, l. ovate-oblong sessile pellucid-punctate with glandular dots near the

margin, *sep. lanceolate acute fringed with shortly stalked glands*, pet. elliptic entire without dots or glands, styles half the length of the capsule.—*E. B.* 371. *R.* vi. 347.—Fragrant. St. 2 feet high, simple, smooth, slender. Fl. in terminal dense panicles.—Bushy limestone hills. P. VII. VIII. E. I.

11. *H. pul'chrum* (L.); *st.* erect round *glabrous*, *l. cordate amplexicaul pellucid-punctate glabrous*, *sep. broadly ovate blunt fringed with sessile glands*, pet. ovate-lanceolate fringed with glands.—*E. B.* 1227.—St. 12—18 in. high, nearly simple. Fl. in loose, axillary, opposite, and terminal cymes. Buds tipped with red. Anth. red.—Dry heaths, banks, woods. P. VI. VII. E. S. I.

\*\*\*\* *Styles 3; Stam. united throughout their lower half in 3 sets, a scale between each set; pet. equal-sided.* ELODES.

12. *H. elódes* (L.); *st.* ascending round *shaggy rooting below*, *l. roundish-ovate sessile pellucid-punctate shaggy*, *sep. ovate bluntish glabrous fringed with shortly stalked (reddish) glands*, pet. ovate entire, styles nearly as long as the capsules.—*E. B.* 109. *E. palustris* *R.* vi. 342.—St. prostrate below, then ascending and leafy, 6—8 in. long. Fl. in terminal and axillary few-flowered cymes.—Spongy bogs. P. VII. VIII. E. S. I.

### Order XVIII. ACERACEÆ.

Cal. 5-, rarely 5—9-parted, imbricate. Pet. the same number, inserted round an hypogynous disk. Stam. generally 8, inserted on the hypogynous disk. Ovary 2-lobed, 2-celled. Style 1. Stigmas 2. Fruit winged, separating into two indehiscent nuts (samaras) each with 1 cell and 1—2 seeds. Embryo curved, albumen 0.—Trees with opposite leaves. Stip. 0.

1. ACER. Fl. some imperfect. Calyx 5-parted. Pet. 5. Stam. usually 8, longer in the male flowers.

#### 1. A' CER *Linn.* Maple.

1. *A. campes'tre* (L.); *l.* 5-lobed, lobes entirely or slightly cut, *corymbs erect*, *sep. and pet. linear hairy*, wings of the fruit horizontally diverging, ovary downy, *stam. of the male flowers as long as the corolla.*—*E. B.* 304.—A small tree with corky fissured bark.—Woods and hedges. T. V. VI. *Maple.* E. S. I.

†2. *A. Pseudo-plat'anus* (L.); *l.* 4-lobed, unequally serrate, *racemes pendulous*, ovary downy with spreading wings, *stam. of the male flowers twice as long as the corolla.*—*E. B.* 303. *R.*

v. 164.—A large handsome tree.—In hedges and plantations even in the North of Scotland. T. V. VI. *Sycamore*. E. S.

### Order XIX. GERANIACEÆ.

Sep. 5, persistent, imbricate. Pet. 5, clawed, twisted in the bud. Stam. generally monadelphous, 2 or 3 times as many as the petals, some often abortive. Fruit of 5 carpels cohering round a long beaked axis, each terminated by an indurated style which finally twists up, separating from the axis and carrying with it the carpel. Seeds solitary, without albumen. Cotyledons convolute, plaited.—L. with stipules.

1. GERANIUM. Sep. 5. Pet. 5. Stam. 10, monadelphous, alternately larger and with glands at their base, all perfect. Carp. rounded at the top; the long ultimately recurved beak glabrous internally.
2. ERODIUM. Sep. 5. Pet. 5. Stam. monadelphous, 5 sterile, 5 fertile with glands at their base. Carp. with 2 lateral depressions at the top; the long ultimately spirally twisted beak bearded internally.

#### 1. GERA'NIUM Linn. Cranesbill.

\* *Root consisting of long fibres springing from a short thick rhizome, perennial.*

†1. *G. phæum* (L.); peduncles 2-flowered, pet. roundish wedgeshaped rather longer than the mucronate sepals, carp. hairy below *transversely wrinkled* above, *seeds punctate-striate*.—*E. B.* 322. *R. v.* 197.—St. erect, 2 feet high. L. 5-lobed; lobes acute, cut, serrate. Fl. purplish black, very rarely white.—In woods and thickets, rare. P. V. VI. E. S.

[*G. nodosum* (L.); peduncles 2-flowered. *pet. obovate long, sepals awned, carp. downy even*, l. 3—5-lobed, lobes ovate acuminate serrate.—*E. B.* 1091.—St. 18 in. high, slender, erect. Fl. pale purple.—In Cumberland, Herts, and Yorksh.—*G. striatum*, which resembles this but has a hairy stem, was found near Filby, Cumb.; Chepstow, and near Plymouth. Colonists.] E.

2. *G. sylvaticum* (L.); peduncles 2-flowered, pet. obovate slightly notched long, sepals awned, carp. hairy even, hairs spreading grandular, *seeds dotted*, l. palmate 7-lobed, lobes cut and serrate, st. erect glandular hairy above, *filaments* of *stam. subulate, fruitstalks erect*.—*E. B.* 121.—St. erect, 2—3 feet high. Fl. purplish blue, claws of the petals bearded, lower half of

filaments hairy. Sometimes the fl. are pale rose-coloured, pet. smaller and nearly entire, and st. more decidedly hairy.—Woods and thickets in the North, rare. P. VI. VII. E. S. I.

3. *G. pratense* (L.); ped. 2-flowered, pet. obovate entire or slightly notched long, sepals awned, carp. hairy even, hairs spreading glandular, seeds minutely netted, l. palmate 7-lobed, lobes cut and serrate, st. diffuse glandular hairy above, hairs deflexed, filaments of stam. filiform with a triangular-ovate base, fr.-st. deflexed.—E. B. 404.—St. 1—3 feet high.—Fl. large, purple, claw of pet. ciliate. Base of filaments slightly hairy.—Moist pastures. P. VI.—VIII. E. S. I.

4. *G. sanguineum* (L.); peduncles mostly single-flowered, pet. obcordate long, sepals awned, carp. smooth crowned with a few bristles, seeds minutely wrinkled and dotted, l. nearly round 7-lobed, lobes deeply 3-fid and cut, st. diffuse hairy, hairs spreading horizontally.—E. B. 272.—Fl. large, purple; filaments dilated at the base.—β. *G. prostratum* (Lindl.); st. dwarf tufted nearly simple decumbent, fl. flesh-coloured. *G. lancastrisense* With.—In dry places, rare. β. Sands in Walney Island, Lancashire. P. VII. E. S. I.

\*\* *Roots fusiform, rhizome wanting, perennial; ped. 2-flowered.*

†5. *G. pyrenæicum* (L.); fruitstalks deflexed, pet. obcordate twice as long as the mucronate sep., claws densely ciliate, carp. smooth with adpressed hairs, seeds smooth, l. reniform 7—9-lobed, lobes of lower l. oblong blunt trifid and toothed at the end, st. erect villose.—E. B. 405. R. v. 191.—Fl. light purple or nearly white. Claws of the pet. with a dense tuft of hairs on each side. Fertile anth. 10. Segments of the upper leaves more acute. St. spreading; 1—3 feet high, clothed with dense short down and long hairs intermixed.—Roadsides and pastures, rare. P. VI. VII. E. S. ? I.

6. *G. molle* (L.); pet. oblong deeply bifid  $\frac{1}{2}$  as long as or  $\frac{1}{2}$  longer than the mucronate sepals, claws ciliate, carp. transversely wrinkled glabrous, seeds smooth, l. roundish-reniform in 7—9 deep wedged shaped segments trifid at the end, st. diffuse pubescent.—E. B. 778. R. v. 191.—Fl. small, purple, or white with lilac claws. Softly pubescent, glandular above.—Dry places. A. IV.—VIII. E. S. I.

7. *G. rotundifolium* (L.); pet. spatulate entire blunt rather longer than the shortly awned sepals, claws glabrous, carp. not wrinkled with spreading hairs, seeds netted, l. reniform in 5—7 broadly wedged shaped incise-crenate segments, st. diffuse pubescent.—E. B. 157. R. v. 190. *G. viscidum* Ehrh.—Fl. small,



flesh-coloured. Peduncles shorter than the leaves.—Old walls and waste places, rare. A. VI. VII. E. I.

\*\*\* *Root fusiform, annual; ped. 2-flowered.*

8. *G. pusillum* (L.); pedicels deflexed after flowering, *pet. notched hardly exceeding the mucronate sepals, claws slightly ciliate, carp. not wrinkled with adpressed hairs, seeds smooth*, l. reniform palmate with 5—7 trifid lobes, *st. diffuse downy*.—*E. B.* 385. *R. v.* 190. *G. rotundifolium* Fries.—St. usually prostrate, clothed only with short down. Fl. small, bluish purple. Claws of the *pet.* only slightly ciliate. Fertile anth. 5. Styles pale flesh-coloured. Ped. shorter than the leaves.—Waste places. A. VI.—IX. E. S. I.

9. *G. dissectum* (L.); *pet. bifid about equalling the awned sepals, claws slightly ciliate, carp. not wrinkled with erect hairs, seeds netted*, l. in 5—7 deep lacinate segments with linear lobes, *st. diffuse hairy*.—*E. B.* 753. *R. v.* 189.—Fl. small, bluish purple. *L. divided almost to the base, longer than the peduncles*.—Waste places. A. VI.—VIII. *Dove's-foot*. E. S. I.

10. *G. columbinum* (L.); *pet. obovate emarginate with a short blunt tooth in the notch about equalling the awned sepals, claws ciliate, carp. not wrinkled with a few minute scattered hairs, seeds netted*, l. in 5—7 deep lacinate segments, *st. diffuse with adpressed hairs*.—*E. B.* 259. *R. v.* 198.—Fl. small rose-coloured. *L. divided almost to their base. Peduncles longer than the leaves; pedicels very long*.—On gravelly and limestone soils. A. VI. VII. E. S. I.

11. *G. lucidum* (L.); *pet. obovate entire, claws glabrous very long nearly equalling the transversely wrinkled pyramidal calyx, carp. netted triply keeled glandular-hairy at the summit, seeds smooth*, l. reniform in 5 blunt incise-crenate mucronate segments, *st. spreading ascending*.—*E. B.* 75. *R. v.* 187.—Fl. small, rose-coloured. St. and l. glabrous and shining, often strongly tinged with red.—Lindley considered his *G. Raii* most allied to this species, differing by its "shaggy calyx and simply keeled fruit."—South coast of England.—Walls and hedgebanks. A. V.—VIII. E. S. I.

12. *G. robertianum* (L.); *pet. obovate entire or slightly emarginate, claws glabrous very long nearly equalling the long-awned hairy and slightly glandular sepals, carp. transversely wrinkled downy, seeds smooth*, l. ternate or quinate, leaflets stalked trifid incise-pinnatifid, *st. spreading erect*.—*E. B.* 1486. *R. v.* 187.—Fl. purple, sometimes white. Cal. with a very few

glandular hairs, not transversely wrinkled.— $\beta$ . *G. purpureum* (Forst.); pet. narrower, sep. glandular-hairy, carp. glabrous and more wrinkled, l. in narrower segments. *E. B. S.* 2648.—Hedgebanks.  $\beta$ . Southern sea-coast. A. V.—IX. E. S. I.

## 2. *ERO'DIUM L'Hérit.* Storksbill.

1. *E. cicutarum* (Sm.); st. procumbent hairy, peduncles many-flowered, perfect *stam. dilated* not toothed below glabrous, beak hairy, a concentric furrow below the circular glandless depression on the carpel, l. pinnate, leaflets sessile pinnatifid cut, *stip. lanceolate*.—*E. B.* 1768.—St. diffuse, leafy, with scattered hairs. Fl. purplish or white. Leaflets very deeply divided, their segments lanceolate or linear, acute. In Jersey specimens the l. are ovate and short, and their segments short broad and bluntish.— $\alpha$ . *pimpinellifolium*; 2 pet. with a spot, lts. ovate incise-pinnatifid with bluntish lobes, furrow on the carp. conspicuous. *E. commixtum* Jord.— $\beta$ . *E. triviale* (Jord.); pet. not spotted, lts. incise-pinnatifid, carp.-furrow faint.— $\gamma$ . *E. pilosum* (Bor.); pet. not spotted, lts. almost pinnate, carp.-furrow obsolete.—Waste ground. A. VI.—IX. E. S. I.

2. *E. moschatum* (Sm.); st. procumbent hairy, peduncles many-flowered, perfect *stam.* toothed at the base glabrous, beak downy, a concentric furrow below the circular *glandular depression* on the carpel, l. pinnate, *leaflets* nearly sessile, *ovate unequally cut, stip. oval*.—*E. B.* 902.—Much larger than the preceding, and diffusing a strong musky scent when handled. Leaflets less deeply cut.—Waste places, rather rare. A. VI. VII. E. I.

3. *E. maritimum* (L'Hér.); st. prostrate slightly hairy, peduncles 1—2-flowered, pet. very minute, a transverse furrow below the semicircular depression on the carpel, l. simple ovate-cordate stalked lobed and crenate.—*E. B.* 646.—St. often very fleshy. Fl. very small. Pet. pale red, very minute, often wanting.—Sandy and gravelly places, particularly near the sea, rare. P. V.—IX. E. I.

## Order XX. BALSAMINACEÆ.

Sep. 5 (2 upper usually wanting), irregular, deciduous, lower spurred, imbricate in the bud. Pet. 5, irregular, the lateral united in pairs. Stam. 5. Anth. 2-celled, opening at the apex by a longitudinal fissure, more or less cohering. Ovary 5-celled. Fr. capsular with 5 elastic valves. Seeds solitary or many, pendulous; albumen 0.—L. with glands in place of stipules.

Plants succulent. Many raphides. Minute but fertile fl. which do not open are found.

1. IMPATIENS. Sep. 3, the lower one hoodlike with a spur. Pet. 3, upper one symmetrical, lateral unequally 2-lobed or each formed of 2 combined. Anth. cohering.

1. IMPA'TIENS Linn. Balsam.

1. *I. Noli-me-tan'gere* (L.); l. ovate coarsely serrate, peduncles many-flowered solitary, spur loosely recurved not emarginate.—*E. B.* 937. *R. v.* 198. b. *St.* 5. 15.—Fl. large, yellow spotted with orange. *St.* 1—2 feet high, tumid at the joinings.—Damp woody places in mountainous districts, rare.—*A.* VI.—IX. *Yellow Balsam.* E. S. I.

\*2. *I. ful'va* (Nutt.); l. ovate coarsely serrate, peduncles about 4-flowered solitary, spur closely reflexed emarginate.—*E. B. S.* 2794.—Fl. orange-yellow spotted with red. Each serrature of the l. with a reflexed glandular tooth. *St.* 2—3 feet high.—An American plant quite naturalized by the Wey and other rivers in Surrey. *A.* VIII. E.

\*3. *I. parvifl'ora* (DC.); l. elliptic serrate, ped. erect 3—12-flowered, spur short straight.—*Sy. E. B.* 315.—Fl. small, yellowish. *St.* 6—18 in. high. L. very acute at both ends.—Waste places. A Russian plant now quite naturalized. *A.* VII.—IX. E.

Order XXI. OXALIDACEÆ.

Sep. 5, equal, persistent, imbricate in the bud. Pet. 5, equal, often cohering at the base, twisted in the bud. Stam. 10, more or less monadelphous, those opposite to the pet. longer than the others. Anth. 2-celled. Ovary 5-celled. Styles 5. Caps. 5—10-valved. Seeds several. Embryo straight, in cartilaginous albumen.—With stipules.

1. OXALIS. Sep. 5, connected below. Pet. 5, often connected below. Stam. 10, monadelphous, 5 outer ones shorter. Styles 5. Seed with an elastic coat.

1. OX'ALIS Linn. Wood-Sorrel.

1. *O. Acetosel'la* (L.); stemless, rhizome creeping toothed, alternate, leaflets obcordate hairy, peduncles with 2 scaly bracts at about the middle 1-flowered, caps. ovoid, seed longitudinally ribbed.—*E. B.* 762. *R. v.* 199.—Fl. white with purple veins

or rarely purple or blue. Cor. about 4 times as long as the calyx.—Woods and shady places. P. V. *Wood-Sorrel*.

E. S. I.

†2. *O. corniculata* (L.); *st. diffuse with procumbent branches pubescent, l. ternate, leaflets obcordate, stipules oblong united to the base of the petioles, peduncles about 2-flowered shorter than the leaves, partial fruitstalks reflexed, caps. narrowly oblong, seeds transversely ribbed, root fibrous.*—*E. B.* 1726. *R.* v. 199.—Fl. yellow. L. mostly in pairs.—Waste ground in Devon and Cornwall. A. VI.—IX. E.

\*3. *O. stricta* (L.); *st. erect, stipules 0, ped. 2—8-fl. longer than the l., fl. cymose, fruitstalks erect, stoloniferous.*—*Sy. E. B.* 312. *R.* v. 199.—Cymes close, terminal; fl. yellow. L. in imperfect whorls.—Naturalized in Cornwall, Devon, Glamorgan, &c. B. VII. VIII. E.

## Order XXII. LINACEÆ.

Sep. 4—5 persistent, imbricate. Pet. 4—5, twisted in the bud, clawed, deciduous. Stam. as many as the pet., connected into an hypogynous ring with intermediate teeth (abortive stamens). Ovary with about as many cells and styles as the sepals, stigmas capitate. Caps. generally tipped with the hardened base of the styles, with 3—5 complete and 4—5 incomplete dissepiments, and no central axis. Seeds 1 in each spurious cell, pendulous, with albumen.—L. without stipules, alternate.

1. LINUM. Sep. 5, entire. Pet. 5. Stam. 5. Styles 5. Caps. with 10 cells and 10 valves.
2. RADIOLA. Sep. 4, connected below, deeply trifid. Pet. 4. Stam. 4. Styles 4. Caps. with 8 cells and 8 valves.

### 1. LINUM Linn. Flax.

\* *Leaves scattered. Margins of the sep. not glandular.*

1. *L. angustifolium* (Huds.); caps. downy within, *sep. elliptic pointed ciliate, l. linear-lanceolate, st. many.*—*E. B.* 381.—Fl. pale blue. St. 1—2 feet long, lax, diffuse, branching irregularly.—Sandy and chalky places in the South. P. VII. E. I.

[*L. usitatis-simum* (L.); caps. glabrous within, *sep. ovate pointed ciliate, l. lanceolate, st. solitary.*—*E. B.* 1357. *St.* 26. 12.—Fl. blue. St. 1—1½ foot high. Sep. 3-veined.—*β. crepitans* (Schub.); smaller and more branched, caps. opening with elasticity, seeds paler.—Escaped from cultivation. A. VII. *Common Flax.*]

2. *L. peren'ne* (L.); sep. obovate obscurely 5-veined glabrous, inner sep. very blunt, l. linear-lanceolate, st. many, fruitstalks erect.—*E. B.* 40.—Fl. blue. St. 1—2 ft. long, erect or decumbent.—Chalky places, rare. P. VI. VII. E.

\*\* *Leaves opposite.*

3. *L. cathar'ticum* (L.); sep. elliptic pointed, l. obovate, upper l. lanceolate.—*E. B.* 382.—Fl. white, small; sep. serrate; pet. acute. St. one or more, slender, 2—6 in. high. Panicle forked, spreading.—In dry pastures. A. VI.—VIII. E. S. I.

## 2. RADI'OLA *Gmel.* Flax-seed. All-seed.

1. *R. millegrána* (Sm.).—*E. B.* 893. *R. linoïdes* DC., Koch.—St. 1—2 in. high, repeatedly forked, with solitary minute white fl. in the forks as well as at the ends of the branches. Sep. deeply and acutely 3-cleft, connected below into a tube.—Damp sandy places. A. VII. VIII. E. S. I.

## Division II. CALYCIFLORÆ.

(Orders XXIII.—XXXIX.)

Petals distinct, and stamens perigynous or epigynous.

A. Petals and stamens perigynous (except in Orders XXVIII., XXIX., XXXI. and XXXV.).

### Order XXIII. CELASTRACEÆ.

Sep. 4—5, imbricate in the bud. Pet. 4—5, inserted in the margin of a fleshy disk surrounding the ovary. Stam. alternate with the petals, inserted in the disk. Ovary sunk in the disk, more or less connected with it, 3—5-celled; cells 1—2-seeded; ovules erect. Embryo straight.—Trees or shrubs. Stip. minute, deciduous.

1. EUONYMUS. Cal. flat, 4—5-lobed; disk peltate. Pet. 4—5, inserted in the margin of the disk. Stam. 4—5, inserted in the disk. Style 1. Caps. 3—5-celled, 3—5-angled, loculicidal. *Seeds solitary in each cell, with a fleshy aril, not truncate at the hile.*—Leaves simple.

1. EUON'YMUS *Linn.* Spindle-tree. Prickwood.

1. *E. europæ'us* (L.); pet. oblong, fl. mostly 4-cleft and 4-androus, branches tetragonous smooth and even, l. ovate-lanceolate minutely serrate, caps. obtusely angular not winged.—

*E. B.* 362. *R.* vi. 309. *St.* 27. 3.—Orange-coloured aril enclosing the seed. Bark green. L. glabrous. Fl. forming small umbels, greenish white. Fruit rose-coloured.—Hedges and woods. Sh. V. VI. E. S. I.

#### Order XXIV. RHAMNACEÆ.

Cal. 4—5-cleft, valvate in the bud. Pet. distinct, inserted in the throat of the calyx. Stam. opposite to the pet. and equalling them in number. Ovary wholly or in part superior, 2—3—4-celled, surrounded by a glandular disk. Seeds solitary, erect. Embryo straight. Fr. fleshy or dry.—Shrubs. Stip. minute.

1. RHAMNUS. Cal. pitcher-shaped, 4—5-cleft. Pet. 4 or 5, or sometimes 0, inserted with the stam. on the margin of the tube of the calyx. Fruit fleshy, with 2—4 cells and as many seeds.

#### 1. RHAM'NUS Linn. Buckthorn.

1. *R. cathar'ticus* (L.); thorns terminal, fl. 4-cleft dioecious, petioles much longer than the stipules, l. roundish-oval sharply toothed, fr. with 4 seeds.—*E. B.* 1629.—Branches opposite. Seratures of the l. incurved, glandular. Notch in the seeds closed. Styles 4, united halfway up.—Hedges and thickets. Sh. V.—VII. *Buckthorn.* E. S. I.

2. *R. Fran'gula* (L.); spineless, fl. 5-cleft perfect, l. elliptic-ovate acuminate narrowed below entire, fr. with 2 seeds, style simple.—*Sy. E. B.* 319.—Branches alternate. Fl. in small clusters greenish white, small.—Hedges and thickets. Sh. V. VI. *Black Alder.* E. S. ? I.

#### Order XXV. LEGUMINOSÆ.

Cal. inferior. Sep. 5, more or less combined, odd one inferior. Cor. papilionaceous (in our plants), inserted into the base of the calyx. Pet. 5, odd one superior and external. Stam. 10 (in our plants), monadelphous or diadelphous. Ovary free, 1-celled. Fruit a legume; placenta on the upper suture; style from the upper suture. Embryo bent over the edge of the cotyledons, or straight.—All our plants have papilionaceous flowers and 10 stamens in one bundle or in two bundles of 9 and 1. L. mostly stipulate; lts. often stipulate.

Tribe I. *LOTEÆ*. Pod continuous. Cotyledons rising above the ground and becoming green leaves. Leaves of 1 or 3 leaflets or pinnate with a terminal leaflet.

\* *Leaflet solitary. Stamens monadelphous.*

1. ULEX. *Cal. of 2 parts; the upper with 2, the lower with 3 minute teeth, a bract on each side at the base. Pod thick, few-seeded, scarcely longer than calyx which nearly equals the corolla.*
2. GENISTA. *Cal. 2-lipped; upper bifid, lower 3-toothed. Style subulate, ascending. Stigma terminal, oblique. Cor. much exceeding calyx.—Many foreign species have ternate leaves.*

\*\* *Leaf of 3 leaflets. Stamens monadelphous.*

3. SAROTHAMNUS. *Cal. 2-lipped, the upper with 2, the lower with 3 teeth. Style long, curved, thickened upwards, channelled within. Stigma terminal, capitate, small. Pod flat.*
4. ONONIS. *Cal. 5-cleft; segments narrow, the lower ones longer. Keel beaked. Style filiform, ascending. Stigma terminal, subcapitate. Pod thick.*

\*\*\* *Leaf of 3 leaflets. Stamens diadelphous; one free.*

5. MEDICAGO. *Cal. with 5 nearly equal teeth. Keel blunt. Filaments of the stamens filiform. Ovaries curved. Pod 1-celled, hooded or spirally twisted. Seeds 1 or many.*
6. MELILOTUS. *Cal. with 5 nearly equal teeth. Keel blunt. Filaments filiform, not adhering to the claws of the petals. Ovary straight. Pod subglobose or oblong, 1-celled, 1—4-seeded, longer than the calyx. Pet. distinct, deciduous. Fl. in long loose racemes.*
7. TRIFOLIUM. *Cal. with 5 unequal teeth. Keel blunt. Filaments slightly enlarged upwards and more or less adhering to the claws of the petals. Pod oval, 1—4-seeded, included in the calyx or slightly protruding. Pet. slightly combined, persistent.—Fl. in close racemes.*
8. FALCATULA. *Cal. of 5 nearly equal teeth. Keel blunt. Filam. filiform, not adhering to the claws of the petals. Pod compressed, truncate, straight, 6—8-seeded, protruding from the calyx. Pet. distinct, deciduous; wings and keel nearly equally long.—Fl. 1, 2 or 3 together. Common peduncle shorter than the petiole.*
9. LOTUS. *Cal. with 5 nearly equal teeth. Keel ascending, with a narrowed point (beak). Wings connivent at their*

upper margin. Longer filaments dilated upwards. Style kneed at the base, filiform-subulate. *Pod linear many-seeded*, 2-valved, imperfectly divided by transverse partitions.

\*\*\*\* *Leaf imparipinnate.* † Stamens monadelphous.

10. ANTHYLLIS. *Cal. tubular inflated*, 5-cleft, segments unequal. *Keel not beaked.* Style filiform. Stigma capitate.

†† Stamens diadelphous; one free.

11. OXYTROPIS. *Cal. with 5 teeth. Keel with a narrow straight point.* Pod imperfectly 2-celled, cells formed by the inflexed margin of the upper suture.

12. ASTRAGALUS. *Cal. with 5 teeth. Keel blunt.* Pods imperfectly 2-celled, cells formed by the inflexed margin of the lower suture.

Tr. II. VICIÆ. Pod continuous. *Stam. diadelphous*; one free. Cotyledons remaining under ground. *L. pinnate without the term. odd leaflet* or apparently simple.

\* *Tube of stam. very obliquely truncate.*

13. VICIA. *Cal. 5-fid or 5-toothed. Style filiform; its upper part hairy all over, or bearded on the underside* and at the same time hairy or glabrous. Pods 1-celled, 2-valved.

\*\* *Tube of stam. transversely truncate.*

14. LATHYRUS. *Cal. 5-fid or 5-toothed. Style flattened upwards, hairy beneath the stigma.* Pods 1-celled, 2-valved.

Tr. III. HEDYSARÆ. Pod dividing transversely into 1-seeded joints. *L. imparipinnate.*

15. ORNITHOPUS. *Cal. long, tubular, with 5 nearly equal teeth, 2 upper ones slightly combined and converging. Keel blunt. Pod long, compressed, of many 1-seeded indehiscent joints equally narrowed on both sides at the joinings.*—Apex of the common peduncle bearing a small pinnate leaf just below the flowers.

16. ARTHROLOBIUM. *Cal. long, tubular, with 5 nearly equal teeth, 2 upper ones combined up to their middle and straight. Keel blunt. Pod long, terete, of many 1-seeded indehiscent*



*joints scarcely narrowed at the joinings.—No leaf at the apex of the peduncles.*

17. HIPPOCREPIS. Cal. short, bellshaped, with 5 nearly equal teeth, 2 upper ones combined up to their middle. Keel narrowed into a beak. Pod long, compressed, of many 1-seeded crescent-shaped joints, so that each pod has many notches on one side.

18. ONOBRYCHIS. Cal. with 5 nearly equal subulate teeth. Keel obliquely truncate, longer than the wings. Pod 1-celled, compressed, indehiscent, 1-seeded, upper suture straight, lower curved toothed winged or crested.

### Tribe I. *Loteæ*.

1. U'LEX Linn. Furze. Whin. Gorse.

1. *U. europæus* (L.); young l. shaggy beneath furrowed, primary spines strong terete-polygonal furrowed rough, st. hairy, fl. lateral, bracts ovate lax, cal. shaggy, wings longer than keel.—*E. B.* 742.—St. shrubby, 4—6 feet high, much branched, spreading. Fl. bright yellow, from both the primary and secondary spines. Cal.-teeth converging. Spines branching at their base and up to about their middle, not exceeding the flowers.—*U. strictus* (Mack.); primary spines small slender tetragonal, plant 1—2 ft. high, with upright branches. Does not come true from seed. *E. B. S.* 2988.—Heaths. Sh. II.—VI. E. S. I.

2. *U. návus* (Forst.); young l. glabrous ciliate furrowed, primary spines slender terete striate smooth, st. hairy, fl. lateral and terminal, bracts very minute adpressed, cal. finely downy, wings shorter than keel.—*E. B.* 743.—St. shrubby procumbent. Primary spines short, spreading, branched at their base only. Fl. half the size of those of *U. europæus*, from the primary spines and exceeded by them, pale; pet. scarcely separated when full-blown. Cal.-teeth diverging. Pod persistent for nearly a year.—Heaths. Sh. VIII. IX. E.

3. *U. Gal'lii* (Planch.); young l. glabrous ciliate furrowed, primary spines strong deflexed subterete striate smooth, st. hairy, fl. lateral and terminal, bracts very minute adpressed, cal. finely downy, wings longer than keel.—*E. B. S.* 2987.—St. 2—5 ft. high. Fl. orange, pet. more or less divaricate; wings not straight and thus often seeming shorter than keel. Cal.-teeth diverging. Pods bursting in spring. A dwarf form is often

taken for *U. nanus*, a tall one for *U. europæus*.—Heaths. Sh. VIII.—XI. E. I.

## 2. GENIS'TA Linn.

1. *G. pilósa* (L.); st. procumbent without thorns, l. obovate-lanceolate blunt, *stipules ovate blunt*, branches peduncles calyx standard keel and underside of the l. silky, *peduncles lateral accompanied by a tuft of leaves, pods hairy*.—*E. B.* 208.—Fl. small, yellow, collected towards the end of the branches. St. much branched, furrowed, woody.—Dry sandy and gravelly heaths, rare. Sh. V. E.

2. *G. tinctoria* (L.); st. with erect branches without thorns, l. lanceolate or elliptic hairy at the edges, *stipules minute subulate, fl. racemose*, cor. and *Pods glabrous*.—*E. B.* 44.—Branches erect, 1—2 feet high, striate, glabrous, downy above. Fl. yellow. Keel as long as the standard.— $\beta$ . *G. humifusa* (Dicks.); st. and branches procumbent, l. ovate or oblong, pods hairy on the back of each valve. St. angular, 6—10 in. long.—In pastures and thickets.  $\beta$ . In the Lizard district, Cornwall. Sh. VII.—IX. *Dyer's-weed*. E. S. I.

3. *G. anglica* (L.); st. *spinous* leafless below, flowering branches glabrous without thorns, l. ovate-lanceolate, *stip. 0, fl. solitary* in the axils of the l., cor. and pods glabrous.—*E. B.* 132.—St. 1 foot high, round, leafless, with short leafy branches bearing the yellow flowers. Keel longer than the standard.—Moist peaty heaths. Sh. V. VI. *Needle Whin*. E. S.

## 3. SAROTHAM'NUS Wimm. Broom.

1. *S. scoparius* (Koch).—*Spartium* L., *E. B.* 1339.—St. 2—3 feet high, angular, glabrous; or prostrate and spreading (by Kynance Cove, Cornwall). L. ternate or simple, obovate. Fl. axillary, solitary or in pairs, shortly stalked, large, bright yellow. Pods dark brown, hairy at the edges; seeds many.—Dry hills and heaths. Sh. V. VI. E. S. I.

## 4. ONO'NIS Linn. Rest-harrow.

1. *O. arven'sis* (L.); stoloniferous, st. procumbent uniformly hairy, fl. axillary solitary stalked, leaflets broadly oblong, *Pods ovate erect falling short of the calyx*.—*E. B. S.* 2659. *O. repens Koch*.—Shrubby. Usually without spines. St. rooting at their base. Wings equalling the keel; standard a little longer. Seeds tubercular.— $\alpha$ . glandular, fl.-l. equalling or surpassing

cal., pod shorter than calyx.—*β. maritima*; glandular-villose, fl.-l. falling short of cal., pod as long or longer than calyx.—Barren sandy places. P. VI.—IX. E. S. I.

2. *O. campes'tris* (Koch); not stoloniferous, st. erect or ascending bifariously hairy, fl. axillary solitary stalked, leaflets oblong, pods ovate erect *exceeding the calyx*.—*E. B.* 682.—Shrubby. Usually spinous. St. mostly erect. Wings falling short of the keel, which falls short of standard. Seeds tubercular.—Barren places. P. VI.—IX. E. S.

3. *O. reclináta* (L.); viscid, pubescent, st. ascending, fl. axillary, pedicels 1-flowered shorter than the l. fl. or pod without bracts, cor. about equal to the calyx, leaflets obovate-cuneate serrate at the tip, stipules ovate, pods *cylindrical reflexed*, seeds 14—18 tubercular.—*E. B. S.* 2838.—St. 5—6 in. high, much branched.—Sandy places. With *Bupleurum aristatum* at Berry Head, Devon. *Mr. E. M. Holmes!* (*J. of B.* vi. 58). Channel Isles. \*Galloway. A. VII. E.

### 5. MEDICA'GO Linn.

\* *Pods without spines.*

(Sp. 1, 2, and 3 are said to be sometimes monadelphous.)

\*1. *M. satíva* (L. ; st. erect, racemes many-flowered, pods compressed *loosely spiral with 2 or 3 turns downy with adpressed hairs*, pedicels shorter than the calyx or bract, leaflets obovate-oblong dentate above emarginate mucronate.—*E. B.* 1749.—St. angular when young. Pods twisted into a loose open spiral. Fl. large, yellow or violet.—Hedgebanks and borders of fields, scarcely naturalized. P. VI. VII. *Lucerne.* E. S.

2. *M. sylves'tris* (Fries); st. rather quadrangular pithy ascending, racemes many-flowered, pods *forming one complete flat ring*, pedicels shorter than the cal. longer than the bract, lts. obovate-oblong dentate above emarginate mucronate.—*E. B. S.* 2980.—Fl. large, yellow or blackish green with darker streaks.—Sandy and gravelly places in Norfolk and Suffolk. P. VI. VII. E.

3. *M. falcáta* (L.); st. usually terete nearly solid prostrate, racemes many-flowered, pods *straightly sickle-shaped twisted not forming a ring*, pedicels shorter than the cal. longer than the bract, lts. obovate-oblong dentate emarginate mucronate.—*E. B.* 1016.—Fl. large, yellow.—Sandy and gravelly places in Norfolk and Suffolk. P. VI. VII. *Yellow Medick.* E.

4. *M. lupulina* (L.); spikes many-flowered dense oval, pods compressed kidney-shaped with a spiral point with longitudinal branched prominent veins, stip. obliquely ovate slightly toothed, leaflets roundish-ovate denticulate emarginate mucronate.—*E. B.* 971.—St. procumbent or ascending, spreading widely. Pods 1-seeded, glabrous or slightly hairy. Fl. small, yellow.—Waste ground. A. or B. V.—VIII. *Black Medick.* E. S. I.

\*\* *Pods edged with spines.*

5. *M. maculata* (Sibth.); peduncles 1—4-flowered, pods compactly spiral compressed of 2—6 turns veined with 4 ridges and a central furrow on the edge, spines in 2 rows divergent subulate curved, leaflets triangular-obcordate, stip. toothed.—*E. B.* 1616.—Lts. with a purple spot in the centre. Edge of pods broad; spines springing from the margin and the ridge next to it on each side, compressed and furrowed on both sides, variable in length.—On a gravelly soil. A. V.—VIII. E. S. I.

6. *M. minima* (Lam.); peduncles 1—6-flowered, pods compactly spiral of 4 turns smooth with a thin edge, spines in 2 rows divergent subulate hooked, leaflets obovate, stip. nearly entire.—*E. B. S.* 2635.—Edge of the pods with 3 ridges, the central one so prominent as to be easily taken for the true margin, no central furrow but the central ridge common to the 2 rows of spines. Sides of the pods smooth. Spines varying considerably in length and the whole plant in hairiness.—In sandy fields in the South-east, rare. A. V. E.

7. *M. denticulata* (Willd.); peduncles 1—5-flowered, pods rather loosely spiral consisting of 2 or 3 turns deeply netted with a thin edge, spines in 2 rows divergent subulate hooked, leaflets obcordate, stip. laciniate.—*E. B. S.* 2634.—Edge of the pods as in *M. minima*. Spines about equalling the diameter of the pod. Glabrous.— $\beta$ . *M. apiculata* (Willd.); spines very short without hooks, often scarcely longer than their own breadth so as to appear little more than tubercles, ped. 3—10-flowered.—On sandy ground near the sea, rare. A. V.—VIII. E. I.

6. MELLILOTUS Linn. Melilot.

1. *M. officinalis* (Willd.); wings keel and standard equal, pods ovoid acute compressed transversely wrinkled hairy, leaflets serrate truncate narrowly ovate, stip. setaceous entire.—*E. B.* 1340.—*M. macrorrhiza* Pers., Koch.—St. erect, 2—3 feet high. Fl. in lateral racemes, yellow.—Waste places. B. ? VI.—VIII. Common Melilot. E. S. I.

2. *M. arven'sis* (Willd.); wings and standard equal longer than the keel, pods ovoid blunt mucronate rounded and slightly keeled on the back transversely plicate glabrous, lts. obcordate or oblong serrate uppermost lanceolate, stip. awlshaped entire.—*E. B. S.* 2960. *M. officinalis* Koch.—St. erect. Fl. yellowish, in long racemes. Pods brown.—Waste places. Cambridge; Thetford, &c. B. VI.—VIII. E. I.

†3. *M. al'ba* (Lam.); wings and keel equal but shorter than the standard, pods ovate blunt mucronate netted glabrous, leaflets obovate the upper ones oblong serrate blunt, stip. awlshaped entire.—*M. leucantha* Koch, *E. B. S.* 2689.—St. erect. Fl. white.—Sandy and gravelly places near the sea, rare. B. VII. VIII. E. S. I.

\*4. *M. parvifl'ora* (Desf.); wings and keel equal but shorter than the standard, pods globular-ovoid blunt mucronate netted glabrous, lts. obovate serrate at the end, stip. awlshaped entire.—*Sy. E. B.* 344.—Slender. Smaller in all its parts than the other species, 6—15 in. high. Fl. very small, pale yellow; cal.-teeth triangular. Pods olive-green.—Waste places. A. VII. VIII. E.

## 7. TRIFO'LIUM Linn. Clover.

\* *Fl. sessile; cal. with an elevated thickened often hairy line or ring of hairs in its throat, not inflated.*

1. *T. pratense* (L.); heads ovate dense sessile, cal. 10-veined hairy not half so long as the corolla, teeth setaceous ciliate, free part of stip. blunt ovate abruptly bristle-pointed, leaflets oval emarginate upper ones entire apiculate.—*E. B.* 1770. St. 15. 11.—Lts. of cultivated plant usually all quite entire. Stip. adpressed to petiole; veins much branched and anastomosing. Upper part of cal. usually hairy; teeth 5, 4 nearly equal in length to the tube, the lower one twice as long. Heads of fl. sometimes slightly stalked. Fl. purplish, sometimes white. St. erect.—*β. parviflorum*; head stalked, calyx-teeth as long as or longer than the corolla.—Mountainous pastures, fields. *β* in dry places. P. V.—IX. Purple Clover. E. S. I.

2. *T. médium* (Huds.); heads subglobose lax stalked, calyx 10-veined glabrous not half as long as the corolla, teeth setaceous hairy, free part of stip. lanceolate acuminate, leaflets elliptic or lanceolate apiculate.—*E. B.* 190. St. 15. 13.—Stip. spreading; veins branching, parallel, scarcely at all joining. Cal. glabrous; teeth ciliate, 4 of them as long or rather longer than the tube, the fifth  $\frac{1}{3}$  longer. Heads of fl. large. Fl. purplish. St. ascending, zigzag.—Dry elevated pastures. P. VI.—IX. E. S. I.

3. *T. ochroleucum* (L.); heads subglobose dense stalked solitary terminal, cal. 10-veined pubescent about half as long as the corolla, teeth erect in fruit subulate lower one rather longer than the tube the others two-thirds shorter, stip. lanceolate-subulate, leaflets elliptic-oblong the lower one emarginate or cordate.—*E. B.* 1224. *St.* 15. 15.—Cal. with acute teeth having 1 strong vein. *St.* 1½ foot high, erect. Lower l. on very long stalks. Fl. cream-coloured, at length turning brown.—Dry gravelly soils and clays of the East of England. P. VI. VII. E.

4. *T. incarnatum* (L.); heads ovate at length cylindrical stalked solitary terminal, cal. 10-veined hairy, teeth patent in fruit lanceolate-subulate nearly equal rather longer than their tube and falling short of the cor., stip. ovate blunt, leaflets obovate retuse or obcordate, st. erect and together with the l. and stip. villose.—Mouth of fr.-cal; hairy. Stip. sometimes rather acute especially the lower ones.—*a*; hairs of stem patent, heads oblong, tips of cal.-teeth slightly hairy, fl. reddish purple.—*β. T. Molinieri* (Balb.); st. with adpressed hairs, heads conical, cal.-teeth glabrous, fl. nearly white. *E. B. S.* 2950.—Var. *a* is cultivated. *β* native near Lizard Point, Cornwall. A. VI. VII. E.

\*5. *T. stellatum* (L.); heads globose stalked terminal, calyx 10-veined hairy, teeth subulate from a broad base equal exceeding the corolla patent in fruit 3-veined and netted, throat closed with hairs, stip. ovate rather acute denticulate, leaflets obcordate, st. spreading and together with the l. and stip. villose.—*E. B.* 1545. *St.* 16. 5.—Fl. cream-coloured, small. Calyx of the fruit remarkably large, its teeth spreading like a star. *St.* short.—Shingly beach near Shoreham. A. VI. VII. E.

6. *T. arven'se* (L.); heads nearly cylindrical stalked very hairy, cal. 10-veined, teeth subulate-setaceous hairy nearly equal exceeding the cor. at length slightly spreading, stip. ovate or lanceolate acuminate, leaflets linear-oblong.—*E. B.* 944. *St.* 16. 3.—*St.* erect or, in a maritime form (*Ray Syn.* t. 14. f. 2), procumbent with globose heads; st. and l. finely hairy. Points of the lower stip. sometimes very slender. Fl. small, almost concealed by the very hairy calyx. Seeds oval, greenish yellow; radicle not prominent.—Sandy fields. A. VII.—IX. *Hare's-foot Trefoil.* E. S. I.

7. *T. striatum* (L.); heads ovate or oblong sessile terminal and lateral solitary or the terminal in pairs, calyx 10-veined hairy, teeth subulate unequal straight mucronate about equalling the cor., tube ventricose in fruit, stip. ovate cuspidate, leaflets obcordate or obovate, veins equal and straight at the margins.

—*E. B.* 1843. *St.* 16. 6, 7.—*St.* procumbent, 4—10 in. long and as well as the l. silky. *Stip.* with reddish veins. *Fl.* small. *Seeds* oval, brownish yellow; radicle not prominent.—*β. erectum* (Leight.): *st.* erect, heads long subconical lateral shortly stalked, cor. longer than calyx.—Dry and sandy places. A. VI. VII. E. S. I.

8. *T. scabrum* (L.); heads ovate sessile terminal and lateral solitary, cal. 10-veined hairy, *teeth lanceolate* mucronate about equalling the cor. with 1 strong prominent vein at length patent, tube cylindrical in fruit, *stip.* ovate cuspidate, leaflets obovate, veins thickened and curved at the margins.—*E. B.* 903.—*St.* procumbent, spreading. *Fl.* small. Cal. of the fruit very rigid. *Seeds* oblong, reddish yellow; radicle not prominent.—Dry sandy places. A. V.—VII. E. S. I.

9. *T. Bocconi* (Savi); heads oblong-ovate sessile terminal usually 2 together, cal. 10-veined hairy, teeth lanceolate-subulate mucronate about as long as the cor. with 1 strong prominent vein erect, tube cylindrical in fruit, *stip. oblong with a long subulate point*, leaflets oblong obovate or roundish obovate in the lower leaves, veins equal and straight at the margins, seeds with the radicle slightly prominent.—*E. B. S.* 2868.—*St.* 2—6 in. high, erect. *Fl.* small, pale yellow. *Seeds* oval, brownish yellow. Heads dense, somewhat conical.—In dry places, near the Lizard Point, Cornwall. A. VII. E.

10. *T. maritimum* (Huds.); heads ovate-globose stalked terminal, cal. strongly veined, *teeth ciliate at first subulate erect* falling short of the cor. afterwards broad leaflike acute spreading 1-veined the lower one longest and 3-veined, tube hairy above obconical in fruit, *stip. broadly subulate very long*, leaflets oblong-obovate.—*E. B.* 220.—*St.* spreading, usually procumbent. *Fl.* pale red, small.—Muddy salt marshes. A. VI. VII. E. I.

\*\* *Fl. sessile; throat of the calyx naked within; heads few-flowered, at length producing thick stellate fibres (abortive calyces) from their centre which ultimately fold over the fruit.*

11. *T. subterraneum* (L.); heads 2—5-flowered erect deflexed in fruit, calyx glabrous, teeth filiform hairy nearly equal shorter than the corolla, tube inflated in fruit and at length split longitudinally, *abortive calyces many slender with 5 points*, *stip.* ovate pointed, leaflets obcordate.—*E. B.* 1048.—*St.* prostrate and as well as the l. hairy. *Fl.* white, considerably longer than the calyx. The abortive calyces are remarkably characteristic of this species.—Dry gravelly places. A. V. VI. E. I.

\*\*\* *Fl. sessile or stalked; throat of the calyx naked within, not inflated; pods 2—4-seeded.*

12. *T. glomeratum* (L.); heads globose sessile terminal and axillary, calyx sessile 10-veined, teeth ovate very acute reviny nearly equal reflexed, stip. ovate taper-pointed, leaflets obcordate the upper ones obovate, seeds 2.—*E. B.* 1063.—St. procumbent. Fl. rose-coloured; standard persistent, striate. Seeds transversely ovate-reniform, radicle prominent.—Gravelly places East and South of England, rare. Wicklow, I. A. VI. E. I.

13. *T. strictum* (L.); heads globose axillary, calyx sessile 13-veined, teeth subulate unequal somewhat spreading, stip. broad pointed serrate, leaflets obovate the upper ones oblong, seeds 2.—*E. B. S.* 2949.—St. diffuse, short. Ped. short. Pods slightly projecting; seeds ovate, radicle slightly prominent.—Jersey. Near the Lizard Point, Cornwall. A. VI.—VIII. E.

14. *T. suffocatum* (L.); heads roundish sessile axillary; cal. sessile, teeth lanceolate acute recurved longer than the corolla, stip. ovate pointed, leaflets obcordate, seeds 2.—*E. B.* 1049.—St. short, usually buried in the sand. Fl. small, erect. Cal. scarcely striate. Seeds roundish, radicle prominent.—Sandy sea-shores, rare. A. VI. E.

15. *T. répens* (L.); heads roundish, ped. axillary longer than the leaves, fl. stalked at length deflexed, calyx glabrous half as long as the corolla, teeth lanceolate unequal erect, stip. ovate abruptly cuspidate, leaflets obovate or obcordate, seeds 4, stems creeping.—*E. B.* 1769.—Fl. white, sometimes pink, rarely lilac-purple (*T. Townsendii*); standard striate. L. often with a dark spot at their base. Pod covered by the faded corolla.—Meadows and pastures. P.V.—IX. Dutch or White Clover. E. S. I. \*

\*16. *T. hybridum* (L.); heads globular depressed, ped. axillary longer than the leaves, fl. stalked at length deflexed, cal. glabrous half as long as the (pinkish) corolla, teeth nearly equal subulate erect, stip. ovate-lanceolate attenuate, leaflets obovate-lanceolate, seeds 2—4, st. erect not rooting.—*Sy. E. B.* 361.—Fl. white or pale pink, turning brown. St. 1—2 ft. high, branched, wavy.—Introduced. P. VII.—IX. Alsike Clover. E. S.

\*\*\*\* *Fl. sessile; throat of the calyx naked within, inflated after flowering and arched above.*

17. *T. fragiferum* (L.); heads globose, peduncles axillary exceeding the leaves, involucre multifid equalling the calyx, calyx of the fruit membranous netted downy, stip. ovate with a long



slender *point*, leaflets obovate emarginate minutely serrate, seeds 2.—*E. B.* 1050. *St.* 16. 8.—St. creeping. Fl. purplish red. Heads large, remarkable when in fruit for their curious calyces enclosing the pods.—Damp pastures. P. VII. VIII. E. S. I.

[*T. resupinatum* (L.); heads hemispherical at length globose. peduncles axillary short, *bracts minute*, calyx of the fruit membranous netted woolly, *stip. subulate-lanceolate from an ovate base*, leaflets obovate minutely serrate, seeds 2.—*Sy. F. B.* 364. *St.* 16. 9.—St. prostrate or ascending. Fl. small, reversed in position. Pod included.—Introduced. A. VII.] E.

\*\*\*\*\* *Fl. stalked, yellow; throat of the calyx naked within, not inflated; cor. persistent; standard covering the pod.*

18. *T. procumbens* (L.); heads oval dense with about 40 fl., ped. axillary equalling or exceeding the leaves, fl. at length reflexed, *standard dilated and deflexed in front (not folded)* striate much exceeding the pod, style much shorter than the pod, *stip. ovate acute entire*, lts. obovate emarginate, central petiole longest, seeds oval, radicle scarcely prominent.—*E. B.* 945. *St.* 15. 15.—Primary stem erect, branches procumbent or ascending. Fl. turning tawny. Radicle causing a slight irregularity in the otherwise regularly oval seeds.—Dry pastures. A. VI.—VIII. E. S. I.

19. *T. minus* (Sm.); heads close about 12 fl., ped. axillary straight, peduncles very short, fl. at length reflexed, *standard folded furrowed truncate* covering the pod, style much shorter than the pod, *stip. ovate falling short of petioles*, lts. obcordate intermediate one stalked.—*E. B.* 1256. *T. dubium* Sibth. *T. filiforme* Koch.—St. wiry, ascending or prostrate. Fl. turning dark brown.—Dry places. A. VI.—VIII. E. S. I.

20. *T. filiforme* (L.); fl. few (2—7) in lax racemes, ped. axillary flexuose, pedicels as long as the calyx-tube, fl. at length reflexed, *standard folded not furrowed deeply notched* scarcely covering the pod.—*E. B.* 1257. *T. micranthum* Koch.—St. prostrate, very slender. Standard much narrower in proportion than in *T. minus*.—Dry places. A. VI. VII. E. S. I.

## 8. FALCAT'ULA Brot.

1. *F. ornithopodioides* (Bab.); clusters stalked axillary of 1—3 stalked fl., calyx glabrous, teeth slender acute nearly equal erect, *stip. ovate with long taper points*, leaflets obcordate, seeds 6—8.—*E. B.* 1047. *Trigonella* DC.—St. prostrate. Fl. small, pet. all distinct. Pod compressed, blunt, transversely furrowed, slightly hairy, curved, longer than the calyx, opening with 2 valves.—Dry gravelly places, rare. A. VI. VII. E. I.

9. *Lo'tus* Linn. Bird's-foot Trefoil.

1. *L. corniculátus* (L.); claw of the standard obovate transversely vaulted, *calyx-teeth adpressed in the bud subulate from a triangular base, points of the 2 upper teeth converging, heads 5—10-flowered.*—*E. B.* 2090.—Glabrous or slightly hairy. St. ascending. Angle between the 2 upper calyx-teeth rounded.—*a*; st. short, lts. obovate thin, stip. rather narrowly ovate.—*β. villosus* (Ser.); upper part of st. l. and cal. hairy with long spreading hairs.—*γ. crassifolius* (Pers.); pilose, st. cæspitose, leaflets obovate fleshy, stip. ovate.—*δ. L. tenuis* (W. and K.); glabrous or slightly hairy, st. filiform long procumbent or ascending, leaflets linear or linear-obovate, stip.  $\frac{1}{2}$ -ovate. *E. B. S.* 2615. Perhaps a distinct species.—Pastures, dry banks, &c. P. VI. —VIII. E. S. I.

2. *L. májor* (Scop.); claw of the standard linear, *calyx-teeth spreading like a star in the bud subulate from a triangular base, two upper teeth diverging, heads 8—12-flowered*, leaflets obovate, stip. roundish-ovate.—*E. B.* 2091. *L. uliginosus* Schkuhr.—Hairy. St. usually erect, 1—3 feet high. Angle between the 2 upper calyx-teeth acute. Sometimes glabrous, the margins and veins of the l. stip. bracts and sep. ciliate, st. erect or procumbent.—In damp places. P. VII. VIII. E. S. I.

3. *L. angustis'simus* (L.); claw of the standard linear, calyx-teeth straight in the bud subulate, *pod linear straight 6 times as long as the calyx, heads about 2-flowered.*—*E. B.* 925. *L. angustissimus* (L.!), *L. diffusus* (Ser.).—Standard broader than long, not exceeding the wings, fading to a greenish colour. Peduncle of the fl. as long as the l., of the fr. twice as long; leaflets and stip. ovate-lanceolate acute; st. procumbent.—*β. Seringianus* (Bab.); peduncle of the fl. and fr. as long as the leaves, leaflets obovate-oblong, stip. ovate acute, st. ascending. *L. angustissimus* (Ser.).—South of England near the sea. A. VII. VIII. E.

4. *L. his'pidus* (Desf.); claw of the standard subulate, calyx-teeth straight in the bud subulate, *pod rugose terete twice as long as the calyx, heads few (3—4)-flowered*, leaflets obovate-lanceolate, stip. half cordate, st. procumbent.—*E. B. S.* 2823.—Standard longer than broad, exceeding the wings, not turning green.—Near the sea in Devon, Dorset, and Cornwall. A. VII. VIII. E.

10. *Anthyl'lis* Linn. Lady's Finger.

1. *A. vulnerária* (L.); herbaceous, l. pinnate, leaflets unequal,

heads of fl. in pairs, calyx of 5 ovate pointed teeth.—*E. B.* 104. *St.* 49. 4, 5.—Pod semiorbicular, long-stalked, upper suture arched outwards, 1-seeded. *St.* 6—12 in. high, silky. Root-l. simple, oval. Fl. yellow, in terminal pairs of crowded many-flowered heads.— $\beta$ . *A. Dilleni* (Schulte); plant smaller, fl. red. *Dill. Eth.* 320.—Dry pastures. P. VI.—VIII. E. S. I.

### 11. OXYTROPIS *Cand.*

1. *O. Hal'leri* (Bunge); stemless, leaflets ovate acute in about 12 pairs, *peduncles* exceeding the leaves *erect silky*, bracts equaling the calyx, *Pods erect* ovate-oblong inflated *silky* imperfectly 2-celled.—*Astragalus uralensis* Sm. *E. B.* 466.—Rootstock woody, branched. Pods abrupt with a very oblique acute point. *Fl. bluish purple*. More silky and hairy than Sp. 2.—“Pl. uralensis ab europæa differt.” *Koch.*—Dry hilly pastures in Scotland. P. VII. S.

2. *O. campes'tris* (DC.); stemless, leaflets lanceolate in about 12 pairs, *peduncles* rather exceeding the l. *ascending hairy*, bracts equaling the calyx, *Pods erect* ovate inflated *hairy* imperfectly 2-celled.—*E. B.* 2522. *St.* 19. 12.—Rootstock woody, prostrate. Pods narrowed upwards with a slightly oblique point. *Fl. yellowish* tinged with purple.—Clova Mountains. P. VII. S.

### 12. ASTRAG'ALUS *Linn.* Milk-vetch.

1. *A. hypoglot'tis* (L.); st. prostrate, *stip. united*, leaflets blunt in 8—10 pairs, racemes ovate, *peduncles exceeding the leaves*, *Pods* ovate hairy *erect* stalked in the calyx.—*E. B.* 274.—*Stip.* opposite to the leaves. *St.* a few inches long, slender. Leaflets small. Fl. in rather large heads, ascending, purple. Ovary twice as long as its stalk.—Chalky and gravelly places, rare. P. VI. VII. E. S. I.

2. *A. alpinus* (L.); st. prostrate, *stip. ovate free*, leaflets elliptic blunt in 10—12 pairs, racemes short close, *peduncles* equaling the leaves, *Pods* oblong hairy narrowed at both ends *pendulous* stalked in the calyx.—*E. B. S.* 2717. *St.* 19. 13.—*Stip.* sometimes slightly connected at the base. *St.* a few inches long, slender. Fl. few, drooping, white tipped with purple.—Lofty mountains. Glen Dole, Clova. Little Craigindal, Braemar. P. VII. S.

3. *A. glycyphyl'los* (L.); st. prostrate, *stip. ovate-lanceolate* free, leaflets ovate in 5—6 pairs, racemes ovate, *peduncles* falling much short of the leaves, *Pods linear incurved erect glabrous*.—*E. B.* 203.—*St.* 2—3 feet long, scarcely branched, nearly

glabrous. Fl. in short dense racemes, dull yellow. Pods an inch long.—Thickets on a chalky or gravelly soil, rare. P. VI. *Wild Liquorice.* E. S.

## Tribe II. *Vicieæ*.

### 13. VI'CIA, Linn. Vetch.

\* *Upper part of the style equally hairy all over.*

† Peduncles long, few-flowered; cal. not gibbous at the base on the upperside.—Annuals. ERVUM. *Tares.*

1. *V. hirsúta* (Koch); *ped.* 1—6-*fl.* about equalling the leaves, leaflets in 6—8 pairs linear-oblong truncate mucronate, calyx-teeth equal as long as their tube the 2 upper ones converging, *Pods sessile oblong 2-seeded hairy, hile long linear.*—*Ervum* Sm. *E. B.* 970. *Cracca minor* Godr.—Stip. 2-lobed, outer lobe trifid with setaceous segments, inner lanceolate. Fl. small, pale blue, standard entire. Calyx-teeth subulate. Pod obliquely truncate; its upper suture nearly straight and prominent at the end. Seeds globose, compressed, red with darker spots, smooth. Ped. rarely 1—2-flowered and pods glabrous.—Corn-fields and hedges. A. VI.—VIII. *Hairy Tare.* E. S. I.

2. *V. tetrasper'ma* (Moench); *ped.* 1—2-*fl.* about equalling the leaves, leaflets linear-oblong blunt mucronate in 4—6 pairs, calyx-teeth unequal shorter than the tube the 2 upper ones shortest “diverging,” *Pods shortly stalked linear-oblong about 4-seeded glabrous, hile oblong.*—*Ervum* Sm. *E. B.* 1223. *St.* 32. 14.—Stipules half-arrowshaped. Fl. small, pale blue; standard with blue streaks, emarginate. Calyx-teeth long-triangular. Pod rounded; its upper suture decurved at the end. Seeds 3—5, globose, dull brown, slightly rough.—Fields and hedges. A. VI.—VIII. *Smooth Tare.* E. S. I.

3. *V. grávilis* (Lois.); *ped.* 1—4-*fl.* becoming twice as long as the leaves, leaflets linear acute in 3—4 pairs, calyx-teeth unequal shorter than their tube, the 2 upper ones shortest, *Pods linear 5—8-seeded glabrous, hile roundish-oval.*—*E. B. S.* 2904.—Stip. half-arrowshaped. *Fl. twice as large as those of V. tetrasperma,* pale blue; standard emarginate. Calyx-teeth long-triangular, “two upper slightly converging.” Upper suture of pod slightly decurved at the end. Seeds globose, variegated with dark brown and yellow, smooth, *half as large and hile half as long as those of Sp. 2.*—Fields and hedges in the South. A. VI.—VIII. E.

†† Peduncles long, many-flowered; calyx gibbous at the base on the upperside.—Perennial.

4. *V. sylvatica* (L.); ped. exceeding the leaves, leaflets elliptic blunt mucronate in about 8 pairs, *stip.* lunate *deeply toothed at the base, teeth setaceous, calyx-teeth* shorter than their tube, *subulate*.—*E. B.* 79. *St.* 31. 3.—*St.* many feet long, climbing by branched tendrils. Fl. many, cream-coloured and streaked with blue or purple. Hile extending about half round the seed.—Woods and thickets.—P. VII. VIII. *Wood Vetch.* E. S. I.

5. *V. Or'obus* (DC.); leaflets ovate-oblong or ovate-lanceolate mucronate in many (7—10) pairs, *stip.* half-arrowshaped *slightly toothed at the base, calyx-teeth* longer than their tube, 2 *upper ones triangular*, the others triangular-subulate.—*Orobus sylvaticus* L. *E. B.* 518.—*St.* ascending, 1—1½ foot long. Tendrils reduced to a short slender point. Fl. many, cream-coloured streaked with purple. Pods linear-oblong. Hile extending about ½ round the seed.—Northern rocky woods. P. V. VI. *Bitter Vetch.* E. S. I.

\*\* *Upper part of the style hairy all over, the hairs rather longer below the stigma but scarcely bearded; calyx gibbous at the base.*

6. *V. Crac'ca* (L.); ped. long many-flowered, lts. lanceolate mucronate silky in about 10 pairs, *stip.* half-arrowshaped *entire*, calyx-teeth shorter than their tube, upper pair minute, others subulate, standard sinuate at about the middle of each side, its limb and claw equally long, pods linear-oblong smooth.—*E. B.* 1168. *St.* 31. 6.—*St.* 3—4 feet long. Fl. blue varied with purple. Seeds subglobose, black. Hile linear, extending ½ round the seed.—Hedges. P. VI.—VIII. E. S. I.

\*\*\* *Style bearded below the stigma, in other respects glabrous or uniformly hairy all over its upper part; calyx gibbous at the base on the upperside.*

7. *V. bithyn'ica* (L.); ped. falling short of the leaves 1—2-flowered, lts. of upper l. in 2 pairs *elliptic-lanceolate mucronate or linear-lanceolate acute*, *stip.* half-arrowshaped toothed, calyx-teeth longer than their tube lanceolate-subulate, pods linear-oblong *hairy*.—*E. B.* 1842. *St.* 32. 5.—*St.* 12—18 in. long. Fl. almost always solitary, purple. "Seeds globose, speckled with black and grey; hile oval." Upper part of the style hairy all over. The inland plants have broader leaflets and more cut *stip.* than those found near the sea.—Bushy places on a gravelly soil, rare. P. VII. VIII. E.

8. *V. sépium* (L.); fl. 4—6 in small axillary nearly sessile clusters, leaflets in 4—8 pairs ovate blunt mucronate gradually smaller upwards on the petiole, stip. half-arrowshaped undivided or lobed, calyx-teeth unequal shorter than their tube, 2 upper ones curved upwards, pods linear-oblong glabrous.—*E. B.* 1515. *St.* 31. 16.—St. about 2 feet high. Fl. purplish. Calyx hairy. Hile linear, extending about  $\frac{3}{4}$  round the seed. Upper part of the style nearly or quite glabrous, bearded. L. more or less hairy, leaflets sometimes ovate-lanceolate truncate.—Woods and hedges. P. VI.—VIII. E. S. I.

9. *V. hybrida* (L.); fl. solitary axillary, leaflets obovate emarginate apiculate in 5—7 pairs, calyx-teeth unequal spreading subulate longer than their tube, standard hairy, pod ovate-oblong hairy.—*E. B.* 482. *St.* 32. 3.—St. ascending, 1 foot long. Fl. reddish yellow. Hairs on the pods simple. "Seeds round with a short hile."—On Glastonbury Tor Hill. P. VI. VII. E.

10. *V. lútea* (L.); fl. solitary axillary, leaflets elliptic-lanceolate acute or rounded at the end apiculate in 5—8 pairs, calyx-teeth unequal, upper ones very short and curved upwards, lower one longer than the tube, standard glabrous, pods elliptic-oblong hairy.—*E. B.* 481. *St.* 31. 13.—St. procumbent, 1—2 feet long. Fl. sulphur-coloured. Hairs on the pods bulbous. Seeds round, compressed, with a short hile. L. varying greatly in hairiness.—Pebbly and sandy ground near the sea. P. VI.—VIII. E. S.

11. *V. sativa* (L.); fl. axillary solitary or in pairs, leaflets in 5—7 pairs elliptic-oblong retuse or obcordate mucronate, upper ones narrower or linear truncate mucronate, calyx-teeth equal lanceolate-subulate long equalling their tube, standard glabrous, pods linear slightly silky, seeds globose smooth.—Seeds slightly compressed; hile linear, occupying about  $\frac{1}{4}$  of the circumference.—*a. V. sativa* (Sm.); leaflets all elliptic- or ovate-oblong, the lower ones shorter and broader, fl. usually in pairs, pods mostly parallel to the st., st. 1—1 $\frac{1}{2}$  foot high. *Sy. E. B.* 392. *St.* 31. 10.—*β. V. angustifolia* (Roth); leaflets of the upper l. linear-lanceolate, lower ones obovate retuse or obcordate, fl. solitary or in pairs, pods mostly patent, st. slender. *E. B. S.* 2614. *St.* 31. 11.—*γ. V. Bobartii* (Forst.); leaflets of the upper l. linear, fl. solitary, pods patent, st. prostrate. *E. B. S.* 2708. *V. angustifolia* Sm.—*a* is only known in cultivation. *β* and *γ* in dry places. A. V. VI. Common Vetch. E. S. I.

\*\*\*\* Style bearded below the stigma; calyx not gibbous.

12. *V. lathyroides* (L.); fl. axillary solitary, leaflets in 1—3 pairs obovate or oblong retuse mucronate, calyx-teeth subulate

straight as long as their tube, pods linear glabrous, *seeds nearly cubical tubercular*, hile short oblong.—*E. B.* 30. *St.* 31. 12.—*St.* procumbent, 3—5 in. long. Fl. small, purple.—Dry gravelly and sandy places. A. V. VI. E. S. I.

#### 14. LATH'YRUS Linn. Vetchling.

\* *No true leaflets.*

1. *L. Aph'aca* (L.); ped. 1-fl., *petioles leafless forming tendrils, stip. very large leaflike* hastate-ovate.—*E. B.* 1167.—*St.* weak, climbing. Fl. yellow, rarely 2 together. Remarkable for its want of l., which are replaced by the stipules, rarely 1 or 2 elliptic leaflets occur. Pods subfalcate; seed smooth, compressed.—Sandy and gravelly fields in the South, rare. A. V. —VIII. E.

2. *L. Nissólia* (L.); ped. long 1—2-flowered, *petioles leaflike* but leafless linear-lanceolate, no *tendrils, stip. minute* subulate.—*E. B.* 112.—*St.* mostly erect. Petioles grasslike. Fl. purple on long stalks. Pods cylindrical; seeds tubercled, round; hile small, oval.—Bushy grassy places, rare. A. VI. E.

\*\* *Petioles with one pair of leaflets and a tendril.*

[*L. sphær'icus* (Retz.); ped. 1-fl. with a long point shorter than the petiole, lts. linear-lanceolate, seeds globose; has been found in Hertfordshire.]

3. *L. hirsútus* (L.); ped. 2-flowered, *lts. linear-lanceolate*, pods hairy, *seeds globose tubercled*.—*E. B.* 1255.—*St.* winged, climbing to the height of 1 or 2 feet. Pods linear-oblong, covered with bulbous hairs. Fl. 2 or 1, pale blue with a crimson standard. Hile oblong.—Rare. Essex. Surrey. A. VI. VII. E.

4. *L. pratén'sis* (L.); *st. angular, ped. many-flowered*, lts. lanceolate mucronate slightly silky, *calyx-teeth subulate*, pods obliquely veined, seeds globose smooth.—*E. B.* 670.—Creeping. *St.* 2—3 feet high, climbing, not winged. Pods linear-oblong, compressed. Fl. racemose, drooping, bright yellow. *Hile small*, oblong.—Moist meadows and pastures. P. VII. VIII. E. S. I.

†5. *L. tuberósus* (L.); *st. angular*, ped. long many-flowered, *lts. obovate-oblong* mucronate, upper *cal.-teeth triangular*, pod net-veined, seeds globose smooth.—*Sy. E. B.* 401.—Creeping and tuberous. *St.* about 2 ft. high, not winged. Fl. 2—5 together, purple. Pods linear-oblong, subcylindric. Hile small.—Hedges and fields. Abundant at Fyfield, Essex. P. VIII. E.

6. *L. sylves'tris* (L.); *st. winged*, ped. many-flowered, *lts. linear-lanceolate* or lanceolate, *calyx-teeth triangular-subulate*, 2 upper ones short, pods net-veined, seeds compressed roughish nearly *half surrounded by the hile*.—*E. B.* 805.—*St.* climbing to the height of 5—6 feet. Pods linear-oblong curved. Fl. greenish yellow variegated with purple. Broader-leaved varieties often pass for *L. latifolius*, which differs by its elliptic pointed lts. and rougher seeds with a shorter hile, but is not a native.—Woods and thickets. P. VII.—IX. E. S.

\*\*\* *Petioles with 2 or more pairs of leaflets and a tendril.*

7. *L. palus'tris* (L.); *st. winged*, ped. long many-flowered, *lts. in 2 or 3 pairs linear-lanceolate acute*, stip. half-arrowshaped lanceolate, pods linear-oblong compressed, seeds globose smooth  $\frac{1}{4}$  surrounded by the hile.—*E. B.* 169.—*St.* 2—3 feet high. Fl. bluish purple.—Boggy meadows, rare. P. VI. VII. E. S. I.

8. *L. marit'imus* (Big.); *st. angular not winged*, ped. short many-flowered, *lts. in 3 or 4 pairs oval*, stip. large oval cordate-hastate, pods oblong obliquely net-veined, seeds globose  $\frac{1}{3}$  surrounded by the hile.—*Pisum* L. *E. B.* 1046.—*St.* prostrate. Leaflets large, blunt but apiculate; petioles often recurved. Fl. purple variegated.— $\beta$ . *acutifolius* (Bab.); leaflets elliptic-lanceolate acute, petioles straight, stems slender straggling.—Pebbly sea-shores, rare.  $\beta$ . Burrafirth, Unst, Shetland. P. VII. VIII. *Sea Pea*. E. S. I.

\*\*\*\* *L. pinnate, without tendrils.* OROBUS.

9. *L. macrorrh'izus* (Wimm.); *st. simple winged*, peduncles 2—4-flowered, l. of 2—3 pairs of oblong or lanceolate blunt apiculate leaflets *without tendrils*, stip. half-arrowshaped broad, pods cylindrical, seeds globose  $\frac{1}{3}$  surrounded by the hile.—*O. tuberosus* L. *E. B.* 1153.—Root tuberous. L. glaucous beneath. Fl. purple, variegated with red and blue.— $\beta$ . *temi-folius*; lts. linear.—Woods and thickets in hilly countries. P. V.—VII. E. S. I.

10. *L. n'iger* (Wimm.); *st. branched not winged*, ped. many-flowered, l. of 3—6 pairs of lanceolate or oblong leaflets *without tendrils*, stip. linear-subulate the lower ones half-arrowshaped, pods slightly compressed, seeds oval  $\frac{1}{3}$  surrounded by the hile.—*E. B. S.* 2788.—Turns black in drying. Fl. variegated with red, blue, and purple. Pods linear. Seed dark brown, perfectly smooth.—Rocky woods in Scotland, rare. P. VI. VII. S.



Tribe III. *Hedysarææ*.15. ORNI'THOPUS *Linn.* Bird's foot.

1. *O. perpusillus* (L.); ped. exceeding the leaves, calyx-teeth triangular acute  $\frac{1}{3}$  the length of their tube, beak scarcely as long as a joint of the pod.—*E. B.* 369.—A small prostrate plant, 3—12 in. long. L. with  $5\frac{1}{2}$ — $12\frac{1}{2}$  pairs of elliptic downy leaflets. Fl. small; calyx hairy, cor. white with crimson veins. Pods curved, joints beadlike wrinkled lengthwise.—Dry sandy and gravelly places. A. V.—VII. E. S. I.

16. ARTHROLOBIUM *Desv.*

1. *A. ebracteatum* (DC.); ped. about equalling the l. 2—4-flowered, stip. minute distinct, l. pinnate with many pairs of elliptic-oblong leaflets, the lowest pair remote from the stem.—*E. B. S.* 2844.—St. prostrate, filiform. Fl. small, yellow, standard red externally. Pod. curved upwards, joints cylindrical rugose.—Channel and Scilly Islands. A. VI. VII. E.

17. HIPPOCREPIS *Linn.* Horseshoe Vetch.

1. *H. comosa* (L.); pods umbellate, their joints rough curved neither dilated nor bordered, joinings glabrous, peduncles longer than the leaves.—*E. B.* 31.—St. procumbent, often a foot long. Fl. yellow. Leaflets 7—13, obovate, blunt or emarginate, apiculate.—Dry calcareous banks. P. V.—VIII. E.

18. ONOBRYCHIS *Gaert.* Sainfoin. Cock's-head.

1. *O. sativa* (Lam.); wings shorter than the calyx, keel about as long as the standard, st. ascending, pods with netted spinous elevations on the disk and short sharp flat teeth on the lower suture.—*E. B.* 96. *St.* 19. 10.—St. often 2 feet long. Fl. in long dense racemes, crimson, variegated. Tube of the calyx silky, short; teeth very long. Leaflets elliptic-oblong, mucronate, entire glabrous above, in about  $12\frac{1}{2}$  pairs.—On chalky and limestone hills. P. VI. VII. E.

## Order XXVI. ROSACEÆ.

Cal. 4—5-parted, or 8—10-parted in 2 rows, free, or adhering to and enclosing the ovary; odd lobe superior. Pet. usually 5, equal, perigynous. Stam. perigynous, usually indefinite. Carp. several or solitary, distinct or connate or adnate to the calyx.

Styles distinct, often lateral. Fruit various. Seeds nearly without albumen, embryo straight.—L. alternate, usually compound, with stipules.

### Suborder I. AMYGDALÆÆ.

*Fruit a drupe.* Calyx deciduous, inferior, quite free from the solitary ovary. Stipules free.

1. PRUNUS. Drupe fleshy, indehiscent; its nut smooth or furrowed.

### Suborder II. ROSEÆÆ.

Carpels several, distinct from each other and from the calyx. Stipules united to the petiole.

Tribe I. *SPIRÆIDÆ.* *Carpels* (follicles) several, *not included in calyx-tube*; seeds 1—6, suspended from the inner edges of the follicles. Sep. persistent, in one row.

2. SPIRÆA. Cal. 5-cleft. Stam. many, inserted with the pet. on a disk adhering to the calyx. Follicles 1 or more, usually distinct. Seeds 2—6.

Tr. II. *SANGUISORBEÆ.* *Carpels* 1—3, *enclosed in the dry calyx-tube*, which is narrowed at the top.

3. SANGUISORBA. *Fl. perfect.* Cal. 4-cleft, with 2 or 3 scales at its base, tube quadrangular. Pet. 0. *Stam.* 4, opposite to the segments of the calyx. Nuts 1—2. Style terminal. Stigma capitate, covered with oblong spreading prominences. Seed suspended.
4. POTERIUM. *Fl. monœcious or polygamous.* Cal. 4-cleft, with 3 scales at its base, tube quadrangular. Pet. 0. *Stam.* 20—30. Nuts 2—3. Style terminal. Stigma brush-like, with filiform divisions. Seed suspended.
5. AGRIMONIA. *Fl. perfect.* *Calyx 5-cleft, without scales; tube turbinate, armed with hooked bristles above.* Pet. 5. *Stam.* 12—20, inserted with the pet. into a glandular ring in the throat of the calyx. Nuts 1—3. Style terminal. Seed suspended.
6. ALCHEMILLA. *Fl. perfect.* *Cal. 8-parted, the alternate parts smaller, contracted at the throat, unarmed.* Pet. 0. *Stam.* 1—4, inserted into a ring in the throat of the calyx and opposite to the smaller segments. Nuts 1—5. *Style basal.* Seeds ascending.

Tr. III. *DRYADEÆ*. Fruit not included in calyx-tube, of 5 or more small dry nuts (in *Rubus* drupes) inserted on a dry or succulent receptacle. Calyx persistent, open, nearly flat.

\* *Attachment of the seed near to that of the style, radicle superior.*

7. *SIBBALDIA*. *Cal.* concave, 10-parted, in 2 series, 5 exterior parts smaller. *Pet.* 5. *Stam.* 5. Style lateral. *Fr.* of 5—10 small nuts placed on a dry receptacle. Seed ascending.

8. *POTENTILLA*. *Cal.* concave, 8—10-parted, in two series, 5 exterior parts smaller. *Pet.* 4—5. *Stam.* many. Style lateral or nearly terminal. *Fr.* of many small nuts placed upon a flattish dry receptacle. Seed pendulous or ascending.

9. *COMARUM*. *Cal.*, *cor.*, *stam.*, and pistils as in *Potentilla*. *Receptacle* ultimately large, spongy, persistent. Style lateral near the top of the nut. Seed pendulous.

10. *FRAGARIA*. *Cal.*, *cor.*, *stam.*, and pistils as in *Potentilla*. *Receptacle* large, succulent, pulpy, deciduous. Style lateral near the base of the nut. Seed ascending.

11. *RUBUS*. *Cal.* concave or flattish, 5-parted. *Pet.* 5. *Stam.* many. Styles nearly terminal. *Carp.* many, succulent (drupes), placed upon a hemispherical or conical spongy receptacle. Seed pendulous.

\*\* *Attachment of the seed distant from that of the style, radicle inferior. Nuts with long awnlike styles.*

12. *DRYAS*. *Cal.* 8—10-cleft, in one row. *Pet.* 8—9. *Stam.* many. *Fr.* of many small nuts, tipped with the persistent hairy styles, which are not jointed. Receptacle flat, dry. Seed ascending.

13. *GEUM*. *Cal.* 10-cleft, in 2 rows, the outer parts smaller. *Pet.* 5. *Stam.* many. *Fr.* of many small nuts, tipped with the persistent jointed styles hooked at the joining. Receptacle elongated, dry. Seed ascending.

Tr. IV. *ROSIDÆ*. Fruit formed of many small dry nuts enclosed in the fleshy calyx-tube.

14. *ROSA*. *Cal.* urceolate, contracted at the mouth, ultimately fleshy, 5-fid. *Pet.* 5. *Stam.* many, inserted with the petals on the rim of the tube of the calyx.

## Suborder III. POMEÆ.

Calyx-tube thick and fleshy, in fruit adhering to the carpels and forming a 1—5-celled pome: thus fl. appearing superior.

15. CRATÆGUS. Calyx-segments 5, acute. Pet. 5. Styles 1—5. Fr. oval or round, *concealing the upper end of the 1—5 bony 1—2-seeded carpels.*
16. COTONEASTER. Calyx-segments 5. Pet. 5. Styles 2—5. Fr. turbinate, *its nuts adhering to the sides of the calyx but not cohering at the centre.*—Stam. erect, as long as the teeth of the calyx.
17. MESPILUS. Calyx-segments 5, leaflike. Pet. 5. Styles 2—5. Fr. turbinate with *the upper end of the bony carpels exposed*; disk dilated, almost as broad as the fruit.
18. PYRUS. Cal. 5-toothed. Pet. 5. Styles 2—5. Fr. *fleshy with 5 cartilaginous or membranous distinct 2-seeded cells.* Testa cartilaginous.

## Suborder I. Amygdaleæ.

## 1. PRU'NUS.

\* *Young leaves convolute. Drupe glaucous.*

1. *P. communis* (Huds.); ped. solitary or in pairs, l. elliptic or ovate-lanceolate rather downy beneath.—*a. P. spinosa* (L.); branches spinous, ped. glabrous, l. usually glabrous, fr. globose. *Sy. E. B.* 408.—*β. P. insititia* (L.); branches slightly spinous, ped. and underside of l. usually downy, fr. globose. *E. B.* 841.—\**γ. P. domestica* (L.); branches without spines, ped. glabrous, l. hairy about the midrib beneath, fr. oblong. *E. B.* 1783.—*a* is a shrub with crooked and much armed black branches and *fl. before the leaves*; *β* a taller shrub with straight and slightly armed brown branches and *fl. and l. usually together*; *γ* a small tree with straightish branches.—Thickets. *γ* not indigenous. *Sh. IV. V. a. Sloe. Blackthorn. β. Bullace. γ. Plum. E. S. I.*

\*\* *Young leaves conduplicate. Drupe not glaucous.* CERASUS.

2. *P. Pádus* (L.); arborescent, l. obovate-lanceolate finely serrate glabrous, *fl. in pendulous racemes*, fr. roundish-oblong.—*E. B.* 1383.—A small tree. L. minutely doubly serrate. Fl. white, many, in a lax raceme. Fr. black, harsh, bitter; nut wrinkled.—Woods and hedges. *T. V. Bird-Cherry. E. S. I.*

3. *P. A'vium* (L.); arborescent, *l. long-stalked drooping* oblong-ovate suddenly cuspidate incise-serrate downy beneath, calyx-tube constricted below the entire sepals, pet. flaccid, "fr. heart-shaped."—*E. B.* 706.—A tree 20—30 feet high. Outer scales of the leaf-buds deflexed. Flower-buds not leafy. Fl. in umbels. Pet. bifid, with a minute claw. "Fr. firm, bitter, black or red." *Borr.*—Woods. T. V. *Wild Cherry.* *Gean.* E. S. I.

4. *P. Cer'asus* (L.); fruticose, *l. short-stalked not drooping* oblong-ovate or ovate-lanceolate doubly crenate-serrate glabrous, calyx-tube not constricted, "pet. firm," fruit round.—*E. B. S.* 2863.—An erect bushy shrub 3—8 feet high. Umbels scattered. Outer scales of the leaf-buds erect. Inner scales of the flower-buds leaflike. Sep. crenate-serrate. Pet. submarginate, with a claw. "Fr. juicy, acid, always red."—Hedges. Sh. V. *Dwarf Cherry.* E. I.

## Suborder II. *Rosæ.* Tribe I. *Spiræidæ.*

### 2. SPIRÆA *Linn.*

‡1. *S. salicifolia* (L.); shrubby, *stip.* 0, *l.* elliptic-lanceolate unequally serrate glabrous, racemes terminal compound, *stam.* exceeding the petals.—*E. B.* 1468.—A shrub 4—5 feet high with smooth round wandlike branches. Fl. flesh-coloured, in dense erect racemes.—Damp places in the North and in Wales, rare. Sh. VII. E. S.

2. *S. Ulm'aria* (L.); herbaceous, *stip.* rounded toothed, *l.* interruptedly pinnate, leaflets ovate undivided, the terminal one larger palmately 3—5-lobed, fl. in compound cymes, caps. glabrous twisted together.—*E. B.* 960. *St.* 18. 8.—*St.* about 3 feet high, angular, branched. L. with a few large serrate leaflets and very minute intermediate ones, downy beneath. Cymes with long side branches. Fl. yellowish, sweet-scented. Pet. roundish.—Meadows and by water. P. VI.—VIII. *Meadow-sweet.* E. S. I.

3. *S. Filipen'dula* (L.); herbaceous, *stip.* of the root-leaves linear acute entire, those of the stem rounded and cut, *l.* interruptedly pinnate, leaflets all oblong deeply cut and serrate, fl. in a paniced cyme, caps. hairy, straight but adpressed.—*E. B.* 284. *St.* 18. 7.—Root producing knobs. *St.* 1—1½ foot high, round, simple, paniced at the top. L. mostly radical, spreading; leaflets small, many, intermediate ones small. Fl. yellowish-white

tinged with red. Pet. obovate.—Dry chalky and limestone pastures. P. VI. VII. *Dropwort*. E. S. I.

### Tribe II. *Sanguisorbea*.

#### 3. SANGUISOR'BA Linn. Great Burnet.

1. *S. officinalis* (L.); spikes ovate-oblong, stam. about as long as the calyx, leaflets cordate-oblong.—*E. B.* 1312.—L. pinnate, glabrous; leaflets about 13, stalked, opposite, blunt, coarsely serrate. Spikes rarely long, cylindrical. Fr. oblong, winged chiefly in its upper half, transverse section terete, 4 wings thin.—Damp meadows. P. VI.—VIII. E. S. I.

#### 4. POTE'RIMUM Linn. Lesser Burnet.

1. *P. Sanguisor'ba* (L.); herbaceous, st. slightly angular, cal. of the fruit hardened *quadrangular with 4 thin entire wings and netted veins*.—*E. B.* 860.—L. pinnate, with many small ovate coarsely serrate subsessile leaflets glabrous or slightly hairy beneath. Lower part of the stems and petioles often downy.—On a dry calcareous soil. P. VI.—VIII. E. S. I.

‡2. *P. muricatum* (Spach); herbaceous, st. slightly angular, calyx of fruit hardened *4-winged pitted*, pits with elevated and *denticulate* edges.—*E. B. S.* 2989.—*a. P. platylophum* (Jord.); fr.-cal. with rather blunt denticulate wings, pits with sharply denticulate edges.—*β. P. stenolophum* (Jord.); fr.-cal. with sharp entire wings, pits with rather bluntly denticulate edges.—Both of these closely resemble Sp. 1, but are usually larger in all their parts. Leaflets usually oval.—Dry places. P. VI. VII. E.

#### 5. AGRIMO'NIA Linn. Agrimony.

1. *A. Eupatoria* (L.); *cal.-tube of fr. obconic* furrowed to the base, *exterior spines spreading*, l. interruptedly pinnate serrate shaggy beneath.—*E. B.* 1335. *St.* 59. 4.—*St.* erect, about 2 feet high. Spikes long with distant yellow flowers. Leaflets deeply serrate.—Fields and roadsides. P. VI. VII. E. S. I.

2. *A. odorata* (Mill.); *cal.-tube of fr. bellshaped* not furrowed, *exterior spines declining*, l. interruptedly pinnate hairy and with minute glands beneath.—*E. B. S.* 2982.—Scented. Spikes long. Fl. yellow; pet. obovate-lanceolate, wedged below, distant, spreading, flat, turning up at the end. Styles spreading. Leaflets deeply and sharply serrate throughout. Cal.-tube of

the arge fruits rarely with shallow furrows in its upper half. Taller than Sp. 1, usually more branched; l. and lts., fl. and fr. larger.—Waste places, rare. P. VI. VII. E. S. I.

### 6. *ALCHEMILLA* Linn. Lady's Mantle.

1. *A. vulgaris* (L.); *l. reniform* or nearly orbicular plaited 7—9-lobed, lobes rounded serrate throughout green beneath, fl. in terminal corymbs.—*E. B.* 597.—Slightly hairy. Fl. yellowish green. L. large on long stalks, st.-l. sessile with large notched connate stipules.— $\beta$ . *A. montana* (Willd.); st.-l. and petioles silky, old l. wavy with broad waves.—Moist pastures. P. VI.—VIII. *Common Lady's Mantle.* E. S. I.

2. *A. alpina* (L.); *radical l. digitate, divisions 5—7 separated* to their base oblong blunt adpressed serrate at the end white and silky beneath, fl. in interrupted spikes of small lateral and terminal corymbs, achene oblong-ovoid suddenly acuminate, st. slightly branched simple below.—*Sy. E. B.* 425. *St.* 51. 2.—*St.*, cal. and underside of the l. beautifully silky. Leaflets rarely slightly combined, *outer ones* of the radical l. usually nearly *opposite* to each other. Branches usually undivided, ascending.—Mountains. P. VI. VII. *Alpine Lady's Mantle.* E. S. I.

3. *A. conjuncta* (Bab.); *radical l. suborbicular peltate-palmate, divisions 5—7 much connected below oblong blunt adpressed serrate at the end white and very silky beneath, fl. in small lateral and terminal corymbs, achene ovate-ovoid gradually acuminate, st. much branched.*—*E. B. S.* 2983. *A. argentea* (Don); not *Lam.*—Often taken for *A. alpina* but its lts. are much larger, their lobes broader and placed in the radical leaves so that *the 2 external ones almost if not quite touch each other so as to present the appearance of a peltate leaf*; st.-branches long alternate spreading and often again subdivided; fl. in small nearly simple distant corymbs. [Faroe Isles. *A. fissa* Fl. Dan. 2101. Dauphine.] Clova! Glen Sannox, Arran! P. VI. VII. S.

4. *A. arvensis* (L.); *l. palmate 3-fid wedgeshaped below hairy, lobes with 3—6 teeth at the end, fl. sessile axillary.*—*Aphanes* Linn. *E. B.* 1011.—*St.* prostrate or ascending, 4—5 in. long. Fl. very small, greenish, in small hairy inconspicuous tufts.—Dry fields on sand and gravel. A. V.—VIII. *Parsley Piert.* E. S. I.

Tribe III. *Dryadeæ*.7. SIBBAL'DIA *Linn.*

1. *S. procumbens* (L.); l. ternate, leaflets wedgeshaped with 3 teeth at the end, fl. corymbose, pet. lanceolate.—*E. B.* 897. *St.* 17. 5. *Pot. Sibbaldii*, Hall.—L. pilose on both sides. St. woody, procumbent. Pet. very small, yellow. Pistils and stam. very variable in number.—Dry summits of Scottish mountains. Above Highcup Scar, Teesdale. P. VII. E. S.

8. POTENTIL'LA *Linn.* Cinque-foil.

\* *Hairs on the receptacle shorter than the glabrous carpels.*

† Leaves pinnate.

1. *P. rupes'tris* (L.); *st. erect* dichotomous, leaflets roundish-ovate unequally cut and serrate 5—7 on the lower leaves, on the uppermost 3.—*E. B.* 2058.—Fl.-shoots annual. *Pet. white*, much longer than the calyx. Base of l. wedgeshaped. *St.* 1—2 feet high.—On Craig Breidden, Montgomeryshire. P. V. VI. E:

2. *P. anser'ina* (L.); *st. creeping*, l. interruptedly pinnate, leaflets many oblong acutely serrate silky beneath, peduncles solitary.—*E. B.* 861. *St.* 4. 7.—*Fl. yellow*, large. L. green above, white and silky beneath; or white and densely silky on both sides.—Roadsides. P. VI. VII. *Silver Weed.* E. S. I.

†† Leaves digitate.

3. *P. argen'tea* (L.); *st. ascending*, l. *quinate*, leaflets obovate-cuneate incise-serrate *white and downy beneath* their margins revolute.—*E. B.* 89. *St.* 17. 7.—*Fl. yellow*, small, in terminal corymbs.—Dry gravelly places. P. VI. VII. E. S.

4. *P. ver'na* (L.); *st. prostrate*, lower l. of 5—7 *obovate leaflets* serrate towards the end bristly on the margin and ribs beneath, teeth 2—4 on each side, *lower stipules narrowly linear.*—*E. B.* 37. *St.* 17. 8.—St. woody, about 5 in. long. *Fl. yellow*, solitary or 2 or 3 together. The terminal tooth of the l. usually smallest and shortest.—Dry pastures. P. IV. V. E. S.

5. *P. macul'ata* (Pourr.); *st. ascending*, lower l. *quinate*, leaflets obovate-cuneate somewhat hairy deeply cut in the upper half, teeth about 4 on each side, *stip. all ovate.*—*P. salisburgensis* (Haenke). *St.* 17. 10. *Sy. E. B.* 429.—Larger than the pre-



ceding. All the teeth of l. equal. Outer sep. oblong, blunt; inner twice as broad, ovate, acute.—Mountains. P. VI. VII.

E. S.

6. *P. reptans* (L.); *st. filiform* prostrate *rooting*, l. quinate stalked, leaflets obovate bluntly serrate, peduncles solitary, pet. obcordate, *carpels asperous*.—*E. B.* 862.—L. on long stalks, often with a bunch of small l. in their axils, sometimes solitary, usually in pairs. Leaflets blunt, rough or hairy on their ribs and margins. Fl. on long stalks, yellow.—Sometimes the l. and cal. are covered with long silky hairs on both sides.—Roadsides and banks. P. VI.—IX.

E. S. I.

7. *P. Tormentilla* (Nesl.); *st. procumbent* or ascending, l. ternate sessile or shortly stalked, lower l. quinate on long stalks, lts. lanceolate or obovate-cuneate incise-serrate, pet. obcordate, *carpels longitudinally wrinkled*.—*Tormentilla officinalis* L. *E. B.* 863. *St.* 34. 12.—Rootstock large, woody. L. all nearly sessile, except the lowest, which are often ternate; lts. acute, rather hairy. Stip. deeply cut. Fl. small, yellow, usually with 8 sep. and 4 pet.—*P. mixta* (Nolte); l. stalked, lts. obovate-oblong acutely serrate, stip. entire. A hybrid between Sp. 6 and 7.—*β. P. procumbens* (Sibth.); leaflets obovate-cuneate deeply cut, l. 5-nate or 3-nate stalked, stip. entire or trifid. Fl. usually larger. *Sy. E. B.* 431. *T. reptans* L.—In dry places. *β.* Woods and hedgebanks. P. VI.—VIII.

E. S. I.

\*\* *Hairs on the receptacle long, carpels hairy at the scar or all over.*

8. *P. fruticosa* (L.); *shrubby*, l. *pinnate*, leaflets mostly 5 oblong acute entire hairy with revolute margins.—*E. B.* 88.—*St.* 3—4 feet high. Fl. large, yellow, terminal.—Teesdale. Wastdale. Galway. Clare. Sh. VI. VII.

E. I.

9. *P. fragarias'trum* (Ehrh.); *st. procumbent*, l. *ternate* greyish green, *leaflets roundish obovate serrate* silky on both sides, pet. narrowly obcordate not contiguous, carp. glabrous except at the scar smooth or wrinkled transversely.—*E. B.* 1785.—Hairs on the upperside of l. bulbous-based. Fl. small, white.—Woods, banks. P. IV. V. *Barren Strawberry*.

E. S. I.

[*P. Norvégica* (L.). Mr. G. S. Gibson found this in Burwell Fen, Cambridgeshire, in 1868! *J. of B.* vi. 302.]

## 9. COM'ARUM Linn. Marsh Cinque-foil.

1. *C. palus'tre* (L.).—*E. B.* 172.—*St.* ascending, 1 foot high,

reddish. L. pinnate. Leaflets 5—7, elliptic-oblong, acute, sharply serrate. Fl. several, dark purple; cal. purple within; pet. small. Differing from *Potentilla* by its enlarged spongy receptacle.—Marshes and peaty bogs. P. VII. E. S. I.

### 10. FRAGA'RIA Linn. Strawberry.

1. *F. ves'ca* (L.); cal. of the fruit spreading or reflexed, *hairs* on the peduncles spreading those of the *pedicels adpressed upwards* silky.—*E. B.* 1524. *E. B. S.* 2742.—Stoloniferous. Stole continued by an axillary shoot at each rosette (a sympode); one scale between each two rosettes. L. bright green. Flowering stems short, erect, mostly simple, few-flowered. Lts. sessile. Hairs on pedicel of first fl. spreading, on the underside of l. adpressed. Carp. smooth, glabrous, on all parts of the receptacle, superficial. Pet. about as long as broad, white throughout with 2 slight notches at the end, contiguous; claw indistinct.—Woods and thickets. P. V. VI. *Wood Strawberry.* E. I. S.

[*F. collina* (Ehrh.) has a stole formed by a continuous axis; no scale except between its base and the first rosette.]

†2. *F. elátior* (Ehrh.); cal. of the fruit spreading or reflexed, *hairs* on the peduncles and *pedicels spreading and somewhat deflexed*.—*E. B.* 2197. *F. moschata* Lindl.—Fl. imperfectly dioecious. Pet.  $\frac{1}{2}$  broader than long, white, entire; claw distinct, bright yellow. “Base of receptacle without carpels.” Larger and more hairy than *F. vesca*.—Woods, rare. P. VI.—IX. *Hautboy Strawberry.* E. S.

### 11. RU'BUS Linn.<sup>1</sup> Bramble.

#### A. FRUTESCENTES.

Stem shrubby. Leaves subquinate. Stipules linear, affixed to the petioles. Flowers subpanicked. Succulent carpels forming a compound many-seeded berry. Receptacle conical.

<sup>1</sup> In the descriptions by *stem* is meant the *barren stem* of the year; the prickles are called *patent* when they spread at right angles to the st. and *subpatent* when a little declining; the shape of the leaflets, unless otherwise stated, is taken from those of the barren stem. The term *seta* is used to express a *hair or bristle tipped with a gland*; *aciculæ* are *strong bristles*.—By *R. G.* the plates in Weille and Nees's *Rubi Germanici* are intended.—See Babington's *British Rubi*.

When the continental plants are better known it is feared that considerable changes of nomenclature will be necessary.

- i. *Idæi*. Stems suberect, biennial. Ripe fruit separating from its receptacle.—Leaves often pinnate.

1. *R. Idæ'us* (L.); st. erect round pruinose, prickles setaceous straight, l. 5-pinnate or ternate white beneath, term. lt. long-stalked, interm. lts. sessile not imbricate, prickles of fl.-shoot and ped. deflexed, fl. axillary and terminal corymbose.—*E. B.* 2442. *R. G.* 47.—Creeping. St. 4—6 ft. high; prickles small, usually many. L. usually pinnate, rarely ternate. Fr. red or amber-coloured.—*β. R. Leesii* (Bab.); l. ternate, lts. all roundly ovate subsessile imbricate. *E. B. S.* 2981.—*γ. rotundifolius*; l. like those of *β* but term. lt. long-stalked, l. of flowering shoot similar but upper ones simple.—Damp edges of woods and heaths. Sh. VI. E. S. I.

- ii. *Fruticosi*. Stems biennial or subperennial. Ripe fruit not separating from its receptacle. Leaves digitate, pedate or rarely subpinnate.—*R. fruticosus* *Hook.*

- (1) *Suberecti*. Stems usually suberect, glabrous or slightly pilose, not setose nor felted. Prickles equal.—*Sepals* densely white-felted within, pilose externally *with a narrow border of white felt.*

2. *R. suberec'tus* (Anders.); st. erect obtuse-angled at the top, prickles few small uniform from a large compressed base confined to the angles of the stem, l. 3—5—7-nate, lts. flexible flat, term. lt. cordate-acuminate, basal lts. subsessile *those of fl.-shoot narrowed to the base*, fl. racemose or subpanicked, sep. reflexed.—*E. B.* 2572.—St. 3—6 ft. high. Prickles conical, scarcely longer than the longitudinal extent of their base. L. rarely ternate; lts. thin, unequally serrate. Stam. pale, exceeding styles. Fr. dark red.—Boggy woods and heaths. Sh. VI. VII. E. S. I.

3. *R. fis'sus* (Lindl.); st. erect or subarcuate obtuse-angled, prickles many straight or deflexed *from an oblong slightly dilated base not confined to the angles*, l. 5—7-nate, lts. coriaceous plicate, term. lt. cordate-ovate, basal lts. sessile, lateral lts. of fl.-shoot often gibbous at the base, pan. simple racemose-corymbose, *fr.-cal. erect-patent.*—Creeping extensively. St. 1—2 ft. high. Prickles much longer than the longitudinal extent of their base. Lts. unequally serrate. Stam. slightly exceeding styles. Fr. dark red.—Wet places. Sh. VI.—VIII. E. I.

4. *R. plicátus* (W. & N.); st. suberect obtuse-angled, prickles straight or deflexed from an oblong dilated base confined to the angles, l. quinate, lts. more or less *plicate thin pilose not felted beneath*, term. lt. cordate-acuminate, basal lts. usually subsessile,

*lateral lts. of fl.-shoot rhomboidal-ovate dilated at the base, fl. racemose or corymbose, rachis and ped. pilose not felted, fr.-cal. reflexed.*—*E. B. S.* 2714. *R. G.* 1. *R. fruticosus Arrh.*—St. rarely 4 ft. high. Prickles unequal, usually conical, much longer than the longitudinal extent of their base. Upper l. sometimes pinnate-septenate. Cal. bright green externally. Stam. falling short of styles.—Heaths. Sh. VI. VII. E. S. I.

5. *R. affinis* (W. & N.); st. suberect or subarcuate angular smooth, prickles strong slightly deflexed or declining from a dilated compressed base confined to the angles, l. quinate, *lts. coriaceous wavy towards the end green and opaque on both sides subpilose above paler and silky-pubescent beneath*, term lt. cordate-oval cuspidate, basal lts. stalked *those of the fl.-shoot narrowed at the base, branches of the compound leafy pan. corymbose erect-patent often long, sep. acuminate greenish white-felted with pale edges externally patent.*—*R. G.* 3.—St. often arching, but apparently it does not root at the end. Stam. exceeding styles.—*β. R. lentiginosus* (Lees); prickles declining, lts. narrower and nearly glabrous; pet. very small, cal. erect-patent.—Heaths and open woods. Sh. VII. VIII. E. S. I.

6. *R. hemistémon* (Müll.); st. arcuate angular nearly or quite glabrous with subsessile glands, prickles slender short declining from a short oblong base, l. quinate, *lts. coriaceous irregularly dentate green on both sides paler with many short hairs beneath*, term. lt. oblong-ovate acuminate, *basal lts. very shortly stalked those of the flowering shoot rather dilated at the base*, pan. oblong abrupt often nearly subracemose with long 1—2-fl. branches, term. fl. nearly sessile, rachis and peduncles hairy not felted, *sep. ovate-acuminate greenish subtomentose clasping the fruit.*—St. often arching. Prickles rather unequal. Stam. white falling short of "pale green" styles.—Hedges and thickets. Warw., Card., Aberd. Sh. VII. VIII. E. S.

(2) *Rhamnifolia*. Stems arching more or less, rooting at the end, slightly pilose, not setose nor felted nor glaucous. Prickles usually confined to the angles of the stem, nearly equal, from a depressed and compressed base.

7. *R. Lindleiánnus* (Lees); st. erect-arcuate smooth shining prickles strong declining compressed dilated below, l. quinate, *lts. subcoriaceous shining above pale green pilose (often slightly felted) beneath*, term. lt. obovate or roundish acuminate, basal lts. stalked not overlapping the interm. lts., *branches of the compound leafy pan. patent or divaricate short corymbose, rachis polished most prickly in the middle lts. top and the pedicels felted lts. thorns strong declining.*—*R. nitidus* Bell-Salt., Bab. (not W.

& N.).—St. angular throughout, appearing as if varnished. Term. lt. often much narrowed below. Pan. often very compound, close, usually long, blunt and convex at the end. Pet. not contiguous, oblong, white. Stam. slightly exceeding styles; both greenish. Fr. small.—Hedges and borders of thickets. Sh. VII. VIII. E. S.

8. *R. rhamnifolius* (W. & N.); st. arcuate angular furrowed upwards, prickles strong patent or declining, l. quinate, lts. coriaceous flat opaque above greenish-white-felted beneath, term. lt. ovate or cordate subcuspidate, basal lts. stalked not imbricate, pan. felted often dense and blunt at the end with axillary racemose few-flowered distant branches and strong declining prickles.—*E. B. S.* 2604. *R. G.* 6. *R. cordifolius R. G.* 5.—St. usually bright red. Prickles yellow or tipped with red. Lts. hard but felted beneath, finely serrate. Sep. dull green externally. Pet. roundish, white. Stam. exceeding styles; both greenish. Primordial fr. oblong.—Hedges and thickets. Sh. VII. VIII. E.

9. *R. incurvatus* (Bab.); st. arcuate-prostrate angular, prickles strong patent or declining, l. quinate concave, lts. coriaceous acuminate with incurved wavy edges shining and subglabrous above greenish-white-felted beneath, term. lt. cordate-ovate, pan. narrow leafy below with short approximate patent corymbose branches its top and pedicels hairy and felted its prickles slender deflexed, sepals ovate acuminate.—St. purple, strong. Lts. flat except at the edges, doubly dentate, soft beneath. Pet. roundly obovate, pink. Stam. exceeding styles; both pink. Primordial fr. almost hemispherical.—Heaths and open woods. Sh. VII. E. S.

10. *R. imbricatus* (Hort); st. arcuate-prostrate angular, prickles small strong declining from a long compressed base, l. convex quinate, lts. convex coriaceous opaque and subglabrous above paler with scattered hairs beneath cuspidate imbricate, term. lt. roundly cordate-obovate, pan. narrow leafy below with ascending distant long racemose branches its top and pedicels hairy scarcely felted its prickles slender declining, sep. abruptly cuspidate.—St. purplish red. "Basal lts. overlap the intern., intern. the term. leaflet." Pet. obovate, white. Styles greenish yellow. Primordial fr. subglobose.— $\beta$ . *R. ramosus* (Blox.); st. erect-arcuate, l. not imbricate broadly ovate shining.—Top of pan. felted; pet. pink. Fil. white, styles brownish pink. *J. of B.* ix. 330.—By the Wye below Monmouth.  $\beta$ . Devon, Cornw., and Warw. Sh. VI. VII. E.

11. *R. latifolius* (Bab.); st. arcuate-prostrate angular furrowed, prickles small slender compressed slightly declining from a very long compressed base, l. quinate, lts. thin pilose on both

sides coarsely and doubly dentate not felted beneath, *term. lt. cordate acuminate*, basal lts. sessile imbricate, pan. short leafy hairy with ascending few-flowered corymbose branches its top and pedicels felted and hairy its prickles slender short declining.—St. strong, green. Lts. very broad. Pet. ovate, clawed. Primordial fruit apparently about hemispherical.—Open woods. Cramond Bridge and Colinton near Edinburgh. Acharn, Perthshire. Sh. VII. VIII. S.

- (3) *Villicaules*. Stems arching more or less, rooting, pilose or bald, often felted, having subsessile glands and rarely a few setæ and aciculi. Prickles confined to the angles of the stem, nearly equal, or a few smaller scattered on the faces. Basal leaflets stalked, not overlapping the intermediate leaflets.

- a. *Discolores*. St. with strong equal prickles and closely adpressed pubescence. Leaves white-felted beneath.

12. *R. dis'color* (W. & N.); *st. arcuate-prostrate* angular furrowed *stellately downy*, prickles declining or deflexed from a much dilated compressed base, l. quinate, lts. coriaceous convex rugose above *finely white-felted* beneath, *term. lt. obovate-cuspidate*, pan. long narrow felted its few lower branches axillary many-flowered its prickles strong hooked, *cal. finely white-felted*.—*R. G.* 20. *R. fruticosus Sm. E. B.* 715.—St. nearly prostrate unless supported, often nearly glabrous and glaucous. Lts. usually hairless but with fine hard felt beneath, usually with decurved edges. Pet. pink. Fil. and styles purple; anth. greenish; *stam. exceeding styles*. Drupes many, small, acid.—*β. pubigerus*; *st. angular stellately hairy* with spreading hairs, prickles slender from a dilated oblong depressed scarcely compressed base patent or deflexed.—Hedges and thickets. Common. Sh. VII. VIII. E. S. I.

13. *R. thyrsoideus* (Wimm. ?); *st. erect-arcuate* angular furrowed *subglabrous*, prickles declining or deflexed from a much-dilated compressed base, l. quinate, lts. flat rather coriaceous glabrous above *hairy and greenish-white-felted beneath*, *term. lt. cordate-ovate-acuminate*, pan. long narrow its lower branches many axillary few-flowered its prickles strong hooked, *cal. hairy and felted*.—*R. fruticosus R. G.* 7. *R. candicans Focke*.—St. arching highly, self-supporting. Lts. wavy and a little turned up at the edge, hairy and softly but often finely felted beneath. Pet. white. Fil. white; anth. rather fuscous. Drupes few, subacid.—Hedges and thickets. Sh. VII. VIII. E. I.

- b. *Sylvatici*. Stem with moderate mostly equal prickles and (often dense) hairy-woolly patent (but often deciduous) pubescence. Leaves usually green but sometimes white-felted beneath.

14. *R. leucostachys* (Sm.); *st. arcuate-prostrate* angular hairy woolly and felted, prickles many subpatent slender from a dilated compressed base, l. quinate, *lts.* flat hairy and shining and *softly yellowish or whitely felted beneath*, *term. lt.* obovate ovate or roundish cuspidate, *pan. long* narrow felted hairy setose its branches short few-flowered its prickles slender declining or angularly deflexed, cal. green-felted hairy setose aciculate.—*a*; *st. arcuate-prostrate*, prickles mostly on the angles of the stem equal, *lts.* coriaceous obovate sublobate-serrate yellowish-white-felted and shining beneath.—*E. B. S.* 2631.—*St.* nearly prostrate, covered with loose spreading clustered hairs. *Pan.* very narrow, long. *Pet.* pinkish. *Fr.* purplish black.—*β. R. vestitus* (Weihe); *st. arcuate*, prickles unequal scattered, *lts.* cordate-roundish cuspidate irregularly dentate pale green beneath. *R. G.* 33. *St.* often nearly round and bearing a few aciculi and setæ. *Lts.* rather thin.—Hedges, thickets and woods. Sh. VII. VIII. E. I.

15. *R. hirtifolius* (Müll.); *st. arcuate-prostrate* angular pilose, prickles nearly equal small declining from a dilated compressed base, l. quinate, *lts.* flat pale green and pilose only on the veins beneath, *term. lt.* obovate acuminate, panicle narrowly pyramidal setose its branches few-flowered its prickles few slender, sepals greenish felted and with pale edges externally, *stam.* exceeding the styles.—*Pan.*-branches often 1-fl. *Sep.* ovate, leaf-pointed.—*R. pyramidalis* Kaltenh. ?—Bushy and heathy places. Near Plymouth. Isle of Wight. Sh. VII. VIII. E.

16. *R. Grabows'kii* (Weihe); *st. arcuate* angular subglabrous, prickles many equal declining or deflexed much dilated and compressed below, l. quinate, *lts.* plicate opaque and glabrous above ashy-felted beneath irregularly dentate imbricate, *term. lt.* cordate abruptly cuspidate (on the fl.-shoot much dilated below), *pan.* long leafy below its branches ascending racemose-corymbose its prickles many deflexed, cal. ashy-felted and hairy.—*R. carpinifolius* Borr. *E. B. S.* 2664 ? *R. montanus* Focke—*St.* with a few often clustered hairs. *Pan.* hairy but not felted. "Stam. white scarcely exceeding the green styles." *Fr.* sparingly produced.—Hedges and thickets. Sh. VII. VIII. E.

17. *R. Coleman'ni* (Blox.); *st. arcuate* angular subglabrous, prickles many nearly equal declining compressed and dilated below, l. quinate, *lts.* convex opaque above green and hairy on the veins beneath irregularly dentate imbricate, *term. lt.* roundly

cordate acuminate (on the fl.-shoot roundish or broadly oval), pan. long leafy below its branches ascending corymbose or the axillary ones racemose its prickles many slender deflexed or declining with many setæ and aciculi, sep. ashy-felted hairy.—*R. fusco-ater*  $\beta$ . *Colemanni* Bab.—St. with a few aciculi and setæ and scattered hairs. Pan. not felted. Pet. white. The position of this plant is doubtful; but it seems to be most allied to *R. Grabowskii* of our species.—Hedges near Coventry and Packington. Sh. VII. VIII. E.

18. *R. Salteri* (Bab.); *st. arcuate-prostrate angular furrowed subglabrous, prickles subpatent slender* compressed from a dilated compressed base, l. quinate, *lts. thin coarsely and doubly patentely dentate* green on both sides hairy only on the veins beneath, term. lt. broadly obovate cuspidate-acuminate subcordate, pan. long lax hairy its ultra-axillary branches short few-flowered corymbose patent its rachis wavy its prickles slender declining, sep. hairy felted erect-patent.—St. green, becoming quite glabrous, with a few aciculi and setæ occasionally. Lts. hard and rough beneath.—*a. Salteri*; lts. lobed or doubly serrate, rachis of cylindrical panicle nearly straight, branches corymbose patent, sep. erect-patent. A few sunken setæ on the pan. and calyx. Pet. white.— $\beta$ . *R. calvatus* (Blox.); lts. coarsely dentate the teeth distant with recurved tips interstices denticulate, rachis of lax pan. wavy branches subracemose and ascending the uppermost corymbose and patent, sep. loosely reflexed. Many sunken setæ on the panicle. Pet. pinkish or deep rose-coloured.—These plants seem to be the extremes of one species.—Open woods and hedges, rare. Sh. VII. VIII. E. I.

19. *R. carpinifolius* (W. & N.); *st. erect-arcuate angular striate* with patent clustered hairs, prickles slender conical compressed declining from a dilated compressed base, l. quinate, lts. thin irregularly but finely serrate pilose above rather paler or whitish and densely hairy beneath, term. lt. obovate-acuminate or cuspidate, *pan. narrow* racemose hairy setose its lower branches axillary short few-flowered its prickles deflexed or declining, sep. hairy setose loosely reflexed from the fruit.—*R. G.* 13.—St. forming a large lofty arch. *L. with very fine* but irregular acute *teeth* remarkably directed forwards. Pan. often nearly simple, cylindrical. Pet. white or reddish.—Open places in hilly districts. Sh. VII. VIII. E. S. I.

20. *R. villicaulis* (W. & N.); *st. arcuate* angular patentely pilose, prickles strong conical-compressed subpatent from a dilated compressed base, l. quinate, lts. thin dentate-serrate pale green and shining and often densely hairy but the hairs only on the veins beneath, term. lt. obovate or roundly cordate-obovate sub-



acuminate, *pan. open compound* leafy hairy subsetose felted its branches corymbose its prickles slender declining or deflexed, sep. hairy setose aciculate loosely reflexed from the fruit.—*R. G. 17. R. villicaulis* and *R. pampinosus* Bab.—St. becoming nearly naked, rarely a few setæ and aciculi.— $\beta$ . *R. adscitus* (Genev.); st. aciculate and setose. *R. vulgaris* (Lindl.). Heads of setæ very deciduous.—Woods and hedges. Sh. VII. VIII.  
E. S. I.

21. *R. macrophyllus* (Weihe); *st. arcuate-prostrate* angular patently pilose, prickles short slender conical-compressed declining from a large dilated compressed base, l. quinate, *lts. doubly and patently dentate* or irregularly dentate-serrate pilose above pale green felted or velvety or hairy only on the veins beneath, term. lt. elliptic or roundly obovate or obovate cuspidate or acuminate more or less cordate, *pan. hairy* felted setose its branches short few-flowered corymbose the lower axillary subracemose and ascending its prickles declining, sep. ovate-attenuate hairy felted setose loosely reflexed from the fruit.—The following forms have been considered species.—*a. R. umbrosus* (Arrh.); prickles slender from a large base, l. quinate, *lts. doubly and patently dentate* velvety or slightly felted beneath, term. lt. broadly obovate cuspidate, *pan. with slender prickles, tip of sep. linear*, cor. rose-coloured. Stam. white, scarcely exceeding the dull styles. *R. carpinifolius* (Blox.). Term. lt. sometimes divided into three.— $\beta$ . *R. macrophyllus* (W. & N.); prickles small short from a very large base, l. 5—3-nate, *lts. irregularly dentate-serrate* hairy only on the veins or rarely felted beneath, term. lt. elliptic or broadly obovate, *pan. with slender prickles, sep. leaf-pointed*, cor. white. *E. B. S. 2625. R. G. 12.* Lts. often very large. St. often with a few aciculi and setæ. Stam. pinkish, exceeding the pink styles. A very variable plant.— $\gamma$ . *R. Schlechtendalii* (W. & N.); prickles short small from a very large base, l. usually quinate, *lts. doubly and patently dentate* usually hairy only on the veins not felted beneath, *term. lt. long obovate* acuminate wedgeshaped or subcordate at the base, *pan. with strong prickles*, sep. with a linear point, cor. white. *R. G. 11.* Hardly distinguishable from var.  $\beta$ .— $\delta$ . *R. amplificatus* (Lees); prickles short from a very large base, l. usually quinate, *lts. somewhat doubly patently dentate hairy only on the veins beneath* not felted, term. lt. broadly obovate acuminate, *pan. very large* its prickles strong compressed from a very large base, sep. usually leaf-pointed, cor. white or pinkish. Remarkable for the very strong prickles with exceedingly long compressed bases on its panicle.— $\epsilon$ . *glabratus* (Bab.); prickles short from a very large base, l. quinate, *lts. irregularly or rather doubly dentate slightly hairy only on the veins beneath*, term. lt.

*roundly cordate* or broadly obovate subcordate, pan. with slender prickles, sep. (apparently) leaf-pointed. L. nearly glabrous beneath. Term. lt. very round.—Woods and thickets. Sh. VII. VIII. E. S. I.

c. *Spectabiles*. Stem with rather unequal prickles and scattered aciculi and setæ and usually dense pubescence.

22. *R. mucronulátus* (Bor.); *st.* arcuate *subterete* patently pilose, *prickles* few conical *slender* declining from an oblong dilated base, l. quinate, *lts.* thick *finely dentate-serrate* rough and pilose above hairy only on the (reddish) veins beneath, *term. lt.* broadly obovate *cuspidate* cordate below, pan. narrow lax leafy hairy setose felted its branches long 1—3-flowered its prickles small slender declining, sep. ovate-attenuate hairy setose felted with a linear point.—*Sy. E. B.* 451. *R. mucronatus* Blox. (not Ser.).—*St.* becoming nearly naked; a few setæ and aciculi. Pet. pale pink. Pink *stam.* exceeding green styles.—*R. festivus* (Warr.) seems to differ only by angular *st.* and much thinner doubly dentate leaves.—Banks and hedges. Sh. VII. VIII. E. S.

23. *R. Sprengel'ii* (Weihe); *st.* prostrate *terete* pilose, *prickles unequal deflexed* from a large compressed base, l. 3—5-nate pedate, *lts.* thin green with scattered hairs only on the veins beneath, *term. lt.* elliptic-acuminate, pan. lax hairy felted setose its axillary branches patent few-flowered the ultra-axillary divaricate its prickles few slender deflexed, sep. ovate-acuminate erect-patent often leaf-pointed.—*a.* *R. Borreri* (Bell-Salt.); *st.* arcuate-prostrate thick with scattered aciculi and setæ, prickles unequal, l. usually quinate, pan. rather thyrsoid or with a subcorymbose top, dull *stam.* falling short of styles.—*R. rubicolor* (Blox.) seems an extreme form of this.—*β.* *R. Sprengelii* (Weihe); *st.* usually quite prostrate slender, prickles small, aciculi and setæ very few, l. usually ternate, *lts.* flexible, pan. lax few-flowered pyramidally subcorymbose; rose-coloured *stam.* exceeding styles. *R. G.* 10.—Heaths and woods. Sh. VI. VII. E.

d. *Radulæ*. Stem rough with small elevated rigid points on which the deciduous setæ and aciculi were seated; prickles nearly equal.

24. *R. Bloxámii* (Lees); *st.* arcuate-prostrate angular slightly furrowed, *prickles small subpatent* unequal, aciculi and setæ short many, hairs many, l. 5- or 3-nate, *lts.* coarsely doubly *dentate green and pilose on both sides*, *term. lt.* roundly obovate *cuspidate*, pan. long leafy to the top felted its short branches and top corymbose its prickles slender declining, sep. ovate-acuminate loosely reflexed

from the fruit.—St. thick, rarely furrowed. L. subpedate. Upper floral l. simple. Pet. white. Pale Stam. falling short of styles.—Woods. Sh. VII. VIII. E.

25. *R. Hystrix* (Weihe); st. arcuate-prostrate angular slightly furrowed, prickles slender declining from a dilated compressed base, aciculi and hairs few short equalling the many setæ and much shorter than the prickles, l. quinate-pedate, *lts. coarsely and rather doubly and patently dentate green and pilose on both sides, term. lt. oblong-obovate* acuminate, pan. long leafy its branches long racemose ascending but the uppermost and ultra-axillary patent or divaricate its rachis wavy its prickles strong declining the uppermost slender, *sep. lanceolate-attenuate loosely adpressed to the fruit.*—*R. G.* 41.—*Lts.* flat, but wavy at the edges, hairy but not felted beneath. Pet. pink. Pale Stam. exceeding pink styles.—Hedges and thickets. Sh. VII. VIII. E. I.

26. *R. rosæus* (Weihe); st. arcuate-prostrate angular, *prickles* slender nearly equal declining from a dilated compressed base a few shorter slightly exceeding the nearly equal aciculi setæ and hairs, l. quinate-pedate or ternate, *lts. doubly-dentate-serrate pilose above paler and pilose only on the veins beneath, term. lt. obovate- or oblong-acuminate usually subcordate below, pan. subpyramidal truncate* leafy below its branches racemose ascending or the ultra-axillary patent corymbose or simple its rachis more or less wavy its prickles slender declining, *sep. lanceolate-attenuate loosely adpressed to the fruit.*—*R. G.* 56.—Perhaps not distinct from *R. Hystrix*. Its more elegant pyramidal and abrupt panicle and the more unequal prickles less markedly separated from the aciculi are the chief differences. “Pet. pale pink. Pale Stam. exceeding pink-based styles.”—Woods and hedges. Sh. VII. VIII. E. I.

27. *R. præruptorum* (Boul.); st. arcuate-prostrate subterete, *prickles* many slender unequal declining *slightly dilated at their base*, setæ hairs and very slender aciculi unequal many, l. quinate-pedate or ternate, *lts. coarsely unequally and doubly serrate pilose above paler and pilose only on the veins beneath, term. lt. obovate-acuminate, pan. long narrow leafy below its branches corymbose its rachis straight its prickles slender declining its aciculi setæ and hairs many and unequal, sep. ovate-attenuate aciculate with long setæ* felted loosely reflexed from the fruit.—*R. G.* 42. *R. hirtus*  $\beta$ . *Menkii* Bab. *R. pygmæus* Bab. not Weihe.—St. perhaps prostrate. Pan. rather long with distant short axillary branches, uppermost very short and often divaricate. Pet. white or pinkish. “Pale Stam. exceeding pink-based styles.”—Hedges. South-eastern counties. Sh. VII. VIII. E.

28. *R. scaber* (Weihe); st. arcuate-prostrate subangular subsulcate, prickles strong short nearly equal declining or deflexed from a long compressed base, aciculi setæ and hairs few very short, l. 3—5-nate, lts. doubly dentate opaque and pilose above pale green and pilose beneath, term. lt. broadly obovate cuspidate or acuminate subcordate below, pan. subpyramidal leafy felted truncate or blunt at the end its axillary branches erect-patent racemose the ultra-axillary racemose-corymbose or simple its prickles short declining or deflexed from a long base its aciculi strong its setæ and hairs nearly equal, sep. ovate-acuminate loosely reflexed from the fruit.—*R. G.* 32.—Stam. inflexed exceeding styles.—The typical plant is slender and elegant, pan. often nearly simple with long peduncles. The large form (*R. Babingtonii* Bell-Salt.) is often an enormous plant with very rough long prostrate stems, a very large panicle with very large and long branches. Pet. white.—Open woods. Sh. VII. VIII. E.

29. *R. rúdis* (Weihe); st. arcuate angular subsulcate, prickles strong conical compressed nearly equal subpatent exceeding the nearly equal and short aciculi setæ and hairs, l. quinate, lts. coarsely and doubly serrate (or lobate-serrate) greenish-white-felted beneath, term. lt. elliptic or broadly oblong-obovate acuminate, pan. long leafy its branches ascending corymbose-racemose the uppermost and ultra-axillary divaricate its rachis straight its prickles strong declining or deflexed from a long base the uppermost slender, sep. ovate-attenuate strongly reflexed.—*R. G.* 40.—The nearly equal not scattered prickles, short aciculi setæ and hairs, jagged lts. felted beneath, and strongly reflexed sepals are marks of this species. Pet. white. Stam. exceeding styles.—Hedges and thickets. Sh. VII. VIII. E. S. I.

30. *R. Rádula* (Weihe); st. arcuate angular, prickles slender declining from a dilated compressed base exceeding the many short unequal aciculi setæ and hairs, l. quinate-pedate, lts. finely but doubly and patently dentate greenish-white-felted beneath, term. lt. obovate acuminate or subcuspidate, pan. long leafy its branches short corymbose ascending its prickles strong declining from a long base the uppermost slender, sep. ovate loosely reflexed from the fruit.—*a. R. Radula* (Weihe); prickles on the st. unequal, term. lt. obovate acuminate. Connivent stam. much exceeding styles. *E. B. S.* 452. *R. G.* 39.—*β. R. Leightonii* (Lees); prickles on the st. nearly equal, term. lt. obovate cuspidate.—*γ. denticulatus* (Bab.); term. lt. broadly quadrangular-obovate cuspidate subcordate below broadly but faintly dentate the teeth denticulate.—Hedges. Sh. VII. VIII. E. S. I.

(4) *Glandulosi*. Stems arcuate-prostrate or prostrate, rooting, hairy. Prickles abundant, very unequal, scattered, passing gradually into abundant aciculi and setæ.

a. *Koehleriani*. Leaves quinate or rarely ternate. Prickles and setæ thickened at their base.

31. *R. Koehleri* (Weihe); st. arcuate-prostrate roundish or angular pilose, prickles very unequal slightly declining from a compressed base, aciculi and setæ very unequal, *lts.* unequally or rather doubly dentate *even above* pale green *hairy on the veins beneath*, term. lt. cordate-ovate, *basal lts. not imbricate* stalked, pan. open leafy its branches short patent corymbose or the axillary branches racemose its prickles many long slender declining its aciculi setæ and hairs many unequal, *sep.* ovate-attenuate *patent or reflexed from the fruit*. Pale *stam.* exceeding pink styles.—a. *R. Koehleri* (Weihe); st. with many prickles aciculi and setæ, *lts. hairy only on the veins and rough beneath*, pan. open truncate often broad-topped its branches long corymbose patent or the axillary branches racemose and ascending, *term. ped. of pan. and branches shorter than the lateral ones.* *R. G.* 25. *E. B. S.* 2605.—β. *infestus* (Bab.); st. with many strong prickles aciculi and setæ, *lts. hairy only on the veins and soft beneath*, pan. broad but compact rounded at the end its branches moderate rather corymbose erect-patent, *term. ped. of pan. and branches shorter than the lateral ones*, prickles of pan. strong deflexed. *R. obliquus* (Blox.) perhaps belongs here.—γ. *R. pallidus* (Weihe); st. with fewer but strong prickles aciculi and setæ, *lts. often slightly felted hairy on the veins and soft beneath*, pan. narrow its branches short corymbose-racemose patent or the axillary branches ascending, *term. ped. of branches often longer than the lateral ones.* *R. G.* 29.—δ. *R. cavatifolius* (Müll.); st. subglabrous, aciculi and setæ unequal inconspicuous, hairs very few, *lts. not felted* but hairy on the veins beneath, *term. lt. cordate cuspidate-attenuate*, pan. abrupt with very short terminal peduncles.—All are usually very prickly; their very strong prickles pass very gradually into aciculi, and those into setæ; st. often quite covered by their broad bases. Pet. white.—Hedges and thickets. Sh. VII. VIII. E. S. I.

32. *R. fusco-ater* (Weihe); st. arcuate-prostrate angular hairy, prickles unequal slightly declining from a very large compressed base, setæ and strong unequal aciculi many, *lts.* irregularly or rather doubly dentate *even above* green and hairy beneath, term. lt. broadly cordate-ovate acuminate or subcuspidate, *basal lts. stalked imbricate*, pan. long subpyramidal leafy below its branches patent corymbose or the axillary branches erect-patent racemose

its prickles many unequal longest at about the middle of the fl.-shoot its hairs setæ and aciculi many unequal, *sep.* ovate-attenuate setose aciculate *patent or adpressed to the fruit.*—*R. G.* 26.—The lts. are all imbricate and rather rough beneath. Pet. pink. Stam. incurved exceeding styles.—Heaths. Sh. VII. VIII. E.

33. *R. emersisty'lus* (Müll.); st. arcuate-prostrate angular hairy, prickles slightly declining unequal from a compressed base a few smaller, *setæ aciculi and hairs very short nearly equal, lts.* rather irregularly and rather doubly dentate *thick* even above green and hairy on the veins beneath, *term. lt. orbicular-ovate attenuate with a cordate base,* basal lts. very shortly stalked imbricate, pan. often leafy nearly to the top its *axillary branches corymbose* its prickles small unequal declining its setæ and hairs many unequal, *sep.* ovate-attenuate with a rather leaflike point setose aciculate *patent or adpressed to the fruit.*—*R. Briggsii* (Blox.) *J. of B.* vii. t. 88.—Stam. exceeding styles.—Heaths in the South. Sh. VII. VIII. E.

34. *R. diversifolius* (Lindl.); st. arcuate prostrate angular *sparingly pilose,* prickles unequal subpatent from a compressed base, aciculi and setæ many unequal, *lts.* often irregularly or towards their tip rather doubly dentate wavy at the edge *rugose* above pale green hairy and felted beneath, *term. lt. broadly cordate obovate-acuminate basal lts. subsessile imbricate, pan. long leafy nearly to the top* its branches erect-patent subracemose its prickles longest at about the middle of the fl.-shoot its hairs setæ and aciculi short equal, *sep. ovate acute* felted setose *patent or loosely adpressed to the fruit.*—*R. fusco-ater* Bab. formerly. *J. of B.* viii. t. 107.—A very prickly plant which differs much in appearance from *R. fusco-ater.* Lts. usually soft beneath. Pet. white. Stam. much exceeding styles.—Hedges. Sh. VII.\* VIII. E.

35. *R. mutabilis* (Genev.); stem arcuate-prostrate angular sparingly pilose and setose, prickles moderate unequal subpatent from a long compressed base, aciculi strong very unequal mostly short, l. quinate, lts. doubly or mostly lobate-dentate pilose above hairy or felted *beneath, term. lt. obovate acuminate, basal lts. stalked not imbricate, pan. long narrow pyramidal leafy* nearly to the top its *branches and branchlets subcorymbose subpatent its rachis very prickly* aciculate and setose its prickles many very strong from large compressed bases, *sep.* ovate felted setose reflexed with rather leaflike points.—Pet. dull white. “Filam. pink. Styles greenish.”—Hedgebanks in the South. Sh. VII. VIII. E.

36. *R. Lejeun'ii* (Weihe); st. arcuate-prostrate subangular with scattered hairs and setæ, prickles mostly small a few longer declining from a long compressed base, *aciculi very short*, l. quinate-pedate, lts. opaque and pilose above paler and hairy only on the veins beneath lobate-serrate towards the tip, *basal lts. stalked* not imbricate, term. lt. obovate-acuminate, pan. broad hairy leafy setose its top corymbose its axillary *branches ascending* subracemose its prickles slender declining its setæ many unequal, *sep. ovate* felted setose loosely adpressed to the fruit.—*R. G.* 31.—Sep. often having a short linear point; its allies when furnished with an appendage to the sepals have it leaf-like.—Hedges. Sh. VII. VIII. E.

b. *Bellardiani*. Leaves ternate or quinate-pedate; basal leaflets not imbricate, stalked. Prickles mostly confined to the angles of the very hairy aciculate and setose stems.

37. *R. pyramidalis* (Bab. not Kaltenb.); st. very nearly prostrate roundly angular, *prickles many short strong* much declining or deflexed from a very large compressed base, hairs few, aciculi and setæ many nearly equal, l. ternate or rarely quinate-pedate, *lts. convex* irregularly dentate-serrate nearly equal opaque and pilose above paler and *pilose beneath*, term. lt. obovate-cuspidate, *pan. pyramidal* leafy below its top and branches racemose felted its *rachis straight rigid* its prickles slender declining its hairs and setæ many unequal, *sep. lanceolate-attenuate* loosely adpressed to the fruit.—St. usually quite prostrate. Pan. very pyramidal. Pet. very narrow, greenish white. Stam. white exceeding pinkish styles.—Edges of woods, rare. Abundant at Llanberis, N. Wales. Sh. VII. VIII. E.

38. *R. Gun'theri* (Weihe); st. arcuate-prostrate terete, *prickles slender unequal* declining from a large subcompressed base, aciculi setæ and hairs short nearly equal, l. ternate or rarely quinate-pedate, *lts. flat* irregularly or doubly dentate-serrate nearly equal opaque and pilose above pilose greenish white or slightly felted beneath, term. lt. obovate-acuminate, pan. narrow leafy its branches distant ascending racemose few-flowered its *rachis wavy* its prickles few slender declining its hairs and setæ many equal, *sep. ovate-lanceolate* loosely reflexed from the fruit.—*R. G.* 21. *R. saltrum* Focke.—Only slightly prickly. Pan. very loose, its rachis forming an angle at each leaf. Pet. narrow, pale pink. Stam. equalling styles.—Shady places. Sh. VII. VIII. E. I

39. *R. humifusus* (Weihe); st. arcuate-prostrate nearly round, prickles many slender very unequal declining from a long compressed base, *hairs setæ and very slender aciculi unequal* many

patent, l. quinate-pedate, *lts. finely but doubly patently dentate* pilose and opaque above pilose greenish white and shining beneath, term. lt. obovate-oblong subcuspidate, pan. broad leafy below its branches corymbose its rachis rather wavy its prickles few slender declining its hairs and setæ many unequal, sep. ovate slightly aciculate shortly setose loosely reflexed from the fruit.—*R. G.* 35. *R. hirtus* a. Bab.—Pan. sometimes nearly simple. Pet. white, large. Stam. exceeding styles. Sometimes the underside of the lts. is felted.—Woods and thickets. Sh. VII. VIII. E. S. I.

40. *R. foliosus* (Weihe); st. arcuate-prostrate angular, prickles many slender unequal declining from a long compressed base, setæ and very slender aciculi scattered unequal, hairs few, l. quinate-pedate, *lts. unequally dentate-serrate* pilose and opaque above *paler and pilose beneath*, term. lt. roundly cordate acuminate, pan. long narrow leafy to the top its branches short erect-patent corymbose its rachis rather wavy its prickles very slender many declining its hairs and setæ many unequal, *sep. ovate-attenuate aciculate setose* hairy laxly reflexed from the fruit.—*R. exsecatus* Müll.—β. *R. adornatus* (Müll.), has term. lts. obovate-acuminate or even narrowed below.—Hartshill Wood, Warw. Plymouth. Sh. VII. VIII. E.

41. *R. glandulosus* (Bell.); st. arcuate-prostrate nearly round, prickles small declining from a long compressed base, aciculi setæ and hairs many nearly equal, l. ternate or rarely quinate, lts. nearly equal oblong cuspidate hairy only on the veins beneath, term. lt. subcordate-ovate-acuminate, pan. felted aciculate very setose its branches erect-patent axillary its top racemose its prickles slender declining, sep. ovate-attenuate aciculate setose felted loosely adpressed to the fruit or patent.—*a. R. Bellardi* (Weihe); l. ternate, lts. nearly equal oblong doubly dentate-serrate shortly pilose on the veins beneath, lateral lts. divaricate, pan. with axillary distant corymbose branches its rachis usually wavy, *stam. slightly exceeding styles.* *R. G.* 44. *R. glandulosus* *E. B. S.* 2883. The divaricate lts. are remarkable, and the very open panicle. Sometimes the l. are quinate. A subvariety, *R. dentatus* Blox., has a slightly angular stem, lts. ovate-acuminate-cuspidate with a cordate base ashy green beneath, lateral lts. patent or ascending.—β. *R. hirtus* (W. & K.); l. quinate coarsely and unequally serrate with long and dense hairs on the veins and shining beneath, term. lt. subcordate-ovate-acuminate, pan. often long its branches racemose or corymbose shortly setose its rachis nearly straight. *R. G.* 43. *R. fuscus* Lees. A subvariety, *R. rotundifolius* Blox., *R. deflexidens* Boul.?, has a slightly angular stem, l. ternate or rarely quinate,



lts. doubly dentate cuspidate, term. lt. nearly round with a subcordate base.— $\gamma$ . *R. Reuteri* (Merc.); l. quinate coarsely and rather doubly dentate-serrate with few hairs on the veins beneath, term. lt. obovate-rhomboidal acuminate, pan. truncate its branches short subcorymbose few-flowered upper 1—3-flowered very aciculate setose and hairy, rachis nearly straight. Much more hairy and setose than the other forms and with some stronger declining or deflexed prickles on the stem.—Woods. Sh. VII. VIII. E. I.

- (5) *Cæsi*. Stems most often arcuate-prostrate, terete or slightly angular, usually with a glaucous bloom; aciculi and setæ few or none; prickles unequal.

42. *R. Balfourianus* (Blox.); st. arcuate-prostrate nearly round patently pilose, aciculi and setæ few, prickles slender unequal scattered patent from an oblong subcompressed base, l. quinate, lts. dentate-serrate green on both sides rugose and pilose above hairy (not felted) beneath, term. lt. cordate or ovate acute, basal lts. sessile imbricate, pan. loose leafy hairy rather setose its branches long distant few-flowered racemose-corymbose erect-patent, *sep. ovate-acuminate erect-patent*, styles pale flesh-coloured, fr. oblong its torus oblong stalked.—A very variable plant, approaching *R. corylifolius* in some states. Pet. pale pink. Stam. exceeding styles.—Hedges. Sh. VII. VIII. E. I.

43. *R. corylifolius* (Sm.); st. arcuate-prostrate nearly round or obtuse-angled nearly glabrous, aciculi setæ and sessile glands very few, prickles subulate slender nearly equal subpatent or rarely deflexed from a long base, l. quinate, lts. doubly serrate green on both sides rugose with scattered hairs above paler and felted beneath, term. lt. roundly cordate or ovate cuspidate or acuminate, basal lts. sessile imbricate, pan. and its branches subcorymbose, *sep. ovate cuspidate reflexed from the fruit*, pet. roundly ovate, styles greenish, torus oblong stalked.—St. slightly glaucous. Pan. usually with 2 or 3 long axillary branches, felted, more or less prominently setose at the top.— $\alpha$ . *R. sublustris* (Lees); st. nearly round reddish green, prickles slender subpatent from a long base, lts. ashy-felted beneath, term. lt. often subtrilobed roundish cordate, rachis straightish with few prickles. *R. corylifolius* Sm. *Sy. E. B.* 455. St. thick. Pet. white. Stam. slightly exceeding styles.— $\beta$ . *conjungens* (Bab.); st. rather angular, reddish green, prickles slender strong subpatent or deflexed from a compressed very long base often with slightly deflexed points, lts. ashy-felted beneath, term. lt. cordate-ovate or broadly obovate with a subcordate base, rachis straightish

with few prickles. *R. Wahlbergii* Bell-Salt. St. round at the base, angular but flat-sided above. Pet. white or pink.—*γ. purpureus* (Bab.); st. angular purple often a little floccose, prickles strong subpatent or deflexed from a long compressed base, lts. pale-green- or white-felted beneath, term. lt. roundly- or subcordate-obovate, rachis rather wavy with many prickles. *R. Wahlbergii* Arrh. St. round at the base, angular and often furrowed above. Pet. white or pink.—Hedges and thickets. Sh. VI.—VIII. E. S. I.

44. *R. althæifolius* (Host); st. prostrate slightly angular with scattered hairs and setæ, prickles many unequal slender patent from an oblong compressed base, l. quinate or ternate, lts. crenately lobed pale green with hairs on the veins loosely white-felted beneath, basal lts. of the ternate l. retrorsely bipartite of the quinate l. sessile imbricate, term. lt. rhomboidal-obovate subcordate below, pan. leafy its axillary branches and top racemose-corymbose with few very short setæ, prickles on the middle of the rachis longest slender, sep. ovate-subacuminate setose loosely adpressed to the (black-blue) fruit, pet. obovate, styles flesh-coloured at their base. *R. deltoideus* Müll. ?—St. round at the base, above bluntly angular or even furrowed. Pan. rather long, open; top formed of irregular corymbs of nearly simple peduncles. Pet. nearly white. Pale stam. about equalling pink styles.—Hedges. Sh. VI.—VIII. E.

45. *R. tuberculatus* (Bab.); st. arcuate-prostrate slightly angular with scattered short hairs and setæ, prickles many unequal slender patent from an oblong tuberculiform base, l. ternate or quinate, lts. rather doubly dentate hairy on the veins beneath green on both sides, basal lts. of the ternate l. bilobate of the quinate l. nearly sessile imbricate, term. lt. roundly cordate subcuspidate, pan. leafy its axillary branches racemose its top corymbose, prickles from the middle to the top of the pan. and ped. slender and longest, sep. ovate acuminate aciculate setose loosely adpressed to the fruit.—*J. of B.* viii. t. 106. *R. nemorosus* *δ. ferox* Leight. Probably *R. teretiusculus* (Kalt.!).—St. very bluntly angular. Pan. short, its branches few-flowered. Pet. pinkish.—Mr. Warren combines my *R. diversifolius* and *R. Briggsii* with this under the name of *R. dumetorum*. His *R. concinnus* is near *R. corylifolius*, but has not felted l. on my authentic specimen. It may be distinct.—Hedges. Sh. VII. VIII. E. I.

46. *R. cæsius* (L.); st. prostrate terete glaucous, prickles small unequal subulate declining or deflexed from a long compressed base, l. ternate or rarely pinnate, lts. unequally cut or coarsely serrate, term. lt. ovate rhomboidal-ovate or 3-lobed,

lateral lts. rather bilobed subsessile, pan. nearly simple often very small, sep. ovate acuminate with a long linear point adpressed to the glaucous fruit, pet. obovate notched, styles greenish.—*a. agrestis*; st. very slender, prickles few small, lts. flat lobate-serrate rather pilose on both sides, term. lt. rhomboidal-ovate acuminate rounded below. *R. cæsius a. aquaticus* and *β. agrestis* R. G. 46. A. Pan. often nearly simple, or its branches rarely more than once divided but often very long.—*β. R. tenuis* (Bell-Salt.); st. very slender, prickles many strong but small nearly equal deflexed, lts. flat (?) doubly serrate pilose on both sides or villose beneath, term. lt. obovate or cordate-obovate acuminate, fr. black “not glaucous.” St. sometimes having a very few aciculi and setæ. Bases of prickles much enlarged.—*γ. ulmifolius*; st. slender purplish, prickles many small deflexed or declining, aciculi setæ and hairs few short, lts. rather rugose lobate-serrate pilose only on the veins or hairy or slightly ashy-felted beneath, term. lt. roundly cordate acuminate often 3-lobed or rarely divided into 3 sessile lts. of which the term. is narrowed below. *R. cæsius β. pseudo-cæsius* R. G. 46. B. f. 1. Often much larger than var. *a* and *β*. St. thicker. L. very broad.—*δ. intermedius*; st. thicker greenish purple, prickles many slender very unequal subpatent, aciculi and setæ few strong very short, lts. lobate-serrate pilose on the veins beneath, term. lt. triangular-cordate acuminate 3-lobed or 3-partite or divided into 3 sessile lts. of which the term. is narrowed to the base.—*ε. R. pseudo-Idæus* (Lej.); st. thick green slightly glaucous, prickles slender violet-coloured subpatent, aciculi and setæ few very short, l. ternate or quinate-pinnate, lts. doubly-serrate ashy-felted beneath, lateral lts. all sessile, term. lt. stalked roundly-cordate. *R. G. 46. B. f. 2.* I have only seen one specimen of this var., found at Hunsdon, Herts.—*ζ. serpens* (Godr.?) ; st. slender green, l. ternate, lts. lobate-serrate pilose on the veins beneath, lateral lts. with a large backward lobe, term. lt. obovate acuminate subcordate below, ped. and sep. very setose felted scarcely hairy. *R. G. 46. C. f. 1.*—Hedges and thickets, common. Sh. VI. VII. Dewberry. E. S. I.

## B. HERBACEI.

Stem nearly or quite herbaceous. Leaves ternate or simple. Stipules usually attached to the stem. Fl. umbellate or nearly solitary. Receptacle flat.

i. *Saxatiles*. Stems slender, prostrate. Flowers umbellate or nearly solitary. Succulent carpels large, few, distinct.

47. *R. saxat'ilis* (L.); st. rooting annual, prickles none or very few minute weak, l. ternate, fl.-shoot erect with a *terminal few-flowered corymb*, pet. lanceolate about equalling the calyx.—*E. B.* 2233. *R. G.* 9.—St. very slender. Fl.-shoots radical, about a span high. Fl. white. Fr. of 1—4 large drupes.—Stony mountains or high hills. P. VII. VIII. E. S. I.

ii. *Arctici*. No sterile stems but a long subterranean rhizome. Flowers terminal, nearly solitary. Succulent carpels adhering together.

48. *R. Chamæmorus* (L.); fl.-shoot erect unarmed 1-flowered herbaceous, fl. diceious, l. simple lobed plicate.—*E. B.* 716. *R. G.* 49.—Rhizome woody. Fl. large, white. Fr. large, red, afterwards orange-yellow.—Alpine turf-bogs. P. VI. *Cloud-berry*. E. S. I.

[*R. arcticus* (L.); st. erect unarmed nearly 1-flowered herbaceous, l. ternate, pet. obovate twice as long as the calyx, stam. connivent, succulent carpels many.—*E. B.* 1585. *R. G.* 48.—Rhizome subterranean. Fl.-shoot 6—10 in. high. Lts. nearly equal. Fl. rose-coloured.—Isle of Mull and Ben y Gloe. Probably a mistake. P. VI.] S.

### Index to the Rubi.

- |                            |                                |                           |
|----------------------------|--------------------------------|---------------------------|
| <i>R. adornatus</i> , 40.  | <i>cordifolius</i> , 8.        | Guntheri, 38.             |
| <i>adscitus</i> , 20.      | <i>corylifolius</i> , 43.      | hemistemon, 6.            |
| <i>affinis</i> , 5.        | <i>deflexidens</i> , 41.       | hirtus, 41.               |
| <i>agrestis</i> , 46.      | <i>deltoides</i> , 44.         | hirtifolius, 15.          |
| <i>althæifolius</i> , 44.  | <i>dentatus</i> , 41.          | <i>hirtus</i> , 39, 41.   |
| <i>arcticus</i> , 48.      | <i>denticulatus</i> , 30.      | <i>hispidus</i> , 46.     |
| <i>amplificatus</i> , 21.  | <i>derasus</i> , 19.           | <i>humifusus</i> , 39. *  |
| <i>Babingtonii</i> , 28.   | <i>discolor</i> , 12.          | Hystrix, 25.              |
| <i>Balfourianus</i> , 42.  | <i>diversifolius</i> , 34.     | Idæus, 1.                 |
| <i>Bellardi</i> , 41.      | <i>dumetorum</i> , 45.         | <i>imbricatus</i> , 10.   |
| <i>Bloxamii</i> , 24.      | <i>emersistylus</i> , 33.      | <i>incurvatus</i> , 9.    |
| <i>Borreri</i> , 23.       | <i>exsecatus</i> , 40.         | <i>infestus</i> , 31.     |
| <i>Briggsii</i> , 33.      | <i>festivus</i> , 22.          | <i>intermedius</i> , 43.  |
| <i>cæsius</i> , 46.        | <i>fissus</i> , 4.             | Koehleri, 31.             |
| <i>calvatus</i> , 18.      | <i>foliosus</i> , 40.          | <i>latifolius</i> , 11.   |
| <i>carpinifolius</i> , 19. | <i>fruticosus</i> , 4, 12, 13. | <i>Leesii</i> , 1.        |
| <i>carpinifolius</i> , 21. | <i>fusco-ater</i> , 32.        | <i>Leightonii</i> , 30.   |
| <i>cavatifolius</i> , 31.  | <i>fusco-ater</i> , 17, 34.    | <i>Lejeunii</i> , 36.     |
| <i>Chamæmorus</i> , 48.    | <i>fuscus</i> , 41.            | <i>lentiginosus</i> , 5.  |
| <i>Colemanni</i> , 17.     | <i>glabratus</i> , 21.         | <i>leucostachys</i> , 14. |
| <i>concinus</i> , 45.      | <i>glandulosus</i> , 41.       | <i>Lindleianus</i> , 7.   |
| <i>conjungens</i> , 43.    | <i>Grabowskii</i> , 16.        | <i>macrophyllus</i> , 21. |

- |                            |                            |                              |
|----------------------------|----------------------------|------------------------------|
| <i>Menkii</i> , 27         | <i>pyramidalis</i> , 37.   | <i>Schlechtendalii</i> , 21. |
| <i>mucronatus</i> , 22.    | <i>pyramidalis</i> , 40.   | <i>serpens</i> , 46.         |
| <i>mucronulatus</i> , 22.  | <i>Radula</i> , 30.        | <i>Sprengelii</i> , 23.      |
| <i>mutabilis</i> , 35.     | <i>ramosus</i> , 10.       | <i>suberectus</i> , 3.       |
| <i>nemorosus</i> , 45.     | <i>Reuteri</i> , 41.       | <i>sublustris</i> , 43.      |
| <i>nitidus</i> , 7.        | <i>rhamnifolius</i> , 8.   | <i>tenuis</i> , 46.          |
| <i>obliquus</i> , 31.      | <i>rosaceus</i> , 26.      | <i>teretiuseculus</i> , 45.  |
| <i>pallidus</i> , 31.      | <i>rotundifolius</i> , 41. | <i>thyrsoides</i> , 13.      |
| <i>pampinosus</i> , 20.    | <i>rotundifolius</i> , 41. | <i>tuberculatus</i> , 45.    |
| <i>plicatus</i> , 5.       | <i>rubicolor</i> , 23.     | <i>ulmifolius</i> , 46.      |
| <i>præruptorum</i> , 27.   | <i>rudis</i> , 29.         | <i>umbrosus</i> , 21.        |
| <i>pseudo-cæsius</i> , 46. | <i>Salteri</i> , 18.       | <i>vestitus</i> , 14.        |
| <i>pseudo-Idæus</i> , 46.  | <i>saltuum</i> , 38.       | <i>villicaulis</i> , 20.     |
| <i>purpureus</i> , 43.     | <i>saxatilis</i> , 47.     | <i>vulgaris</i> , 20.        |
| <i>pygmæus</i> , 27.       | <i>scaber</i> , 28.        | <i>Wahlbergii</i> , 43.      |

## 12. DRY'AS Linn.

1. *D. octopet'ala* (L.); l. crenate-serrate blunt, sep. 3 or 4 times as long as broad more or less pointed, base of the cal. hemispherical.—*E. B.* 451. *St.* 20. 3.—Fl. large, whiter. Pet. 8. L. simple, white with fine dense woolly pubescence beneath. *St.* prostrate, woody. Seeds obovate-oblong apiculate.—*a.*; sep. acute covered with red hairs, l. ovate-oblong, deeply cut into large rounded lobes, underside of the petiole and midrib green bearing minute linear pellucid fringed scales mixed with long hairs, midrib and lateral ribs sparingly hairy and conspicuous.—*β. pilosa* (Bab.); sep. slightly pointed covered with nearly black hairs, l. oblong or ovate-oblong deeply cut into large slightly acute lobes, underside of the petioles and midrib dark red hairy but without scales, midrib and lateral ribs covered with long white hairs and inconspicuous, awn longer and more feathery.—*D. depressa* (Bab.), first separated by the late Dr. J. T. Mackay, has no constant character.—Alpine situations, particularly on limestone. P. VI. VII. E. S. I.

## 13. GE'UM Linn. Avens.

1. *G. urbánum* (L.); fl. erect, *pet.* obovate, cal. of the fruit reflexed, *carpophore* 0, lower joint of the style much longer than the glabrous upper joint, radical l. interruptedly pinnate and lyrate, stem-l. ternate, stip. large rounded lobed and cut.—*E. B.* 1400. *St.* 5. 7.—*St.* 2 feet high. Fl. small, bright yellow, calyx green. Upper joint of the style with a few minute hairs at its base.—Hedges and thickets. P. VI.—VIII. *Wood-Avens.*  
E. S. I.

[*G. intermédiu*m (Ehrh.); fl. erect or nodding, *pet.* roundish with a wedgeshaped claw, cal. of the fruit patent, *carpophore* 0 or

short, lower joint of the style longer than the hairy upper joint, radical l. interruptedly pinnate and lyrate, stem-l. 3-lobed, stipules round toothed.—*Sy. E. B.* 458.—St. 1—2 feet high. Fl. larger than in Sp. 1, less than in Sp. 2, yellow, calyx purplish. Upper joint of style clothed with long hairs but with a rather long glabrous point. Perhaps a hybrid; or rather it consists of extreme forms of Sp. 1 and 2.—Damp woods. P. VI. VII.] E. S.

2. *G. rivale* (L.); fl. nodding, pet. broadly obovate emarginate or orbiculate with a long wedged claw, cal. of the fruit erect, carpophore long, lower joint of the style equalling the long hairy upper joint, radical l. interruptedly pinnate and lyrate, stem-l. ternate, stip. small ovate toothed.—*E. B.* 106. St. 3.—St. about 1 foot high. Fl. large, purplish brown with darker veins, calyx purplish. Carpophore nearly equalling the calyx. Upper joint of the style with a short glabrous point.—Damp woods. P. VI. VII. *Water-Avens.* E. S. I.

#### Tribe IV. *Rosidæ*.

##### 14. *RO'SA* Linn.<sup>1</sup> Rose.

i. SPINOSISSIMÆ. Styles free, scarcely protruding. Sep. mostly persistent. St. short, erect, with many slender unequal prickles lessening gradually into aciculi and setæ.

1. *R. spinosis'sima* (L.); prickles many crowded very unequal mostly straight subulate smallest undistinguishable from aciculi, lts. simply serrate glabrous and glandless on both sides, sep. simple acuminate not glandular on the back, fr. nearly globular erect with no disk.—*E. B.* 187.—St. erect with short compact branches. Fl. solitary, white. Fr. dark purple or black, ripe in September.—Sandy heaths and sea-shore. Sh. V. E. S. I.

2. *R. hiber'nica* (Sm.); prickles scattered unequal gradually passing into aciculi, the larger strong slightly falcate, lts. simply serrate naked or hairy on the veins beneath without glands, sep. pinnate not glandular on the back, fr. roundish with a small disk.—*E. B.* 2196.—St. erect; branches mostly short. Fl. 1—3 rarely more together, pale pink; ped. and cal.-tube quite naked. Fr. rather urceolate, blood-red, with subsistent sepals, ripe in October.—Chiefly in the north. Sh. V.—VIII. E. S. I.

<sup>1</sup> For fuller details concerning the varieties and states of Roses see Baker in *Linn. Journ.* xi. 197. See also *Bull. Soc. Belg.* xv. 202 &c. *Seta*, see note, p. 106. We want a thorough monograph of this genus.

3. *R. involuta* (Sm.); prickles many crowded gradually passing into abundant aciculi, *lts. doubly serrate downy glandular beneath*, sep. simple or pinnate glandular on the back, fr. subrotund with no disk.—*R. Sabini* Woods. *E. B. S.* 2594.—St. erect with short branches. Fl. 1—3; ped. aciculate and setose. Fr. red or pink, with persistent sepals, ripe in Sept. and Oct.—Typical *R. involuta* (*E. B.* 2068) is a smaller plant with l. scarcely at all glandular and very slightly compound serratures, and solitary flowers. *R. Donniana* (*E. B. S.* 2601) has very hairy leaves and nearly simple sepals. *R. Wilsoni* (*E. B. S.* 2723) has simply serrate *lts. hairy on both sides without glands*, sep. nearly simple, fr. ovate-urceolate.—Chiefly in the North. Sh. VI. E. S. I.

ii. VILLOSÆ. Sepals mostly persistent, densely glandular on the back. St. tall, suberect, with uniform nearly straight scattered prickles. L. very hairy and inconspicuously glandular beneath.

[*R. pomifera* (Herrm.); prickles uniform slender straight, *lts. oblong-lanceolate doubly serrate downy on both sides*, sep. quite persistent copiously pinnate, fr. globose with no disk.—St. erect with arching branches. Fl. 1—3; ped. about  $\frac{1}{2}$  an inch long. Fr. bright red with *connivent sepals*, ripe in August.—Not a real native. Sh. VI. ?]

4. *R. mollis* (Sm.); prickles uniform slender nearly straight, *lts. ovate doubly serrate softly downy, especially beneath*, sep. quite persistent slightly pinnate, fr. subglobose with no disk.—*E. B.* 2459.—Fl. 1—3, deep rose-colour; ped. very short. Fr. bright red with *connivent sepals*, ripe in Aug. or July.—Northern counties. Sh. VI. VII. E. S. I.

5. *R. tomentosa* (Sm.); prickles mostly uniform slender straight or slightly curved, *lts. elliptic or subovate doubly serrate downy especially beneath*, sep. subpersistent copiously pinnate, fr. oblong urceolate with a slight disk.—*E. B.* 990.—St. erect with arching branches. Fl. 1—3, pink or white; ped. rather long. Fr. bright red with spreading (or subdeciduous) sepals, ripe in Sept. *R. Sherrardi* Dav., *R. subglobosa* Sm., and *R. farinosa* (Rau) have more globose fruit and, as well as other forms, have been noticed in the north.—Hedges and thickets northwards. Sh. VI. VII. E. S. I.

iii. RUBIGINOSÆ. Sepals subpersistent. St. suberect, with scattered hooked nearly uniform prickles becoming more slender upwards. L. scarcely hairy but very glandular throughout beneath. Fr. with a small disk.

6. *R. rubiginosa* (L.); prickles many the larger hooked the

*smaller subulate unequal* straight, lts. doubly serrate downy beneath, sep. persistent copiously pinnate, primordial fr. pear-shaped, disk very small.—*E. B.* 991.—St. erect, with short straight prickles. Fl. 1—4, rose-coloured; ped. very setose; styles hairy. Fr. subglobose (except the first), ripe in October. Foliage scented.—Bushy places. Sh. V.—VII. E. S.

7. *R. micrantha* (Sm.); prickles uniform hooked, lts. doubly serrate downy and glandular beneath, sep. subpersistent with long leaf-like points and minute lanceolate simple pinnæ, primordial fr. rounded at the base, disk moderate.—*E. B.* 2490.—St. suberect, with long lax branches. Fl. few, pale; ped. very setose; styles glabrous. Fr. ovate-urceolate, bright scarlet, ripe in Sept. or Oct. Foliage scarcely scented, lts. rounded at the base.—Thickets. Not common. Sh. VII. VIII. E. I.

8. *R. inodora* (Fr.); prickles rather unequal hooked, a few aciculi, lts. doubly serrate downy and glandular beneath, sep. subpersistent closely pinnate, primordial fr. ovate or subglobular, disk moderate.—*R. sepium* Borr., *E. B. S.* 2653.—St. suberect, with long lax branches. Fl. 1—3, pink; ped. usually naked; styles hairy. Foliage scarcely scented.—Local. Sh. VI. E.

iv. CANINÆ. Sep. usually naked at the back. St. long, arching, with scattered hooked uniform prickles becoming more slender upward. L. without glands on either surface.

9. *R. canina* (L.); prickles equal hooked, lts. simply or doubly serrate glabrous on both sides or hairy beneath, sep. reflexed deciduous pinnate, fr. ovate-urceolate or roundish, disk flat.—St. erect, with long arching branches. L. flat or keeled, rarely a few glands on the veins. Fl. few; rarely the sep. are subpersistent and ascending; ped. mostly naked; styles distinct, hairy.—Hedges and thickets. Sh. VI. VII. E. S. I.

The following forms may be distinguished:—a. *R. canina* (L.); lts. without glands beneath, sep. reflexed after fall of corolla.—\* *L. glabrous on both sides; ped. without bristles.*—*R. canina* (Woods); lts. elliptic flat simply serrate green or grey, fr. urceolate. *E. B.* 992.—*R. surculosa* (Woods); lts. rounded flat simply serrate green or grey.—*R. sphaerica* (Gren.); lts. rounded below (flat?) simply serrate, bright green, fr. globose.—*R. sarmentacea* (Woods); lts. roundish or elliptic keeled doubly serrate. *E. B. S.* 2595. *R. dumalis* and *R. biserrata* Baker.—\*\* *L. naked above, hairy on the veins beneath, ped. without bristles.*—*R. Forsteri* (Sm.); lts. rounded below usually simply but irregularly serrate (*E. B. S.* 2611) or in *R. arvatica* (Baker) very compound.—\*\*\* *L. hairy on both sides, ped. without bristles.*—*R. dumetorum* (Woods); lts. much rounded below simply ser-



rate. *E. B. S.* 2610.—*R. cæsia* (Borr.); lts. narrowed below doubly serrate. *E. B.* 2367.—*R. tomentella* (Lem.); lts. rounded below doubly serrate very hairy beneath, fr. subglobose.—Forms of most of the above with more or less prickly and setose peduncles have been named but scarcely deserve notice.—β. *R. monticola* (Rap.); lts. without glands beneath, sep. ascending after fall of corolla subpersistent.—*R. coriifolia* (Fr.); lts. usually simply (*R. bractescens* Woods), sometimes (*R. Watsoni*, Bak.) doubly serrate.—Other slightly varying forms are named.—γ. *R. Borreri* (Woods); lts. glandular only on the veins beneath, the surface naked. *R. dumetorum* *E. B.* 2579, *R. Bakeri* (Deseg.) *Sy. E. B.* 473, and *R. marginata* (Wallr.) seem hardly to differ from *R. Borreri*, which itself is not easily distinguishable from *R. tomentella*.

v. SYSTYLÆ. Sepals deciduous. *Styles united into a protruding column.* St. arching or trailing, with uniform strong hooked prickles.

10. *R. bibracteata* (Bast.); prickles equal hooked, lts. simply serrate slightly downy beneath, sep. reflexed deciduous with a tapering point pinnate, fr. ovate with a *prominent disk.* *E. B.* 1895.—*St. erect-arching* to 8—12 ft. in height. Fl. pale pink, 3—6; ped. aciculate and setose; column of styles falling short of stamens, glabrous. Stalks of central germ. as long as their germens.—South of England. Sh. VI. VII. E.

11. *R. arvensis* (Huds.); prickles equal hooked, lts. simply serrate quite glabrous on both sides, sep. reflexed deciduous with a short point simply pinnate, fr. *subglobose* with a convex disk.—*E. B.* 188.—*St. trailing.* Lts. nearly flat. Fl. white, usually 4—6; ped. glandular; column of styles equalling the stamens, glabrous.—Hedges and thickets. Sh. VI. VII. E. S. I.

### *Index to Rosa.*

arvatica, 9.	dumetorum, 9.	sarmentacea, 9.
arvensis, 11.	farinosa, 5.	sepium, 8.
Bakeri, 9.	Forsteri, 9.	Sherrardi, 5.
bibracteata, 11.	hibernica, 2.	sphærica, 9.
biserrata, 9.	inodora, 8.	spinosissima, 1.
bractescens, 9.	involuta, 3.	subglobosa, 5.
Borreri, 9.	marginata, 9.	surculosa, 9.
cæsia, 9.	micrantha, 7.	tomentella, 9.
canina, 9.	mollis, 4.	tomentosa, 5.
coriifolia, 9.	pomifera, 3*.	Watsoni, 9.
Donniana, 3.	rubiginosa, 6.	Wilsoni, 3.
dumalis, 9.	Sabini, 3.	

Suborder III. *Pomeæ*.

## 15. CRATÆGUS Linn. Hawthorn.

1. *C. Oxyacan'tha* (L.); spinose, l. obovate 3—4-lobed cut and serrate cuneate at the base, fl. corymbose, cal. not glandular, styles 1—3.—*a. C. Oxyacantha* (L., Jacq.); l. lobed, *ped. and cal. usually glabrous*, cal.-lobes triangular-acuminate, styles 1—3, fruit oval, nuts 1—3. *Sy. E. B.* 479.—*β. C. monogyna* (Jacq.); l. deeply lobed usually acute, *ped. and cal. villose*, cal.-lobes lanceolate acuminate, style 1 bent, fruit subglobose. *E. B.* 2504.—Hedges and thickets; *β* the more common form and flowering later. T. V. VI. E. S. I.

## 16. COTONEAS'TER Lindl.

1. *C. vulgáris* (Lindl.); l. roundish-ovate rounded at the base, flowerstalks and margins of the calyx downy.—*E. B. S.* 2713.—Pet. rose-coloured. Fr. small pendulous, red.—Cliffs at the Great Orme's Head, Caernarvonshire. Sh. V. E.

## 17. MES'PILUS Linn. Medlar.

1. *M. german'ica* (L.); l. lanceolate undivided downy beneath, fl. solitary.—*E. B.* 1523.—L. entire simply or doubly serrate. In the wild state it is spinous.—Hedges and thickets in Surrey, Sussex, [and Cheshire?]. T. V. VI. E.

## 18. PY'RUS Linn.

1. *P. commúnis* (L.); l. ovate serrate, fl. corymbose, *fruit turbinate, styles distinct*.—*E. B.* 1784.—Branches rather spinous. Germen woolly. Leaves sometimes obovate suddenly contracted into a long very acute point.—*a. P. Pyra'ster* (Bor.); l. acuminate downy beneath when young ultimately glabrous, fr. elongate-pyriform obconical at the base.—*β. P. Ach'ras* (Bor.); mature l. acute or cuspidate slightly downy beneath, fr. globose-pyriform rounded below.—*γ. P. Briggsii* (Syme); l. cordate-ovate nearly glabrous, fr. small globose or pyriform.—Hedges and woods. T. IV. V. *Wild Pear-tree*. E. I.

2. *P. Má'lus* (L.); l. ovate acute serrate, fl. in a sessile umbel, *fr. globose, styles combined below*.—*α. P. acer'ba* (DC.); young branches calyx-tube and underside of the l. glabrous. *E. B.* 179.—*B. mit'is*; the same parts pubescent or woolly. *Sy. E. B.* 490.—Woods and hedges. T. V. *Crab-tree*. F. S. I.

[*P. domes'tica* (Sm.); l. *pinnate* downy beneath serrate, *fl. panicled, fr. obovate*.—Fr. resembling a small pear.—One tree in Wyre Forest; now lost.—T. V. *Service-tree*.] E.

3. *P. Aucupária* (Gaert.); *l. pinnate* downy beneath serrate, *fl. corymbose*, *fr.* (small red) *globose*.—Cormyb 4—6 in. across. Lts. 12—16.—Hilly woods. Mountains. T. V. VI. *Rowan-tree. Mountain-Ash.* E. S. I.

4. *P. fen'nica* (Syme); *l. oblong-serrate* usually pinnate below with 1—4 pairs of free decurrent lts. and 1—2 pairs of nearly free lobes, *lts. and lobes oblong or narrowly elliptical*, underside grey-webbed, *fl. corymbose*.—*Sorbus fennica* Fr.—Lobes blunt or acute as the term. tooth is more or less prominent. Hybrid (?) between Sp. 3 and 5.—Glen Catacol, Arran, S., in small quantity. S.

5. ? *P. scan'dica* (Syme); *l. oblong doubly serrate near the apex pinnatifid below, lobes oblong-lanceolate serrate*, underside white and downy, *fl. corymbose*.—*P. pinnatifida* Sm. *E. B.* 2331. *P. fennica* Bab. not Fr.—Fr. scarlet. Lower part of the *l.* often subpinnate, the lobes becoming more and more combined as they approach the extremity of the leaf which is only deeply and doubly serrate.—Glen Eis-na-Kearradh, Arran, in small quantity. T. V. S.

6. *P. A'ria* (Sm.); *l. of fl.-shoot roundly oval or elliptic*, entire below unequally and doubly serrate or with many small broad lobes especially towards the end, *lateral veins* 9—14 on each side, underside of *l.* uniformly snowy-white-felted, *fl. corymbose*.—*E. B.* 1858.—Lobes deepest a little below the end of the leaf.—Chalky banks and limestone rocks. T. V. E. I.

7. *P. rupic'ola* (Syme); *l. of fl.-shoot obovate narrow and entire below with many small broad lobes on the upper half, deepest near the end, lateral veins* 5—8 on each side, *underside of l. uniformly snowy-white-felted*, *fl. corymbose*.—*Sy. E. B.* 483.—Often the basal half of the edge of *l.* is entire.—Exposed rocks. T. V. E. I.

8. *P. latifolia* (Syme); *l. of fl.-shoot elliptic with triangular-oval acutely serrate lobes deepest near the middle of the sides of leaf, lateral veins* 5—9 on each side, *underside of l. ashy-felted flocculent*, *fl. corymbose*.—*Sy. E. B.* 484. *P. scandica* Bab. not Fr.—*L.* often very acute with very acute lobes which are longer than broad, basal  $\frac{1}{4}$  finely serrate. Fr. dark or reddish brown.—Hilly woods. T. V. E.

9. *P. torminális* (Sm.); *l. ovate or cordate lobed glabrous, lobes triangular acute serrate the lower ones larger and spreading, fl. corymbose*.—*E. B.* 298.—Fr. oval brown.—Woods and hedges in the South. T. IV. V. *Wild Service-tree.* E.

## Order XXVII. LYTHRACEÆ.

Cal. tubular, lobed; valves valvate or distant in the bud, sometimes with intermediate teeth. Pet. between the lobes of the calyx, very deciduous. Stam. inserted in the tube of the cal., below the pet. and equalling them or 2, 3, or 4 times as many. Ovary free, 2—4-celled. Style 1. Caps. membranaceous, usually (by abortion) 1-celled, with many seeds and a central placenta, covered by the calyx. Embryo straight.—Stip. 0.

1. LYTHRUM. *Cal. tubular, cylindrical*, with 8—12 teeth; 4—6 broader, erect; alternate teeth subulate. Pet. 4—6. Stam. as many as or twice the number of the petals. *Style filiform*. Caps. 2-celled many-seeded.
2. PEPLIS. *Cal. bellshaped* with 12 teeth, of which 6 are broader and erect, 6 subulate. Pet. 6, minute, fugacious. Stam. 6. *Style very short*. Caps. 2-celled, many-seeded.

## 1. LYTHRUM Linn. Loosestrife.

1. *L. Salicária* (L.); *l. lanceolate* from a cordate base opposite or whorled, fl. in whorled leafy spikes, bracts 0, subulate calyx-teeth twice as long as the others, stam. 12.—*E. B.* 1061.—Upper l. usually falling short of the flowers, or so large as totally to destroy the spiked appearance of the plant. St. 2—4 feet high and l. nearly glabrous, or downy with crisped hairs. Fl. large, purple or crimson.—Ditch-banks and damp places. P. VII. VIII. *Purple Loosestrife*. E. S. I.

2. *L. Hyssopifolia* (L.); *l. alternate linear-lanceolate* blunt, fl. axillary solitary, bracts 2 minute subulate, calyx-teeth all short, stam 6.—*E. B.* 292.—St. mostly procumbent, spreading, simple or branched. Fl. small, light purple. Glabrous.—Damp places where water has stagnated, rare. A. VI.—X. E. I.\*

## 2. PEP' LIS Linn. Water-Purslane.

1. *P. Portula* (L.); l. opposite obovate stalked, fl. axillary solitary sessile.—*E. B.* 1211. St. 1. 7.—Pet. often wanting. Cal.-tube short, shortly bell-shaped, shorter than the capsule. St. 4—6 in. long, prostrate, creeping.—Damp places. A. VII. VIII. E. S. I.

## Order XXVIII. TAMARISCACEÆ.

Cal. 4—5-parted, persistent, imbricate in the bud. Pet. 4—5, withering, from the base of the calyx. Stam. equal to or twice as many as the pet., from the margin of a shieldlike disk. Caps. 1-celled, 3-valved, many-seeded, loculicidal. Placentas often

only at the base. Seeds ascending, crowned with a tuft of hairs.

1. TAMARIX. Styles 3, patent. Seeds from base of capsule; crowned with a tuft of simple papilliform hairs.

1. TAM'ARIX *Linn.* Tamarisk.

[1. *T. anglica* (Webb); l. glabrous rather narrowed at the base spurred, fl.-buds ovoid, hypogynous ring with 5 angles narrowed into the filaments of the cordate shortly apiculate anthers, caps. roundish-trigonus at the base abruptly narrowed towards the apex.—*T. gallica* Sm. *E. B.* 1318. St. shrubby, with slender leafy branches. L. minute.—Spikes lateral, rather paniced, slender. Fl. small, pink.—South-west coast of England. Planted. Sh. VII.] E.

Order XXIX. ONAGRACEÆ.

Cal. tubular, adnate to the ovary wholly or in part, with 2—4 lobes valvate in the bud. Pet. as many as the calyx-lobes, twisted in the bud, inserted at the top of the tube. Stam. 2, 4, or 8, inserted with the petals. Ovary of several cells, with a central placenta. Style 1, filiform, stigma capitate or lobed. Fr. a berry or capsule with 4 cells. Albumen 0.—L. alternate or opposite, not dotted. Exstipulate. Many raphides.

1. EPILOBIUM. Cal. 4-cleft, deciduous. Pet. 4. Stam. 8. Style filiform, stigma clavate or cruciform. Caps. linear, 4-celled, 4-valved. Seeds many, bearded.—Fl. not yellow.
2. CENOTHERA. Cal.-limb tubular below. Seeds not bearded. In other respects like *Epilobium*.—Pet. yellow.
3. LUDWIGIA. Limb of the cal. 4-cleft, persistent. Pet. 4, or 0 in our plant. Stam. 4. Style filiform, deciduous; stigma capitate. Caps. obovate, 4-valved, 4-celled, many-seeded, with a loculicidal dehiscence.
4. CIRCÆA. Limb of the cal. 2-cleft, deciduous, its tube closed by a cupshaped disk. Pet. 2, obcordate. Stam. 2, alternate with the petals. Style simple; stigma emarginate. Caps. 1- or 2-celled, cells 1-seeded, seeds erect. Fl. white.

1. EPILOBIUM *Linn.* Willow Herb.

A. *Fl. irregular, cal. without a free tube, stam. and style ultimately declining, l. scattered.*

1. *E. angustifolium* (L.); st. erect round, l. lanceolate veined,

pet. obovate shortly clawed, style exceeding the stamens.—Rhizome far-creeping. St. 3—6 ft. high. Fl. crimson.—*a. E. macrocarpum* (Steph.); l. lanceolate, fl.-buds obovate cuspidate, caps. long ( $2\frac{1}{2}$  in.) erect. *Sy. E. B.* 495.—*β. E. brachycarpum* (Leight.); l. lanceolate-attenuate (broadest near their base), fl.-buds narrowly obovate obliquely acute, caps. short (1 in.) spreading. *Sy. E. B.* 496.—Damp shady places. *β.* the cultivated form. P. VII. *Rose-bay.* E. S. I.

2. *E. rosmarinifolium* (Haenke); st. erect round, l. linear not veined, pet. elliptic-oblong not clawed, style equalling the stamens.—*Sy. E. B.* 494. *E. Dodonæi* St. 72. 5.—Rhizome creeping moderately. St. dwarf, often prostrate. L. shortly attenuate at both ends. Fl. rose-coloured or white.—Stony places by the Tarf in Glen Tilt. *The late Mr. John Robertson!* P. VIII. S.

*B. Fl. regular, cal. with a free tube, Stam. and style erect, l. opposite, upper l. lanceolate alternate.*

*a. Turionate, i. e. producing radical suckers.*

3. *E. hirsutum* (L.); st. woolly glandular, l. clasping slightly decurrent oblong-lanceolate denticulate-serrate, buds erect mucronate, sep. lanceolate, seeds tubercular oblong acute below.—*E. B.* 838.—Suckers thick, fleshy, with distant scales; others leafy and ending in rosettes. St. 4—5 ft. high, terete, branched. L. most hairy upon the ribs. Fl. large.—Wet places by ditches and streams. P. VII. VIII. *Great Willow Herb.* E. S. I.

*b. Stoloniferous. \* Stoles autumnal rosulate, st. erect.*

† Stem mostly terete, stigmas 4-cleft.

4. *E. parviflorum* (Schreb.); st. downy, l. sessile lanceolate from a rounded base denticulate, lowest l. shortly stalked, buds erect ovoid, sep. lanceolate, seeds tubercular obovate-oblong rounded below.—*E. B.* 795.—Autumnal rosettes on short stalks or sessile. St. 1—2 ft. high, nearly simple. L. uniformly hairy. Fl. small.—*E. rivulare* (Wahl.) is a subglabrous state.—Damp places. P. VII. VIII. E. S. I.

5. *E. montanum* (L.); st. downy, l. shortly stalked ovate-lanceolate from a rounded base denticulate, buds nodding ovoid, sep. lanceolate, seeds tubercular oblong blunt at both ends or rather narrowed below.—*E. B.* 1177.—Root truncate. Rosettes fleshy, oblong, sessile. St. 6—24 in. high. Edges and veins of l. pubescent. L. sometimes in threes.—Dry places. P. VI. VII. E. S. I.

6. *E. lanceolatum* (S. & M.); st. slightly angular downy, l. stalked lanceolate denticulate-serrate narrowed to an entire base, buds nodding ovoid, sep. broadly linear acuminate, seeds tubercular obovate-oblong subacute below.—*E. B. S.* 2935.—Root not truncate. Rosettes loose, sessile. St. 1—2 ft. high, obscurely or very bluntly angled especially below. L. flaccid, mostly pendulous.—South of England, rare. P. VII.—IX. E.

†† Stem with raised lines, stigma entire.

7. *E. roseum* (Schreb.); rosettes loose, l. stalked ovate narrowed and acute at both ends serrulate, buds usually nodding ovoid acuminate, sep. lanceolate acuminate, seeds obovate-oblong narrowed to a rounded base.—*E. B.* 693.—St. 1—2 ft. high, branching, with two sharp and two blunt angles. Petioles long.—Damp places. P. VII. VIII. E. S. ? I.

8. *E. tetragonum* (L.); rosettes dense subsessile, l. strap-shaped much denticulate-serrate, limb of interm. l. decurrent, buds erect, seeds oblong-obovate tubercular.—*Sy. E. B.* 502. *Curt.* i. 66.—St. 1—2 ft. high, with 2—4 raised lines, usually much branched.—Damp places. P. VII. VIII. E. S.

\*\* *Stoles æstival long-jointed with small leaves, primary stem erect, stigma usually entire.*

9. *E. obscurum* (Schreb.); l. tapering from a rounded base sessile remotely denticulate, lower l. oblong blunt, buds erect, seeds oblong-obovate.—*E. virgatum* G. & G., *Sy. E. B.* 503.—Resembling *E. tetragonum*, but the caps. much shorter and l. very different.—Wet places. P. VII. VIII. E. S. I.

\*\*\* *Stoles æstival long-jointed with very small leaves ending in autumnal bulbs which become detached, base of stem cordlike.*

10. *E. palustre* (L.); stoles long slender, st. rooting near the base, l. narrowly lanceolate from a wedged shaped base sessile not decurrent, top of raceme nodding, sep. lanceolate, seeds subfusiform attenuate and acute below and the testa prolonged above smooth.—*Sy. E. B.* 504.—St. round, often with two rows of down, 6—18 in. high. L. entire or denticulate.—*F. ligulatum* (Baker) is a broad-leaved form.—Bogs. P. VII. VIII. E. S. I.

\*\*\*\* *Stoles or barren stems æstival rosulate.*

11. *E. alpinum* (L.); barren stems short their upper l. closely placed, fl.-st. erect from a short rooting base, l. oval or oblong blunt narrowed below not acuminate, upper l. lanceolate, buds nodding, sep. linear-lanceolate acute, seeds lanceolate-obovate pointed below apiculate.—*E. B.* 2001.—St. filiform, simple,

with two raised lines, 3—4 in. long. L. pale green. Fl. pale. —Higher mountains. (Scandinavian.) P. VII. S.

\*\*\*\*\* *Stoles or barren stems æstival leafy not rosulate.*

12. *E. anagallidifolium* (Lam.); *joints of barren stems all long* with small obovate leaves, fl.-st. erect from a long rooting base, l. oblong blunt narrowed below not acuminate, upper l. lanceolate buds nodding, *sepals oblong blunt*, seeds obovate pointed below apiculate.—*Sy. E. B.* 506.—St. filiform, mostly simple, with 2 raised lines, 3—4 in. long. L. pale green. Fl. pale reddish.—Higher mountains. (French.) P. VII. E. S.

\*\*\*\*\* *Stoles æstival scale-bearing not rosulate.*

13. *E. alsinifolium* (Vill.); *stoles (yellowish) with small roundish distant scales*, st. erect from a long rooting base, l. *ovate-acuminate repand-dentate* shortly stalked, buds nodding, *sep. linear-oblong, seeds subfusiform*.—*Sy. E. B.* 505.—St. mostly simple, rather thick, with 2 raised lines, 3—12 in. long. L. shining, subpellucid. Fl. large, purplish. Caps. upright, long, long-stalked.—Higher mountains. P. VII. E. S.

## 2. CENOETHERA Linn. Evening Primrose.

\*1. *C. bien'nis* (L.); l. obovate-lanceolate flat toothed, pet. longer than the stamens, st.-l. elliptic or ovate-lanceolate, caps. roundly tetragonal *tapering upwards*.—*E. B.* 1534. *St.* 5. 5. —Fl. large, many, bright yellow. St. 2—3 ft. high, leafy. —Sandy coast of Lancashire. (American.) B. VII.—IX. E.

\*2. *C. odorata* (Jacq.); l. subsessile, radical l. linear-lanceolate toothed, st.-l. ovate-lanceolate attenuate, pet. longer than the stamens, caps. clavate.—*Sy. E. B.* 509.—Fl. large, bright yellow. St. 2—3 ft. high. Foliage brighter green and fl. deeper yellow than in Sp. l.—Channel Isles. Coasts of Som. and at Plymouth. (Patagonian.) B. VII.—IX. E.

## 3. LUDWIGIA Linn.

1. *L. palustris* (Ell.); st. procumbent rooting glabrous, l. opposite ovate acute narrowed into a petiole, fl. axillary solitary sessile without petals.—*Isnardia* L. *E. B. S.* 2593. *St.* 22. 3. —St. 6—8 in. long, round, branching, often reddish. Fl. with 2 small bracts at the base. Caps. ovate obtusely quadrangular, with the persistent calyx spreading horizontally.—Pools and marshes. Buxted, Suss., and Petersfield, Hants. Said to be lost in these places. New Forest. Jersey. A. VI. E.



## 4. CIRCÆA Linn.

1. *C. lutetiána* (L.); *l. ovate* or slightly cordate below repand-denticulate *opaque*, *petioles subterete*, *bracteoles* 0, *pet.* deeply emarginate, lobes broadly obovate, *cal.* hairy, *ovary* 2-celled, *fr.* broadly obovate.—*E. B.* 1056. *St.* 23. 1.—*Pet.* as long as the herbaceous sepals, broad below. *Fr.* persistent.—Woods and hedge-banks. P. VI.—VIII. *Enchanter's Nightshade.* E. S. I.

2. *C. alpína* (L.); *l. cordate* acuminate repand-dentate shining, *petioles flat* with membranous wings, *bracteoles setaceous*, *pet.* bifid, lobes oblong, *cal.* glabrous, *ovary* 1-celled.—*E. B.* 1057. *St.* 23. 2.—Usually glabrous. *Pet.* shorter than the membranous sepals, narrow below. *Fr.* soon falling. *Bracteoles* deciduous.— $\beta$ . *C. intermedia* (Ehrh.); *fl.* larger, *sep.* less membranous equalling the petals. Whole plant much larger.—Woods and thickets in hilly districts. P. VII. VIII. E. S. I.

## Order XXX. HALORAGACEÆ.

*Cal.* adnate to the ovary, limb minute. *Pet.* minute, from the throat of the calyx. *Stam.* 1—8, inserted with the petals. *Ovary* of 1 or more cells. *Styles* equal in number to the cells. *Fr.* dry, not bursting, usually crowned with rim of calyx. *Seeds* pendulous; albumen fleshy.—*Stip.* 0.

1. MYRIOPHYLLUM. Monœcious. *Cal.* 4-parted. *Pet.* 4, soon falling, exceeding the calyx in the male, small and reflexed or 0 in the female. *Stam.* 8. *Styles* 4, villose. *Fr.* tetragonal, separable into 4 hard nuts.
2. HIPPURIS. *Calyx-limb* very minute, obsoletely 2-lobed. *Pet.* 0. *Stam.* 1. *Style* filiform. *Stigma* simple, acute. *Fruit* a 1-celled nut.

## 1. MYRIOPHYLLUM Linn. Water-Milfoil.

1. *M. verticillatum* (L.); *fl.* all axillary whorled, bracts pinnatifid.—*E. B.* 218.—*L.* whorled, pinnatifidly divided into setaceous segments. Bracts more or less longer than the flowers, pinnatifid or (*M. pectinatum* DC.) pectinate.—Ponds and ditches, rare. P. VII. VIII. E. I.

2. *M. spicatum* (L.); *fl.* whorled forming a leafless spike, bracts small entire, spike erect when in bud.—*E. B.* 83.—*L.* 4 in a whorl, submersed. Lower bracts often pectinate.—Ponds and ditches. P. VI. VII. E. S. I.

3. *M. alterniflorum* (DC.); sterile fl. alternate about 6 forming a leafless spike, *spike nodding when in bud* afterwards erect, *fertile fl.* about 3 together in *axillary whorls* at the base of the spike.—*E. B. S.* 2854.—L. 3 or 4 in a whorl, submersed.—Ponds and ditches. P. V.—VIII. E. S. I.

## 2. HIPPU'RIS Linn. Mare's-tail.

1. *H. vulgaris* (L.); l. linear 6—12 in a whorl with a hard point.—*E. B.* 763. St. 44. 1.—St. simple, or sometimes branching at the base, erect. Fl. in the axil of each of the upper leaves, often without stamens. In deep water the submersed leaves are long flaccid pellucid and not hard at the end; in running water often wholly submersed flaccid and barren.—In stagnant water and slow streams. P. VI. VII. E. S. I.

## Order XXXI. CUCURBITACEÆ.

Cal. 5-toothed, tube adnate to the ovary. Cor. 5-cleft, often scarcely distinguishable from the calyx, netted with veins. Stam. 5, more or less cohering. Anth. sinuous. Ovary 3—5-celled or spuriously 1-celled; placentas parietal. Style short. Stigmas lobed. Fr. more or less succulent. Seeds flat, in an aril; embryo flat; albumen 0.—Succulent, climbing with extra-axillary tendrils; often monœcious or diœcious. Stip. 0.

1. BRYONIA. Cal. 5-toothed. Cor. 5-cleft. Male: Stam. 5 in 3 bundles. Fem.: Style 3-fid. Fruit a globose few-seeded berry. Seeds oval, compressed, more or less bordered.

## 1. BRYO'NIA Linn. Red Bryony.

1. *B. dioica* (L.); l. palmate 5-lobed dentate rough on both sides with hard points, fl. diœcious, cal. of the fertile fl. half as long as the corolla.—*Sy. E. B.* 517.—St. climbing. Tendrils simple. Fr. red.—[*B. alba* (L.) has the fertile cal. as long as the cor. and black fruit. It is said to be monœcious.]—Hedges and thickets. P. V.—IX. E.

## Order XXXII. PORTULACEÆ.

Sep. 2, rarely 3 or 5, cohering at the base; imbricate in the bud. Pet. usually 5, from the base of the calyx. Stam. indefinite, inserted with the petals, often opposite to the petals; filaments distinct. Ovary 1-celled. Style 0. Stigmas several. Caps. opening transversely or by 3 valves; placenta central. Embryo curved round the albumen.—Stip. scarious.

1. MONTIA. Cal. of 2 sepals, persistent. Cor. 5-parted, with 3 segments smaller than the others, tube split to the base in front. Stam. 3, inserted in the throat and opposite to the smaller segments of the corolla. Ovary turbinate. Style very short. Stigmas 3. Caps. 3-valved, 3-seeded.
2. CLAYTONIA. Cal. of 2 sepals, persistent. Cor. of 5 free petals. Stam. 5, opposite to and adhering to the petals. Style 3-cleft. Caps. globose, 3-valved, 3-seeded.

## 1. MON'TIA Linn. Blinks.

1. *M. fontána* (L.).—*E. B.* 1206. *St.* 11. 1.—L. opposite, spathulate, entire. Valves of the caps. rolled longitudinally inwards after the seeds have fallen.—*a. M. minor* (Gm.); st. short ascending rigid, ped. axillary and terminal, seeds netted-asperous.—*β. M. rivularis* (Gm.); st. flaccid ascending, ped. axillary, seeds netted.—Watery places. *β* in water. A. IV.—VIII. E. S. I.

## 2. CLAYTO'NIA Linn.

\*1. *C. perfoliáta* (Donn); radical l. fleshy rhomboidal-spathulate, upper st.-l. 2 connate, raceme subverticillate with 1 or 2 basal flowers, pet. nearly or quite entire.—*Sy. E. B.* 260.—St. tufted, ascending, 4—12 in. high. Pet. white, just overtopping the calyx.—Naturalized. (American.) A. IV.—VI. E. S.

\*2. *C. alsinoides* (Sims); radical l. ovate-acuminate, upper l. opposite sessile, fl. mostly solitary in raceme, pet. bifid.—St. often 1 ft. high. Fl. more and larger than in *Sp.* 1.—Naturalized. (N.W. American.) A. IV.—VII. E. S.

[*Portulaca oleracea* is a common weed near Richmond, Surrey.]

## Order XXXIII. PARONYCHIACEÆ.

Cal. 5-parted, imbricate in the bud. Pet. usually minute and resembling abortive stam., or 0. Stam. indefinite, opposite the sep. (when the same number), somewhat hypogynous. Ovary free. Styles 2 or 3. Fr. dry, 3-valved or indehiscent. Embryo more or less curved. Albumen mealy.—With stipules.

1. CORRIGIOLA. Sep. 5. *Pet.* 5, oblong, equalling the sepals. Stam. 5. *Stigmas* 3, sessile. Fr. 1-seeded, indehiscent. Seed suspended by its seedstalk, which arises from the base of the capsule.—L. alternate.
2. HERNIARIA. Sep. 5. *Pet.* 5, filiform, inserted with the 5 stam. on a perigynous ring. *Stigmas* 2, nearly sessile. Fr. 1-seeded, indehiscent, membranous.—L. opposite.

3. ILLECEBRUM. *Sep.* 5, thickened, *horned at the back*. Pet. 0 or 5, subulate, inserted with the 5 stam. on a perigynous ring. Stigmas 2. *Fr.* 1-seeded, *furrowed, bursting along the 5 furrows*.

1. CORRIGI'OLA *Linn.* Strapwort.

1. *C. littoralis* (L.); st. leafy amongst the flowers.—*E. B.* 668.—Fl. stalked, white, small, in small clusters. Stem-1. oblong, narrow below. St. many from the crown of the root, prostrate, slender.—Sandy shores. Slapton Sands and near the Start Point, Devon. Helston, Cornwall. A. VII. VIII. E.

2. HERNIA'RIA *Linn.* Rupturewort.

1. *H. glabra* (L.); st. herbaceous prostrate clothed all round with minute decurved hairs, l. narrowed below, clusters of sessile fl. axillary collected on the lateral branches into a slightly leafy spike.—*E. B.* 206.—Pale yellowish green. St. procumbent or subterranean; autumnal shoots ascending, irregularly branched with spreading not imbricate leaves. Lateral branches resembling leafy spikes from the dense aggregation of the clusters. Cal. glabrous or with a very few hairs. L. glabrous.— $\beta$ . *subciliata* (Bab.); l. more or less ciliated.—Rare. Suffolk. Ruan Minor, Cornwall. Wilsford, Lincoln. Six-mile Bottom, Camb. Finchley Common. A. or B. VII. E.

2. *H. ciliata* (Bab.); st. suffrutescent prostrate clothed on its upperside alone with minute decurved hairs, l. orbicular-ovate ciliate, clusters of sessile fl. axillary upon the lateral branches and distinct.—*E. B. S.* 2857.—Dark green. Root strong, woody. St. spreading extensively from the crown of the root but scarcely rooting until the autumn, when they are prostrate and regularly alternately branched and the leaves are imbricate in two rows. Clusters 1—3 together in sma'l distinct bunches. Fl. larger than in Sp. 1. *Sep.* tipped with a strong bristle.—Very rare. Lizard Point. Guernsey. P. VII. VIII. E.

[*H. hirsuta* (L.); st. herbaceous prostrate clothed with straight spreading hairs, l. oblong narrowed at both ends, cal. hairy resembling a minute bur. *E. B.* 1379.—Sandy ground at Christchurch, Hants. *Mr. Townsend.* The Finchley-Common plant was *H. glabra*!]

3. ILLE'CEBRUM *Linn.*

1. *I. verticillatum* (L.).—*E. B.* 895.—St. procumbent, filiform, glabrous. Fl. whorled, axillary, minute, white. L. roundish,

variable in size, shorter or longer than the flowers.—Boggy places in Devon and Cornwall. P. VII. E.

### Order XXXIV. CRASSULACEÆ.

Sep. 3—30, more or less united at the base. Pet. the same number, free or slightly connected, inserted at the base of the calyx. Stam. inserted with the pet. and the same or twice their number. Hypogynous scales 1 at the base of each carpel or inconspicuous. Carpels the same number as and opposite to the pet., free or slightly connected, 1-celled. Fr. of several follicles opening on their face, with slightly albuminous seeds on the inner suture.—Exstipulate.

1. TILLÆA. Sep., pet., and stam. 3—4. Carp. 3—4, constricted in the middle and 2-seeded. Hypogynous scales 0.
2. SEDUM. Sep. and pet. 5, rarely 4 or 6. Stam. 10 or 12. Hypogynous scales entire. Carp. 5 or 6 many-seeded.—*S. Rhodiola* has 4 sep., 4 pet., 8 stam., 4 carp., and is subdieceious.
- [3. SEMPERVIVUM. Sep. 6—10. Pet. the same. Stam. twice as many as the pet. *Hypogynous scales laciniate.* Carp. as many as the petals.]
4. COTYLEDON. Sep. 5. Pet. cohering into a tubular 5-cleft corolla. Stam. 10, inserted on the corolla. Hypogynous scales 5. Carp. 5.

#### 1. TILLÆA Linn.

1. *T. muscosa* (L.); st. branched and decumbent at the base, fl. axillary sessile trifid.—*E. B.* 116. *R. I. t.* 191.—Very minute, about 1 in. long, reddish. L. opposite, oblong, blunt, concave above, connate. Sep. ovate or lanceolate, acute, bristle-pointed. Pet. nearly subulate, white tipped with red.—On barren sandy heaths in the South and East. A. VI. VII. E.

#### 2. SEDUM Linn. Stonecrop.

\* *Rootstock thick, many-headed. Leaves flat. Stems annual.*

† Flowers yellow, subdieceious, 4-parted. RHODIOLA Linn.

1. *S. Rhodiola* (DC.); root fleshy, l. oblong smooth.—*Rhodiola rosea* L. *E. B.* 508.—Fl. in a compact terminal cyme, subdieceious. Stam. 8. L. acuminate, usually dentate in their upper half. Rhizome large, having a remarkable smell. St. 6—8 in. high, simple.—Wet alpine rocks. P. VI. VII. *Rose-root.* E. S. I.

†† Flowers white or purple, perfect, 5-parted.

2. *S. Teléphiium* (L.); uppermost l. long-oval dentate rounded at the base and sessile, lower l. obovate or oblong narrowed below, corymb dense, ovaries flattened and slightly furrowed on the back.—*R. I.* 968. *S. purpurascens* Koch, *Sy. E. B.* 526.—St. 1—2 feet high. L. large, often orbicular. Pet. recurved from the middle, faintly channelled at the end. Stam. 10.—[*S. maximum* Sut., *R. I.* 969, has lower l. broad, uppermost cordate at the base, back of the ovaries convex.]—Hedge-banks and thickets. P. VII. VIII. *Orpine. Live-long.* E. S. I.

3. *S. Fabária* (Koch); l. dentate oblong-lanceolate or lanceolate all narrowed to a slight petiole, corymb dense, ovaries not furrowed.—*Sy. E. B.* 527. *St.* 83. 9.—Resembling *S. Telephiium*. Pet. spreading. Stam. 10.—Hedges and thickets. P. VIII. *Orpine. Live-long.* E. I.

\*\* Root small, weak, without any rooting shoots. L. subterete.

4. *S. villósum* (L.); st. erect, l. linear blunt flat above not produced at the base, pet. ovate acute, l. st. and panicle glandular-pubescent.—*E. B.* 394. *St.* 6. 12.—Pet. rose-coloured with a purple streak. *St.* 3—4. in. high. Seedlings with a rosette.—Wet mountain pastures. B. VI. VII. E. S.

\*\*\* Root small, producing rooting shoots. Leaves subterete.

† Flowers white. Leaves blunt.

5. *S. al'bum* (L.); flowering st. erect, barren st. procumbent rooting, l. oblong subcylindrical flattened above spreading, pet. lanceolate, panicle much branched glabrous.—*E. B.* 1578.—Pet. white. *St.* 4—5 in. high, purplish, leafy.—[*β. S. micranthum* (Bast.); l. oblong-obovate flattened on both sides blunt. *Sy. E. B.* 529.]—Rocks and walls, not common. [*β. Arundel.*] P. VII. VIII. E.

6. *S. dasyphyl'lum* (L.); flowering and barren st. procumbent, l. ovoid fleshy gibbous, pet. and sep. ovate blunt, panicle small glandular-pubescent.—*E. B.* 656.—Pet. white. Sep., pet., and carp. often in sixes. L. very short and thick, glaucous, often tinged with red, opposite on the barren shoots. Flowering st. 3—4 in. long.—Rocks and walls, rare. P. VI. VII. E. I.

7. *S. ang'licum* (Huds.); st. procumbent at the base ascending, l. ovoid fleshy gibbous spurred at the base, pet. lanceolate acute, sep. ovate bluntish, cyme bifid smooth.—*E. B.* 171.—Pet. white spotted with red. L. mostly alternate. Flowering st. 3—6 in. long.—Sandy and rocky places. A. VII. VIII. E. S. I.

†† Flowers yellow. † Leaves blunt.

8. *S. ácre* (L.); l. ovoid thick tumid spurred at the base, pet. lanceolate acute, sep. ovate blunt gibbous at the base, cyme trifid smooth.—*E. B.* 839.—L. closely imbricate on the barren shoots, very acrid.—Walls and dry places. P. VI. VII. *Wall-Pepper.*  
E. S. I.

[*S. sexanguláre* (L.); l. linear terete spurred at the base, pet. lanceolate acute, sep. lanceolate acute not gibbous, cyme trifid smooth.—*E. B.* 1946.—Old walls; a doubtful native. P. VII.]  
E.

†† Leaves acute.

9. *S. reflex'um* (L.); l. subulate scattered spurred at the base convex on both sides the lowermost recurved, fl. cymose, sep. ovate rather acute, pet. lanceolate blunt.—*E. B.* 695.—Barren st. long. L. many, patent or reflexed. Cyme nearly level-topped, its outer branches spreading or recurved. Base of filam. and lateral edges of carp. with glandular hairs.— $\beta$ . *S. albescens* (Haw.); l. more slender glaucous not recurved. *S. glaucum* Sm. *E. B.* 2477.—Walls and rocks, common.  $\beta$ . Dry hills rare. P. VII. VIII.  
E. S. ? I.

10. *S. rupes'tre* (Huds.); l. linear-lanceolate spurred at the base flattened, fl. imperfectly cymose, sep. elliptic, pet. lanceolate.—*E. B.* 170. *S. elegans* *Lej.*—Barren st. short with densely imbricate adpressed glaucous leaves. Fl. corymbose rather than cymose. Filam. and ovaries glabrous.—On limestone rocks. Bristol. Cheddar. Orme's Head, &c. P. VI. VII. E. I.

11. *S. Forsteriánum* (Sm.); l. lanceolate spurred at the base flattened, cyme round-topped compact, sep. ovate, pet. lanceolate.—*E. B.* 1802.—Barren st. short, erect, densely leafy. L. forming small roselike tufts, bright green. Fl. truly cymose. Filam. and ovaries glabrous. Very different from the last in appearance; but there are 2 forms, (1) with short obconic barren stems and green leaves, (2) with very short globular barren stems and glaucous leaves.—Damp rocks in Wales, Shropshire, and West Somerset. P. VI. VII. E.

### 3. SEMPERVIVUM Linn. Houseleek.

[*S. tectorum* (L.); l. glabrous ciliate obovate-oblong, pet. 12 or more entire at the margins, hypogynous scales short convex resembling glands.—*E. B.* 1320.—L. succulent, green with purple tips, forming large roselike tufts.—Walls and roofs (planted). P. VII.]  
E. I.

4. COTYLE'DON *Linn.* Navelwort.

1. *C. Umbilicus* (L.); lower l. peltate concave, bracts entire, fl. pendulous.—*E. B.* 325.—Raceme usually simple. St. 6—12 in. high, leaves mostly collected at its base. Fl. greenish yellow.—Chiefly on rocks and walls in the West. P. VI.—VIII.

E. S. I.

## Order XXXV. RIBESIACEÆ.

Cal. superior, 4—5-cleft, regular. Pet. 4—5, small, inserted at the mouth of the tube and alternating with the stamens. Ovary 1-celled, with 2 opposite parietal placentas. Style 2—4-cleft. Berry many-seeded; cell filled with pulp. Albumen horny. Stip. 0.

1. RIBES. Cal. 5-cleft. Berry many-seeded, crowned with the persistent calyx.

1. RI'BES *Linn.*

\* *Peduncles* 1—3-flowered. *Stems spinous.* Gooseberry.

†1. *R. Grossulária* (L.); ped. with 2 minute bracts, cal. bell-shaped, sep. reflexed oblong, pet. ovate.—*E. B.* 1292 and 2057.—Thorns 1—3, at the base of the young branches. Germen and fruit smooth or pubescent or glandular. L. rounded, 3—5-lobed and cut, glabrous or hairy.—Hedges and thickets, doubtfully native. Sh. IV. V. E. S.

\*\* *Flowers racemose.* *Spines none.* Currants.

2. *R. alpinum* (L.); dioecious, *racemes upright* both in flower and fruit glandular-pilose, *bracts longer than the pedicels lanceolate, cal. glabrous*, l. shining beneath.—*E. B.* 704. St. 5l. 6.—Male raceme dense, of many flowers; fem. of 2—5 fl. Limb of the calyx nearly flat. Berries scarlet. L. with 3 acute deeply serrate lobes.—Woods. North of E., rare. Sh. IV. V. E.

3. *R. nigrum* (L.); *racemes pendulous* downy with a separate simple pedicel at the base of each, *bracts subulate* shorter than the pedicels, *cal. pubescent*, l. glandular-punctate beneath.—*E. B.* 1291.—Racemes lax. Calyx-limb bell-shaped. Berries large, black. L. with 3—5 acute serrate lobes.—In damp and swampy places. Sh. IV. V. *Black Currant.* E.

4. *R. rubrum* (L.); *racemes mostly glabrous and pendulous, bracts shorter than the pedicels ovate, cal. nearly flat glabrous*, l. bluntly 5-lobed.—*E. B.* 1289.—*a. sativum* (R.); l. glabrous, raceme glabrous.—*β. R. petraeum* (Sm. not Wulf.); racemes



slightly downy, in flower upright, in fruit pendulous. *E. B.* 705.—*R. spicatum* (Robs.); racemes erect with fl. and fr., “fl. nearly sessile,” was perhaps a sport from this. *E. B.* 1290.—Woods.  $\beta$ . North of E., S. Sh. IV. V. *Red Currant*. *E. S.*

### Order XXXVI. SAXIFRAGACEÆ.

Cal. 4—5-cleft, superior or inferior. Pet. 4—5, rarely 0. Stam. 5—10, free, perigynous or hypogynous. Glandular disk present or wanting. Ovary of 2 carpels cohering by the inflexed sides or margins. Styles 2, persistent, usually diverging. Seeds many. Albumen fleshy.—Stip. 0.

1. **SAXIFRAGA.** Cal. 5-fid or 5-parted, more or less adnate to the ovary or free. Cor. of 5 petals. Stam. 10, rarely 5. Styles 2, persistent. Caps. 2-celled, with 2 beaks, opening by a pore between the beaks.
2. **CHRYSOSPLENIUM.** Cal. 4-fid, half superior. Cor. 0. Stam. 8, rarely 10. Styles 2. Caps. 1-celled, with 2 beaks, opening into the form of a cup.
3. **PARNASSIA.** Cal. 5-cleft, inferior. Pet. 5. Stam. 5, perigynous, with 5 scales bearing glandular bristles interposed. Stigm. 4, sessile. Caps. 1-celled, with 4 valves.

#### 1. SAXIFRAGA *Linn.* Saxifrage.

*Sec.* 1. With decumbent barren shoots at the base.

\* *Cal. reflexed inferior, fl. paniced, flowering stems leafless.*

† Filaments enlarged upwards. **ROBERTSONIA** *Haw.*

1. *S. umbrósa* (L.); *l. obovate* with cartilaginous crenatures or sharp notches tapering at the base into *dilated footstalks with flat edges*, panicle racemose.— $\alpha$ ; *l. crenate* or dentate spreading. *E. B.* 663.— $\beta$ . *S. punctata* (Haw.); *l. nearly round acutely serrate* erect. *R. I. t.* 622, 623.— $\gamma$ . *S. serratifolia* (Mack.); *l. oblong acutely serrate* erect. *E. B. S.* 2891.—West of Ireland. Yorkshire (var.  $\alpha$ ), a doubtful native. P. VI. *London Pride*. *St. Patrick's Cabbage*. [E.] I.

2. *S. elegans* (Mack.); *l. round smooth shining acutely serrate, footstalks broad convex beneath with flat edges*, panicle racemose.—*E. B. S.* 2892.—*L. not tapering into the footstalks*.—

Very rare. Top of Turk Mountain, Killarney, and Connor Hill, Kerry. P. VI. I.

3. *S. hirsuta* (L.); l. oval sharply serrate, footstalks linear semicylindrical with raised edges hairy, panicles racemose.—*E. B.* 2322.—Footstalks slightly tapering upwards. L. longer than broad.—Very rare. Gap of Dunloe and Connor Hill, Kerry. Hungry Hill, Co. Cork. P. VI. I.

4. *S. Géum* (L.); l. transversely oval or reniform crenate or dentate, footstalks semicylindrical with raised edges hairy, panicle racemose.—*E. B. S.* 2893.—L. always broader than long, usually hairy, sometimes glabrous, often beautifully netted with purple beneath.—West of Ireland. P. VI. I.

[*S. Andrews'ii* (Harv.); l. linear-spathulate dentate-serrate with a membranous margin tapering below into a dilated plane-convex footstalk, panicle racemose.—*Sy. E. B.* 549. *R. Guthriana* Hort.—Sep. patent, oblong, blunt. Pet. thrice as long as cal., oblong, punctate.—Said to have been found at head of Glen Carragh, Kerry. A garden hybrid? P. VI.]

†† Filaments subulate. SPATULARIA Haw.

5. *S. stelláris* (L.); l. oblong wedgeshaped dentate-serrate narrow below scarcely stalked, panicle corymbose of few flowers.—*E. B.* 167. *St.* 35. 3.—Stems tufted. Fl.-stalks 1—5 in. high, naked. Pet. ovate, clawed, white with 2 transverse yellow spots on their lower half.—L. rarely quite entire.—Damp rocks on mountains. P. VII. E. S. I.

\*\* *Cal. at length reflexed inferior, st. leafy, l. undivided.*

6. *S. Hir'culus* (L.); st. erect, barren shoots prostrate filiform, l. alternate lanceolate flat entire, root-l. narrowed into a footstalk, sep. blunt fringed at the margin, pet. blunt with 2 hard points near the base.—*E. B.* 1009. *St.* 35. 8.—Fl. few or solitary, terminal. Pet. obovate, spreading, yellow dotted with red. *St.* 4—8 in. high, downy in the upper part.—Wet moors, rare. P. VIII. E. S. I.

\*\*\* *Cal. erect or spreading half inferior, st. leafy, l. simple toothed or with rigid jointless cilia, also a transparent gland at the tip.*

7. *S. aizoi'des* (L.); st. decumbent below, l. alternate linear-oblong mucronate ciliate entire flat above convex beneath, sep. blunt.—*E. B.* 39. *St.* 35. 9.—Fl. in a leafy panicle, with glu-

tinous downy stalks. Pet. bright yellow often spotted with scarlet. St. 3—6 in. long. L. rigidly ciliate or with several strong teeth.—Wet places on mountains. P. VII.—IX. E. S. I.

\*\*\*\* *Cal. erect or spreading superior or half inferior, st. more or less leafy, l. lobed, cilia all jointed.*—See Baker in *J. of B.* viii. 280, 355.

8. *S. cæspitosa* (L.); root-l. crowded 3—5-cleft blunt veined fringed, fl. 1—5, pet. rounded 3-veined, *cal. half inferior*, sep. blunt.—Barren st. very short. Fl. white, Germen broad and rounded below. Common base of the l. not furrowed. Anth. long, cordate with an open notch. Horns of caps. spreading.—*β. S. incurvifolia* (D. Don); stem-l. more numerous with incurved lobes. *E. B. S.* 2909.—Very rare. Caernarvonshire. Aberdeenshire. Kerry. P. V. VI. E. S.

9. *S. Sternbergii* (Willd.); l. crowded 3—4-cleft, lobes lanceolate pointed fringed, fl. few (usually 3) loosely paniced, *pet. obovate* 3-veined, *cal. half inferior*, *sep. bluntish*.—*S. hirta* Sm. *E. B.* 2291 (not Haw.).—L. on the barren shoots as well as the radical l. 3-cleft, the lateral lobes often divided halfway down, lobes diverging. Barren shoots 3—6 or 8 in. long, weak. Fl. st. hairy glandular, bearing 2 or 3 deeply 3-lobed l. and a few simple linear bracts; lateral fr.-ped. much overtopping the intermediate one. Fl. white. Anth. large, round, with a deep parallel-sided notch. Horns of caps. slightly spreading.—*S. decipiens* (Sm.) is hardly distinguishable, it is found at Llyn y Cwn, N. Wales.—Summit of Brandon Mountain, Kerry; Galty More, Tipperary. West of Scotland. *Mr. G. Don.* P. VII. E. S. I.

10. *S. sponhem'ica* (Gm.); l. of the barren shoots 3—5-lobed with a very broad base fringed, lobes linear acute, fl. few (2—4), *pet. oblong* 3-veined, *cal. half inferior* deeply divided into triangular *subulate sepals*.—*S. affinis* Don. *E. B. S.* 2903. *S. pinnatifida* Haw.—The 5-cleft l. few. St. 1—3 in. high, erect, with a few linear simple leaves, glandular. Fl. white. Anth.  $\frac{1}{2}$  as large as in Sp. 8 and 9, cordate-ovate. Horns of caps. spreading.—The original *S. affinis* has pet. with inflexed sides: *S. platypetala* (*E. B.* 2276) has them broad and flat.—Mountains. P. VII. E. S. I.

11. *S. hypnoï'des* (L.); root-l. 3—5-cleft those of the trailing shoots undivided or 3-cleft, lobes all acute bristle-pointed and fringed, calyx half inferior, *sep. triangular-subulate* or *ovate acute*.—*E. B.* 454.—Fl. white. Anth. very small, oblong-ovate. Horns of caps. divaricate. Differs from the preceding by having usually buds in the axils of the barren shoots.—Mountains. P. V.—VII. E. S. I.

[*S. pedatifida* (Sm.).—*E. B.* 2278.—Probably of garden origin.]

*Sec. 2.* Without barren shoots at the base.

\* *Stem leafy.*

12. *S. tridactylites* (L.); *st.* panicled erect leafy, l. wedge-shaped 3—5-fid with a flat petiole, lowermost l. often simple and spatulate, *peduncles* 1-flowered much longer than the fruit with 2 bracts at the base, cal. superior.—*E. B.* 501. *St.* 33. 15.—Whole plant viscid, 2—4 in. high. Fl. scattered, many, small, white.—Walls and dry banks. A. IV.—VII.  
E. S. I.

13. *S. granulata* (L.); *st.* erect slightly leafy, radical l. reniform crenately lobed with channelled petioles, stem-l. nearly sessile 3—5-fid, fl. in a cymose panicle, cal. half inferior, pet. obovate-oblong 2 or 3 times as long as the sepals, roots bearing many small round downy bulbs.—*E. B.* 500.—*St.* 6—12 in. high. Fl. large, white.—Gravelly banks. P. V.  
E. S. I.

14. *S. cer'nea* (L.); *st.* erect simple 1-flowered leafy, radical l. reniform palmately lobed stalked, upper l. nearly sessile subtrifid, uppermost entire, axils bearing bulbs, cal. quite inferior.—*E. B.* 664.—Rarely flowering; fl. replaced by reddish bulbs. *St.* 3—6 in. high.—Rocks on the top of Ben Lawers, very rare. P. VI.—VIII.  
S.

15. *S. rivularis* (L.); *st.* ascending branched few-flowered leafy, radical l. subreniform stalked with 3—5 rounded lobes, uppermost l. lanceolate entire, cal. half inferior.—*E. B.* 2275.—*St.* 1—2 in. long. Fl. few, stalked, white.—Wet places on the coldest parts of mountains. P. VIII.  
S.

\*\* *Stem leafless.*

16. *S. nivalis* (L.); *st.* erect leafless, l. all radical roundish-obovate dentate-serrate narrowed into a footstalk, fl. in capitate cymes, cal. half inferior, pet. longer than the calyx.—*E. B.* 440. *St.* 35. 4.—*St.* 3—6 in. high, usually simple, sometimes with 1 branch. Fl. in a dense cluster, white.—Alpine rocks. P. VII.  
E. S. I.

*Sec. 3.* Stems procumbent with opposite l. and terminal flowers.

17. *S. oppositifolia* (L.); *st.* procumbent, l. opposite in 4 rows oblong blunt fringed, sep. ciliate without glands, pet. ovate.—*E. B.* 9.—Very different from our other species. L. with a pore at tip. Fl. large, purple.—Damp alpine rocks. P. IV. V.  
E. S. I.

## 2. CHRYSOSPLE'NIUM Linn. Golden Saxifrage.

1. *C. alternifolium* (L.); *l. alternate*, lower l. subreniform hairy crenate upon long stalks.—*Sy. E. B.* 564. *St.* 12.—Crenatures of the lower l. emarginate, upper l. glabrous with the crenatures often rather acute. *St.* erect, 4—5 in. high, *branching only near the top*. Fl. umbellate, nearly sessile, deep yellow. Stam. usually 8.—Boggy places. P. IV. E. S. I.

2. *C. oppositifolium* (L.); *l. opposite*, lower l. roundish-cordate shortly stalked wavy.—*E. B.* 450. *St.* 4. 6.—*St. branching from the base*, 4—6 in. long, decumbent, straggling, rooting. Fl. paler and more scattered than in the last. L. usually glabrous, sometimes slightly hairy. Stam. usually 8.—Damp shady places. P. IV. V. E. S. I.

## 3. PARNAS'SIA Linn.

1. *P. palustris* (L.); filaments of the petal-like scales 9—13, pet. with a short claw, radical l. cordate stalked, st.-l. clasping.—*E. B.* 82.—Pet. white, veined. Glands of the scales yellow. L. mostly radical. *St.* 8—10 in. high.—Wet and boggy ground towards the north. P. VIII.—X. E. S. I.

B. Petals and stamens epigynous, inserted round an epigynous disk. Cal.-tube adnate to the ovary.

## Order XXXVII. UMBELLIFERÆ.

Cal. 5-toothed or entire, adherent to the ovary, limb often scarcely visible. Pet. 5; usually inflexed at the point. Stam. 5, inserted with the pet. round the stylopode. Ovary 2-celled, crowned with a double fleshy disk (stylopode). Styles 2. Fr. consisting of 2 carpels (mericarps) adhering by their face (commissure) to a common axis from which they ultimately separate and become pendulous. Seed solitary, pendulous. Albumen horny.—Inflorescence umbellate. *Æstivation imbricate*, except in *Hydrocotyle* and *Criethnum*.—Each carpel has 5 primary, and often 4 intermediate secondary ridges; and in the substance of the pericarp are usually linear receptacles of oil (*vittæ* or stripes) under the ridges or the spaces between them. These parts are sometimes either wanting or only slightly apparent. The stripes are "solitary" when there is only one in each space between the primary ridges, and "2, 3, &c. together" when 2, 3, or more occur in each space. They and the ridges are best seen by making a horizontal section of the fruit.

## Suborder I. ORTHOSPERMÆ.

Seed or albumen straight or nearly so. Umbels various.

\* *Umbels imperfect or simple; no fr.-stripes. "No carpophore."*

Tribe I. *HYDROCOTYLEÆ*. Fr. laterally flattened, its back even or acute. Umbel irregular or imperfect.

1. *HYDROCOTYLE*. Cal.<sup>1</sup> inconspicuous. Pet. ovate, entire. Fr. of 2 flat nearly orbicular carpels, each with 5 filiform ridges, of which the dorsal and 2 lateral are often inconspicuous; the 2 others arched. Stripes 0. Commissure linear.

Tr. II. *SANICULEÆ*. Fr. ovoid; transverse section nearly round. Umbel simple or imperfect.

2. *SANICULA*. Cal. of 5 leaflike teeth. Pet. erect, obovate, with a long inflexed connivent point. Fr. subglobose, covered with hooked spines; ridges 0; stripes many.
3. *ASTRANTIA*. Cal. of 5 leaflike teeth. Pet. erect, with a long inflexed point. Fr. with 5 plaited dentate ridges; stripes 0.

Tr. III. *ERYNGIÆ*. Fr. ovoid; transverse section nearly round. Fl. in a head-like umbel.

4. *ERYNGIUM*. Cal. of 5 leaflike teeth. Pet. erect, oblong, with a long inflexed point. Fr. covered with chaffy scales, without ridges or stripes.

\*\* *Umbels perfect or compound.*

† Carpels with 5 primary ridges only.

Tr. IV. *AMMINEÆ*. Fr. of two pentagonal carpels with 5 prominent ridges; the commissure about as broad as either of the other four sides.

A. *Leaves compound.*

5. *CICUTA*. Cal. of 5 leaflike teeth. Pet. obcordate with an inflexed point. Fr. subdidymous. Carp. with 5 equal broad flattened ridges, the lateral marginal; stripes solitary.
6. *APIUM*. Cal. inconspicuous. Pet. roundish entire with an oblique acute or small involute point. Fr. roundish ovoid, didymous. Carp. with 5 filiform equal prominent ridges and solitary stripes.—*Inv.* 0.
7. *PETROSELINUM*. Cal. minute or inconspicuous. Pet. roundish entire with a narrow incurved point. Fr. ovoid. Carp. with 5 filiform equal ridges and solitary stripes. Car-

<sup>1</sup> By calyx, throughout this Order, the free margin is intended.

pophore bipartite.—Partial inv. of many; general of few leaves.

8. SISON. Cal. inconspicuous. *Pet. broadly obcordate, deeply notched* with an inflexed point. Fr. ovoid. Carp. with 5 filiform prominent equal ridges and *solitary clavate stripes*.
9. TRINIA. Dioecious. Cal. inconspicuous. *Pet. of the barren fl. lanceolate* with the point inflexed, *of the fertile fl. ovate* with a short inflexed point. Fr. ovoid. Carp. with 5 filiform prominent equal ridges *with a single stripe beneath each of them, but none between*.
10. ÆGOPIDIUM. Cal. inconspicuous. *Pet. obovate, notched*, with an inflexed point. Fr. oblong. Carp. with 5 filiform ridges; the *interstices without stripes*. Stylopode conical.
11. CARUM. Cal. inconspicuous. *Pet. obcordate with an inflexed point*. Fr. oblong. Carpels with 5 filiform ridges; the stripes solitary or 1—3 together. *Stylopode depressed*.
12. PIMPINELLA. Cal. inconspicuous. *Pet. obcordate with an inflexed point*. Fr. ovoid. Carp. with 5 filiform equal ridges; *stripes 3 or more together*. *Stylopode tumid*. Styles of the fr. divaricate or recurved.—*Inv. 0* (rarely of 1 leaf).
13. SIUM. Cal. of 5 small teeth. *Pet. obcordate with an inflexed point*. Fr. ovoid or subdidymous. *Carp. with 5 filiform equal blunt ridges*; stripes 3 or more together. *Stylopode depressed or shortly conical*. Styles of the fr. divaricate or recurved.—*Inv. general and partial*.

B. *Leaves simple.*

14. BUPLEURUM. Cal. inconspicuous. *Pet. roundish entire with a closely involute broad retuse point*. Fr. subdidymous. Carp. with equal, winged, or filiform and sharp, or inconspicuous ridges. *Stylopode depressed*.
- Tr. V. SESELINEÆ. Fr. of two 5-ribbed or pentagonal carpels. Commissure much the broadest side of the carpel.

A. *Stripes solitary, between the ribs.*

15. CENANTHE. Cal. of 5 lanceolate teeth. *Pet. obcordate with an inflexed point*. Fr. ovoid cylindrical or subturbinate, crowned with the long *suberect styles*. Carpels more or less corky, with 5 blunt convex ridges.
16. ÆTHUSA. Cal. inconspicuous. *Pet. obcordate with an acute inflexed point*. Fr. shortly ovoid, crowned with the *reflexed styles*. Carp. with 5 thick acutely keeled ridges.

17. *FÆNICULUM*. Cal. *inconspicuous*. Pet. *roundish entire with a broad blunt inflexed lobe*. Fr. *oblong*. Carp. with 5 prominent bluntly keeled ridges. Stylopode *conical*.
18. *SESELI*. Cal. *with acute teeth*. Pet. *obcordate with a broad inflexed lobe*. Fr. *ovoid or oblong, slightly dorsally compressed*. Carp. with 5 thick blunt ridges. Stylopodes *conical*.

B. *Stripes 2 or more together, between the ribs.*

19. *HALOSCIAS*. Cal. of 5 small persistent teeth. Pet. *ovate with an inflexed lobe and short claw*. Fr. *elliptic, terete or slightly dorsally compressed*. Carp. *with 5 sharp somewhat winged ridges*. Interstices and commissure with many stripes. *Seeds not adhering to the carpel, without stripes*.
20. *SILAUS*. Cal. *inconspicuous*. Pet. *ovate-oblong entire or slightly emarginate with an inflexed lobe, sessile, truncate or appendaged at the base*. Fr. *oblong, terete or slightly dorsally compressed*. Carp. with 5 sharp somewhat winged ridges. Stripes many, *inconspicuous*.
21. *MEUM*. Cal. *inconspicuous*. Pet. *entire, elliptic, acute at both ends, with an incurved point*. Fr. as in *Silaus*.
22. *CRITHMUM*. Cal. *inconspicuous*. Pet. *elliptic with a broad base, entire, involute*. Fr. *oblong, terete*. Carp. with 5 elevated sharp slightly winged ridges. *Seed free; with many stripes*.
- Tr. VI. *ANGELICEÆ*. Fr. of two much and dorsally compressed carpels, with a double wing on each side.
23. *ANGELICA*. Cal. *inconspicuous*. Pet. *lanceolate, entire, acuminate, incurved*. Carp. *with 3 dorsal elevated filiform ridges and 2 marginal ridges dilated into broad wings; interstices with solitary stripes. Seed adhering to the carpel*.
- [24. *ARCHANGELICA*. Cal. *minutely 5-toothed*. Pet. *elliptic, entire, acuminate, incurved*. Carp. *with 3 dorsal elevated thick ridges and two marginal ridges dilated into broad wings. Stripes 0. Seed free, with many fine lines.*]
- Tr. VII. *PEUCEDANEÆ*. Fr. of two much and dorsally compressed carpels, with a single wing on each side; wing flat or thickened towards the edge, formed of those of the two carpels combined.
25. *PEUCEDANUM*. Cal. of 5 teeth or *inconspicuous*. Pet. *obovate or obcordate with an inflexed point*. Fr. with a dilated thin flat margin. Carp. with equidistant ridges, 3



dorsal filiform, 2 lateral close to the base of the dilated margin inconspicuous; stripes solitary.

26. PASTINACA. Cal. of 5 very small or inconspicuous teeth. *Pet. roundish, entire, involute with an acute point. Fr. with a dilated flat margin. Carp. with slender ridges, 3 dorsal equidistant, 2 lateral distant near the outer edge of the dilated margin; stripes linear, solitary.*—Fl. yellow.
27. HERACLEUM. Cal. of 5 minute teeth. *Pet. obcordate with an inflexed point, outer ones radiant. Fr. as in Pastinaca, but the stripes short clubshaped.*—Fl. white.
28. TORDYLIUM. Cal. of 5 awlshaped teeth. *Pet. obcordate with an inflexed lobe, outer ones radiant. Fr. with a thickened wrinkled margin. Carp. with slender ridges, 3 dorsal equidistant, 2 lateral distant close to the thickened margin; stripes 1—3 together.*

†† Carpels with primary and secondary ridges.

[Tr. VIII. SILERINÆ. Fr. dorsally compressed. Carp. with 5 primary ridges, the lateral marginal, and 4 less prominent secondary ridges.]

[29. SILER. Cal. of 5 teeth. *Pet. obovate with an inflexed lobe, regular. Fr. dorsally compressed. Carp. with elevated filiform blunt ridges; one stripe under each secondary ridge and 2 on the commissure.*]

Tr. IX. DAUCINÆ. Fr. somewhat dorsally compressed. Carp. with 5 primary ridges, the lateral ones on the inner face; and 4 secondary, forming rows of prickles.

30. DAUCUS. Cal. of 5 teeth. *Pet. obcordate with an inflexed lobe, exterior usually radiant and bifid. Fr. dorsally compressed. Carp. with bristly primary ridges; secondary ridges equal winged with 1 row of spines.*

#### Suborder II. CAMPYLOSPERMÆ.

Sutural part of seed inflexed at the sides or deeply furrowed lengthwise. Umbels compound or perfect.

Tr. X. CAUCALINÆ. Fr. contracted or rounded. Carp. with the lateral primary ridges on the inner face; 4 secondary more prominent, prickly; stripes solitary.

31. CAUCALIS. Cal. of 5 teeth. *Pet. obcordate with an inflexed point, outer ones radiant and bifid. Fr. slightly laterally compressed. Carp. with filiform bristly primary, and more or less prominent secondary ridges, all bearing 1—3 rows of prickles.*—Umbels sometimes simple.

32. TORILIS. Cal. of 5 teeth. Pet. obcordate with an inflexed point, outer ones radiant and bifid. Fr. slightly laterally compressed. *Carp. with* bristly primary ridges, with *many prickles on the spaces* between them.
- Tr. XI. SCANDICINEÆ. Fr. compressed or contracted on the sides, often beaked or narrowed at the top. *Carpels with primary ridges only.*
33. SCANDIX. Cal. inconspicuous. Pet. obovate with an inflexed point. *Fr. with a very long beak. Carp. with 5 blunt ridges*; stripes none.
34. CHÆROPHYLLUM. Cal. inconspicuous. Pet. obcordate with an inflexed point. *Fr. hardly beaked. Carp. with 5 equal blunt ridges*, often only apparent at the top; stripes solitary or wanting.
35. MYRRHIS. Cal. inconspicuous. Pet. obcordate with an inflexed point. Fr. not beaked. *Carp. formed of a double membrane*; the outer with elevated keeled ridges hollow within, the inner close to the seed; stripes 0.
- Tr. XII. SMYRNIÆ. Fr. ovoid or didymous compressed or contracted on the sides. Carpels with primary ridges only, not beaked nor remarkably narrowed at the top.
36. ECHINOPHORA. Cal. of 5 teeth. Pet. obcordate with an inflexed point, the exterior larger and bifid. Fl. of the ray, sterile on long stalks, fertile central and solitary. Fr. ovoid terete, imbedded in the enlarged prickly receptacle. *Carp. with 5 depressed equal striate wavy ridges*; interstices with single stripes and covered by a cobweblike membrane.
37. CONIUM. Cal. inconspicuous. Pet. obcordate with a short inflexed point. Fr. ovoid, laterally compressed. *Carp. with 5 prominent or wavy crenate ridges*, the lateral marginal; interstices striate; stripes 0\*.
38. PHYSOSPERMUM. Cal. of 5 teeth. Pet. obcordate with an inflexed point. Fr. laterally compressed, didymous. *Carp. reniform-globose, with 5 filiform slender equal ridges, the lateral within the margin*; stripes solitary.
39. SMYRNIUM. Cal. inconspicuous. Pet. lanceolate or elliptic, entire, with an inflexed point. Fr. laterally compressed, didymous. *Carp. reniform-oblong with 3 dorsal prominent sharp ridges, the 2 lateral marginal and inconspicuous*; stripes many.

\* I have never seen any stripes, which are said to be sometimes present.

## Suborder III. CŒLOSPERMÆ.

Seed with the base and apex curved inwards.

Tr. XIII. *CORIANDRÆÆ*. Fr. globose or didymous. Primary ridges of the carpels often inconspicuous, secondary more prominent, all without wings.

[40. *CORIANDRUM*. Cal. of 5 teeth. Pet. obcordate with an inflexed point, outer ones radiant and bifid. Fr. globose. Carp. scarcely separating, the primary ridges inconspicuous, the 4 secondary conspicuous prominent keeled; interstices without stripes; commissure with 2 stripes.]

Suborder I. *Orthospermæ*. Tribe I. *Hydrocotyleæ*.1. *HYDROCOT'YLE* Linn. Pennywort.

1. *H. vulgâris* (L.); l. peltate nearly circular 9-veined doubly crenate, petioles pilose, umbels 3—6-fl., fr. emarginate below.—*E. B.* 751.—Fl. and fr. almost sessile. Umbels or rather heads often proliferous in the centre and bearing a second head. St. creeping extensively. L. upon stalks which considerably exceed the peduncles. Ripe fr. with purplish dots.—Bogs and marshy places. P. V.—VII. E. S. I.

Tribe II. *Saniculeæ*.2. *SANIC'ULA* Linn. Sanicle.

1. *S. europæ'a* (L.); lower l. palmate 3—5-lobed, lobes trifid unequally serrate, fertile fl. sessile, barren fl. slightly stalked.—*E. B.* 98.—Umbels many, capitate, in an irregular slightly umbellate panicle. Styles persistent, reflexed. Fr. covered with hooked prickles. St. ascending about a foot high.—Woods and thickets. P. VI, VII. E. S. I.

3. *ASTRAN'TIA* Linn.

†1. *A. májor* (L.); lower l. palmately 5—7-fid, lobes oblong acutely unequally incise-serrate, inv.-l. entire, cal.-teeth ovate-lanceolate narrowed to an acute point.—*E. B. S.* 2990. St. 29. 8.—Inv. equalling the umbel, usually straw-coloured. Cal.-teeth exceeding petals.—Woods in hilly districts. Above Stokesay Castle, near Ludlow, where Mr. Borrer considered it to have been introduced “ages ago.” P. VI.—VIII. E.

Tribe III. *Eryngiæ*.4. ERYNGIUM *Linn.* Eryngo.

1. *E. maritimum* (L.); radical l. roundish plaited spinous stalked, upper l. amplexicaul palmately lobed, *inv.-l.* 3-lobed spinous exceeding the heads, scales of the receptacle 3-lobed.—*E. B.* 718.—St. 1 foot or more in height, much branched, leafy, rigid, glaucous. Fl. in heads rather than umbels, blue.—Sandy sea-shores. P. VII. VIII. *Sea-Holly*. E. S. I.

2. *E. campestre* (L.); radical l. 2 or 3 times pinnatifid spinous stalked, st.-l. amplexicaul pinnatifid, *inv.-l.* lanceolate spinous longer than the heads, scales of the receptacle undivided.—*E. B.* 57.—More bushy and slender than the last. Pet. purplish or white. Petioles thick, semiterete, channelled.—On waste ground, very rare. P. VII. VIII. E.

Tribe IV. *Ammineæ*.5. CICUTA *Linn.* Water-Hemlock.

1. *C. virôsa* (L.); fibres of the root filiform, l. tripartite, leaflets linear-lanceolate acute serrate decurrent.—*E. B.* 479.—St. 3—4 feet high, very thick terete and hollow below. Lower l. on long stalks; leaflets 1—2 in. long. Umbels large; general *inv.* 0 or of 1 or 2 slender leaflets, partial of many subulate leaflets. Fl. white. Herb poisonous.—Ponds and ditches, not common. P. VII. VIII. *Cowbane*. E. S. I.

6. A'PIUM *Linn.*

\* *Pet. with involute point, cordate base, minute claw. Partial inv. 0.*

1. *A. graveolens* (L.); glabrous, l. pinnate or ternate, leaflets of the upper l. wedged-shaped.—*E. B.* 1210.—St. 1—2 feet high, branched, furrowed, leafy. Umbels terminal or lateral, frequently almost sessile, accompanied by one or two ternate leaves. Fl. small, whitish.—Marshes and ditches, especially near the sea. P. VI.—VIII. *Celery*. E. S. I.

\*\* *Pet. with oblique acute point, and wedged-shaped base. Partial inv. present.*—HELOSCIADIUM Koch.

2. *A. nodiflorum* (R.); st. procumbent at the base and rooting, l. pinnate, leaflets ovate or ovate-lanceolate unequally bluntly serrate, umbels opposite to the l. longer than their pe-

duncles or nearly sessile.—*E. B.* 639.—St. 1—2 feet long.—*a*; general inv. soon falling of 1—3 lts., anth. purple.—*β. A. repens* (R.); creeping, lts. unequally and acutely incise-serrate, umbels shorter than the peduncles, anthers yellow.—*γ. ocreatum*; creeping, l. roundish ovate small bluntly toothed, umbels very shortly stalked.—*E. B.* 1431.—Banks of ditches and brooks. P. VII. VIII. E. S. I.

3. *A. inundatum* (R.); st. creeping or floating, l. pinnate, lts. of lower l. in capillary segments, of upper l. wedgeshaped and trifid, umbels generally with 2 rays.—*E. B.* 227.—Usually submersed, a few of the upper l. and the fl. rising above the water. Partial umbels very small.—Ponds. P. VI. VII. E. S. I.

### 7. PETROSELINUM Hoffm.

\*1. *P. sativum* (Hoffm.); l. tripinnate shining, leaflets of the lower l. ovate-cuneate trifid and toothed of the upper l. ternate lanceolate entire.—*E. B. S.* 2793.—Partial inv. filiform. Fl. yellow. L. greenish.—Rocks and old walls. B. VI.—VIII. Parsley. E.

2. *P. seg'etum* (Koch); lower l. pinnate, leaflets nearly sessile ovate lobed and serrate, upper l. entire or trifid.—*Sison* Sm. *E. B.* 228.—Umbels very irregular. General inv. of 1—2 leaves. Fl. whitish. St. erect, terete, nearly leafless above, 1—1½ foot high, wiry.—Damp calcareous fields and near the sea. B. VIII. IX. E.

### 8. SISON Linn. Stonewort.

1. *S. Amómum* (L.).—*E. B.* 954.—St. erect, panicled, 2—3 feet high. Lower l. pinnate; lts. oblong lobed cut and serrate. Upper l. divided into narrow segments. Partial umbels and fl. small. Much like *Petr. segetum*.—Dampish places on a calcareous soil. B. VIII. E.

### 9. TRI'NIA Hoffm. Honewort.

1. *T. vulgáris* (DC.); glabrous, inv. 0 or of 1 leaf, ridges of the fr. blunt.—*Pimpinella dioica* Sm. *E. B.* 1209, *Sy. E. B.* 579.—L. tripinnate, glaucous green; lts. linear or filiform. Root crowned with the remnants of former leaves. St. branched, erect, 6—8 in high. Dioecious.—Dry limestone hills, rare. P. V. VI. E.

### 10. ÆGOPO'DIUM Linn. Gout-weed.

1. *Æ. Podagrária* (L.).—*E. B.* 940.—St. 1—2 feet high, erect, furrowed. L. 2 or 3 times ternate; leaflets ovate-acumi-

nate, unequal at the base, acutely serrate. Creeping.—Damp places. P. VI. VII. E. S. I.

## 11. CA'RUM Linn.

\* *Root fusiform or fascicled. Stripes 1—2 in each interstice.*

[*C. Car'ui* (L.); *partial involucre* 0, general 0 or of 1 leaf, l. bipinnate, leaflets cut into linear segments.—*E. B.* 1503.—St. 1—2 feet high, branched. Root fusiform. Carp. aromatic.—Meadows and pastures. B. VI. Caraway.] E. S.

1. *C. verticillatum* (Koch); *general and partial involucre*s of many leaves small, l. pinnate, leaflets divided to the base into capillary spreading segments.—*E. B.* 395.—St. 1—1½ foot high. Root fasciculate. *Segments of the leaflets* spreading so as to appear whorled and quite surrounding the petiole. L. mostly long-stalked, radical.—Marshy places in hilly districts, rare. P. VIII. E. S. I.

\*\* *Root a tuber.*—BUNIUM Linn.

† Stripes 3 in each interstice.

2. *C. flexuosum* (Fr.); *general involucre* of 1—3 leaves, partial more numerous, *fr. oval* narrowing upwards crowned with the long stylopode and erect styles.—*E. B.* 988. *B. denudatum* DC.—*Involucre*s sometimes altogether wanting. Root a solitary tuber. St. a foot or more high, very slender below, bearing a few l. with linear segments. Radical l. triternate with long footstalks tapering downwards.—Sandy and gravelly pastures. P. V. VI. Pig-nut. E. S. I.

†† Stripes 1—2 in each interstice.

3. *C. Bulbocastanum* (Koch); *general and partial involucre*s of many leaves, *fr. oblong* crowned with the short stylopode and reflexed styles, interstices with single stripes.—*E. B.* S. 2862.—*Involucre*s always present. Root a solitary tuber. St. about 2 ft. high. Lower l. bipinnate, with a triangular outline, rather many near the base of the stem, rarely 1 or 2 radical upon long footstalks tapering downwards.—Chalky fields in Camb., Bucks., Beds., and Herts. P. VI. VII. E.

## 12. PIMPINEL'LA Linn. Burnet-Saxifrage.

1. *P. mag'na* (L.); l. pinnate, leaflets all ovate serrate somewhat cut the terminal one 3-lobed, *st. angularly striate*.—*E. B.* 408.—St. 1—2 feet high, leafy. Lateral leaflets sometimes 3-lobed. Styles longer than the ovary, as long as or longer than the oval fruit.—β. *P. dissecta* (Retz); l. all divided into long linear cut segments.—Shady hills. P. VII. VIII. E. S. I.

2. *P. Saxifraga* (L.); l. pinnate, leaflets of the lower l. roundish-ovate serrate somewhat cut, *those of the stem-l. bipinnatifid with linear segments, st. terete*, ped. glabrous.—*E. B.* 407.—St. 1—2 feet high, naked above. Styles shorter than the ovary. L. sometimes all pinnatifid.—Dry pastures. P. VII.—IX. E. S. I.

### 13. SI'UM Linn.

\* *Stripes superficial. Lateral ridges marginal. Stylopode depressed.* SIUM Koch.

1. *S. latifolium* (L.); l. pinnate, lts. oblong-lanceolate *evenly serrate* pointed, umbels terminal, inv.-l. many lanceolate.—*E. B.* 204.—St. 3—5 feet high, angular, furrowed, erect. L. of 9—13 very largedistant lts.—Ditches and rivers, rare. P. VII. VIII. E. S. I.

\*\* *Stripes deeply seated. Lateral ridges not marginal. Stylopode shortly conical.* BERULA Koch.

2. *S. angustifolium* (L.); l. pinnate, *leaflets unequally lobed and cut* ovate, of the stem-l. lanceolate, umbels lateral, inv. of many lanceolate entire or cut leaves.—*E. B.* 139.—St. 1—3 feet high, round, striate, erect. Lts. often very deeply cut and lobed.—Ditches. P. VIII. E. S. I.

### 14. BUPLEU'RUM Linn. Hare's-ear.

\* *Fruit granular.*

1. *B. tenuis'simum* (L.); st. branched, l. linear acute, umbels lateral and terminal minute, partial umbels of 3—5 flowers usually overtopped by their involucre, carp. granular between the 5 ridges.—*E. B.* 478.—St. very slender, wiry, 6—12 in. long.—Chiefly in pastures near the sea. A. VIII. IX. E.

\*\* *Fruit not granular.*

2. *B. aristatum* (Bartl.); st. branched, l. linear-lanceolate acuminate 3-veined, l. of the partial involucre elliptic-lanceolate cuspidate somewhat awned with branching veins, pedicels short equal.—*E. B.* 2468.—St. 1—6 in. high.—Park Hill, Torquay, and Berry Head, Devon. Cow Gap near Eastbourne, Suss. Channel Islands. A. VI. E.

3. *B. falcatum* (L.); st. branched, l. 5—7-veined, *lower l. elliptic-oblong* on long stalks, *upper l. linear-lanceolate* acute sessile, partial involucre of 5 lanceolate pointed l. as long as the flowers.—*E. B. S.* 2763.—Pedicels as long as the fruit. St. 1—4 feet high.—Near Ongar, Essex. P. VIII. E.

4. *B. rotundifolium* (L.); st. branched above, l. oval perfoliate, fr. with striate interstices.—*E. B.* 99.—St. 12—18 in. high. General inv. 0. Partial inv. connivent.—Corn-fields on a calcareous soil. A. VII. *Thorough-wax.* E.

### Tribe V. *Seselineæ*.

#### 15. CENANTHE Linn. Water-Dropwort.

\* *Root fascicled, fibres more or less thickened or tuberous.*

1. *C. fistulosa* (L.); stoloniferous; st. and petioles hollow, root-l. 2—3-pinnate with 3-fid leaflets, stem-l. simply pinnate shorter than their petioles, leaflets linear, fr. angular turbinate.—*E. B.* 363.—St. 1—3 feet high, remarkably hollow. Stem-l. distant, with very long stalks. Stoles with simply pinnate leaves. Umbels small, globose in fruit; general involucre 0. Fruit tipped with the long slightly diverging rigid styles.—By ponds and ditches. P. VII.—IX. E. S. I.

2. *C. pimpinelloides* (L.); root of long fibres bearing round or ovoid knobs beyond their middle, root-l. bipinnate with obovate-wedged-shaped 3-lobed lts., st.-l. pinnate with linear acute lts., uppermost l. simple, radiant pet. obcordate divided to the middle, fr. subcylindrical with an enlarged corky base but not narrowed at the top.—*E. B. S.* 2991. *Jacq. Aust.* t. 394.—St.  $\frac{1}{2}$ —3 ft. high, alternately branched. General inv. 0—6-leaved; partial of many l., about as long as the barren fl. *Partial umbels close together forming one compact flat-topped compound umbel.* Fr. nearly cylindrical; cal. erect-patent.—Southern counties. P. VI. VII. E.

3. *C. Lachenalii* (Gmel.); root of long subclavate fleshy fibres tapering at both ends, root-l. bipinnate with oblong entire or wedged-shaped and bluntly 2—3-lobed lts., lower st.-l. 2—3-pinnate with linear acute lts., upper l. simply pinnate, radiant pet. divided to the middle, fr. oblong not corky below rounded and contracted at the top.—*C. pimpinelloides* Sm. *E. B.* 347.—St. 1—3 feet high, slightly branched. General inv. of many leaves, sometimes wanting; partial of many leaves, shorter than the barren flowers. Outer fl. on long stalks, mostly barren; inner fl. fertile, nearly sessile. Radiant pet. roundish obcordate with a short narrow claw. *Partial umbels distinct, spherical.* Fr. crowned with the inflexed calyx. Root-leaves soon vanishing.—Marshes. P. VII.—IX. E. S. I.

4. *C. silaifolia* (Bieb. ?); root of elliptic-oblong knobs, radical l. bipinnate, st.-l. pinnate, lts. all linear acute, external fr. nearly



cylindrical with a corky base, but not narrowed at the top.—*E. B.* 348.—St. 2—3 feet high, branched. Outer fl. stalked, mostly barren; cal. very unequal; radiant pet. small, obcordate with an attenuate base, notch  $\frac{1}{3}$  their length. Fr. usually slightly narrower downwards, in the middle of the umbel much narrower and appearing to want the corky base; cal. erect or inflexed. General involucre 0; partial of many leaves shorter than the flowers. Distinguished from the preceding by its pet., uniform leaflets, want of a general involucre, and fruit.—The true name of this plant is very doubtful.—In freshwater marshes. P. VI. E.

5. *Æ. crocata* (L.); root of large fusiform tubers, radical l. 2—3-pinnate, stem-l. pinnatifid, leaflets stalked roundish or oblong-wedged shaped variously cut those of the upper l. narrower, fr. cylindrical oblong striate longer than its pedicel.—*E. B.* 2313.—Poisonous. St. 3—5 feet high, much branched. L. large, lts. broad. Inv.-l. various in number and shape.—Wet places. P. VII. E. S. I.

\*\* *Root of whorls of slender fibres.* PHELLANDRIUM L.

6. *Æ. Phellan'drium* (Lam.); st. erect fusiform below, l. tripinnate, lts. ovate pinnatifid cut spreading, of the submersed l. multifid with capillary diverging segments, umbels lateral opposite to the leaves, fr. ovate.—*E. B.* 684.—St. 2—3 feet high, very thick below, stoloniferous. Segments of the l. many, fine, acute, pale green; submersed l. dark green. The flowering root dies each year; but the plant is continued by the offsets.—In wet ditches and ponds. B. ? VII.—IX. *Horsebane.* E. S. I.

7. *Æ. fluviat'ilis* (Colem.); st. floating, l. bipinnate, lts. simple or pinnatifid, of the submersed l. cuneate cut pellucid with many parallel veins, umbels lateral opposite to the l., fr. broadly oblong.—*A. N. H.* xi. 188. *E. B. S.* 2944.—A decumbent floating plant well marked by the submersed lts. being divided into finger-like acute broadly linear parallel segments deeply cut at the end. Lts. of the upper l. broader than those of *Æ. Phellan'drium*.—Streams. B. or P. VII.—IX. E. I.

## 16. ÆTHUSA Linn. Fool's Parsley.

1. *Æ. Cynápium* (L.); partial involucre of 3 leaves longer than their umbel, l. all doubly pinnate, leaflets lanceolate decurrent pinnatifid.—*E. B.* 1192.—St. 4—18 in. high. L. deltoïd, dark green, lurid, stinking. General inv. 0; partial long, narrow, pendulous, all on one side. Herb poisonous.—Cultivated land. A. VII. VIII. E. S. I.

17. FŒNIC'ULUM *Hoffm.* Fennel.

1. *F. officinale* (All.) ; st. terete below, l. 3—4 times pinnate, segments all capillary long in the upper l. flaccid, umbels of many rays concave.—*E. B.* 1208.—Involucres 0. St. 3—4 feet high, usually filled with pith, branching. Umbels large. Fl. yellow. Whole herb aromatic. Segments of l. channelled, usually capillary in the wild plant ; but the cultivated plant with awlshaped segments is probably the same species.—Rocks and walls, particularly near the sea. P. VII. VIII. E. I.

18. SES'ELI *Linn.*

1. *S. Libanotis* (Koch) ; l. doubly pinnate cut, segments lanceolate mucronate, the lowermost leaflets crossing, general involucre of many leaves, fr. hairy.—*Athamanta* L. *E. B.* 138.—St. 1—3 feet high, covered at the base with the fibrous remains of decayed petioles. Umbels terminal, convex, with many downy rays.—Chalk hills of Cambr., Herts, and Sussex. P. VII. VIII. E.

19. HALOS'CIAS *Fries.* Scottish Lovage.

1. *H. scoticum* (Fr.) ; l. twice ternate, leaflets ovate somewhat rhomboidal dentate-serrate opaque, involucre of 5—7 linear-lanceolate leaves, cal. 5-toothed.—*Ligusticum* L. *E. B.* 1207.—St. herbaceous, nearly simple, striate, tinged with red, 1—1½ foot high. Lts. large, lobed and cut. Interstices with 3, commissure with 6 stripes. Seed quite free in the carpel.—Rocks on the Northern sea-coast. P. VII. E. S. I.

20. SILA'US *Besser.* Sulphur-wort.

1. *S. pratensis* (Bess.) ; st. angular, radical l. 3—4 times pinnate, leaflets lanceolate entire or bifid, terminal lt. tripartite, general involucre of 1—2 leaves, partial of many leaves.—*E. B.* 2142.—St. 1—2 feet high. L. mostly radical, stem-l. decreasing upwards. Fl. pale yellow.—Damp meadows and pastures. P. VI.—IX. E. S. I.

21. ME'UM *Tourn.* Bald-money.

1. *M. athamaniticum* (Jacq.) ; l. bipinnate, leaflets in many threadshaped acute segments.—*E. B.* 2249.—St. 1—2 feet high, round, clothed at the base with the fibrous remains of the decayed petioles. Fl. many, whitish yellow. General involucre of 2 or 3 leaves, partial more numerous. Highly aromatic.—Dry mountainous pastures. P. VI. VII. E. S.

## 22. CRITH'MUM Linn. Sapphire.

1. *C. maritimum* (L.).—*E. B.* 819.—St. 6—12 in. long. L. fleshy, 2—3-pinnate; leaflets lanceolate, narrowed at both ends, few. Involucre of many lanceolate acute leaves. Fl. whitish.—On rocky sea-coasts. P. VIII. E. S. I.

Tribe VI. *Angeliceæ*.

## 23. ANGEL'ICA Linn.

1. *A. sylvestris* (L.); leaflets equal ovate-lanceolate or ovate incise-serrate not decurrent, lateral lts. rather unequal at the base.—*E. B.* 1128.—St. 2—3 feet high, slightly downy above, purplish. Fl. pinkish white. Inv. deciduous. Lts. often subcordate at the base.—Wet places. P. VII. VIII. E. S. I.

## 24. ARCHANGEL'ICA Hoffm.

[*A. officinalis* (Hoffm.); leaflets all sessile partly decurrent, terminal lt. trifid, foliage stalks and even fl. bright green, l. 2—3 feet wide (*E. B.* 2516), is not a native. Watery places. P. VII.—IX.]

Tribe VII. *Peucedaneæ*.

## 25. PEUCED'ANUM Linn.

1. *P. officinale* (L.); l. 5 times ternate, leaflets linear very long acute flaccid, general involucre 3-leaved deciduous, pedicels much longer than the fruit.—*E. B.* 1767.—Fl. yellow. St. terete, striate, 2—3 feet high. Stripes of the commissure superficial.—Salt marshes. Kent. Essex. P. VII.—IX. E.

2. *P. palustre* (Moench); l. 3-pinnate, leaflets pinnatifid with linear-lanceolate acuminate segments, general involucre of many persistent lanceolate deflexed leaves, st. furrowed.—*Selinum* Sm. *E. B.* 229.—St. erect, 3—5 feet high. Fl. white. Stripes of the commissure deeply seated.—Marshy and fenny places, rare. P. VII. VIII. E. S. ?

†3. *P. Ostrúthium* (Koch); l. biternate, leaflets broadly ovate lobed cut and serrate, sheath very large, general involucre 0, cal.-segments inconspicuous.—*E. B.* 1380.—St. 1—2 feet high. Fl. white.—Moist meadows, rare. P. VI. *Masterwort*. S.

## 26. PASTINA'CA Linn. Parsnep.

1. *P. sativa* (L.); st. angular furrowed, l. pinnate downy be-

neath, leaflets ovate-oblong crenate-serrate often with a lateral lobe at the base, inv. 0, fr. oval.—*E. B.* 556. *T. pratensis* *Jord.*—St. 2—3 feet high. Fl. yellow. L. generally shining above, downy beneath.—Hedgebanks on a calcareous soil. B. VII. E. I.

### 27. HERAC'LEUM *Linn.* Cow-parsnep.

1. *H. Sphondylium* (L.); l. pinnate, leaflets lobed or pinnatifid cut and serrate, fr. at length glabrous.—*E. B.* 939.—St. 4 feet high. Lower l. very large. L. sometimes narrow. Umbels large flattish. Fl. white or reddish, outer fl. radiant.—Hedge-banks. P. VII. *Hog-weed.* E. S. I.

### 28. TORDYLIUM *Linn.* Hartwort.

†1. *T. maximum* (L.); outermost pet. radiant with 2 equal lobes, partial involucre linear shorter than the umbel, fr. hispid the thickened margin slightly crenate.—*E. B.* 1173.—St. 2—4 feet high. Fl. reddish.—Waste ground, very rare. A. VII. E.

[*T. officinale* (L.), *E. B.* 2440, was a mistake.]

## Tribe VIII. *Silerineæ.*

### 29. SILER *Scop.*

[*S. trilobum* (Scop.).—*R.* xxi. 153. *J. of B.* ix. t. 118.—St. solid, glaucous when young. Radical l. triternate, irregularly and coarsely serrate, long-stalked, dark green, glabrous. St.-l. ternate. Umbels terminal, large. Inv. of 1—3, partial of 5—8 bracts. Fr. large, crowned with persistent reflexed styles.—Naturalized at Cherry Hinton, Camb. P. V. VI.] E.

## Tribe IX. *Daucineæ.*

### 30. DAUCUS *Linn.* Carrot.

1. *D. Caróta* (L.); radical l. with an oblong narrow outline bipinnate with incise-dentate lts. and acute segments, upper l. broader below with lanceolate segments.—*E. B.* 1174.—St. 2—3 ft. high, hairy; branches ascending. Umbel of fr. usually concave. Prickles of fr. slender, mostly distinct, about equalling its breadth, spreading, tipped with 1—3 recurved minute bristles.—Pastures. B. VI.—VIII. E. S. I.

2. *D. gum'mifer* (Lam.); radical l. triangular broad 2—3-pinnate with ovate cut or pinnatifid lts. and blunt mucronate segments, upper l. narrower below.—*D. maritimus* With. (not Lam.).

*E. B.* 2560.—St. short, very hispid below; branches divaricate. Prickles of fr. usually flattened and often united below and shorter than its breadth, incurved, tipped with one bristle. L. shining above, rather fleshy. Umbel of fr. usually convex.—The forms of *Daucus* on the South coast deserve further study.—Sea-coasts in the South-west, rare. B. VII. VIII. E. I.

## Suborder II. *Campylospermæ*. Tribe X. *Caucalineæ*.

### 31. CAUCALIS Hoffm. Hen's-foot.

1. *C. daucoides* (L.); l. bipinnate, leaflets pinnatifid with linear-acute segments, general involucre 0, partial umbels of few fl. with involucre of 3—5 leaves, secondary ridges of the fr. each with one row of glabrous hooked prickles.—*E. B.* 197.—St. 6—12 in. high, furrowed, hairy at the joints. General umbels 3-cleft; partial bearing about 3 large oblong very prickly fruits. Fl. small, reddish.—Corn-fields on a chalky soil. A. VI. E.

†2. *C. latifolia* (L.); l. pinnate, leaflets lanceolate decurrent coarsely serrate, inv.-l. oblong membranous, secondary ridges of the fr. with 2 or 3 rows of retrorsely scabrous prickles.—*E. B.* 198. *Turgenia* Hoffm.—St. 1—2 feet high, rough. General umbels about 3-cleft; partial bearing about 5 large oblong very prickly fruits. Fl. large, pink.—Corn-fields, mostly on a chalky soil, very rare. Formerly abundant in Cambridgeshire. A. VII. E.

### 32. TORILIS Adans. Hedge-Parsley.

1. *T. Anthriscus* (Gaert.); l. bipinnate, leaflets ovate-oblong incise-serrate, umbels long-stalked terminal, general involucre of many leaves, fr. with subulate incurved prickles not hooked at the tip.—*E. B.* 987.—St. erect, 1—3 feet high. Umbels on long stalks. Fr. densely prickly. Fl. small, white or reddish.—Hedges and banks. A. VII. VIII. E. S. I.

2. *T. infesta* (Spr.); l. bipinnate, leaflets ovate-lanceolate incise-serrate, umbels long-stalked terminal, general involucre of one leaf or 0, fr. with spreading asperous prickles hooked at the tip.—*E. B.* 1314. *T. helvetica* Gm.—St. erect, usually much and densely branched, 6—18 in. high. Umbels on long stalks. Fr. densely prickly, primary ridges with adpressed prickles. Fl. small, white or reddish. Styles scarcely twice as long as the stylopode.—Fields. A. VII. VIII. E.

3. *T. nodosa* (Gaert.); lower l. bipinnate, upper l. pinnate, lts. deeply narrowly and uniformly pinnate, umbels nearly sessile dense lateral, outer carpels with bristles hooked at the tip, inner often warted.—*E. B.* 199.—St. diffuse, often prostrate. Umbels

very small, nearly globular.—Banks and dry places.—A. V.—VII. E. S. I.

### Tribe XI. *Scandicinea*.

#### 33. SCAN'DIX Linn. Shepherd's Needle.

1. *S. Pecten-Ven'eri* (L.); fr. rough dorsally compressed glabrous with bristly edges, beak three times as long as fruit, lts. of partial inv. entire or bifid longer than the pedicels.—*E. B.* 1397.—St. often a foot high. L. light green, triply pinnate; segments short linear. Umbels 1—2 together, small. Fl. often slightly radiant. Styles always straight. Stylopode purple. Fr. and beak nearly 2 in. long. Partial involucre sometimes much divided.—Fields. A. VI.—IX. E. S. I.

#### 34. CHÆROPHYL'LUM Linn. Chervil.

\* *Ridges only apparent on the beak of the fr. ; no stripes.*  
ANTHRISCUS Hoffm.

1. *C. sylves'tre* (L.); st. hairy below glabrous upwards swollen below the joinings, umbels terminal stalked, l. bipinnate, leaflets pinnatifid, fr. smooth and shining narrow to its tip.—*Anthriscus* Hoffm. *E. B.* 752.—St. 3 feet high, erect, leafy, furrowed, hollow, branched. Partial involucre of several ovate-lanceolate ciliate leaflets. Umbels at first drooping. Pet. oblong-obovate, scarcely emarginate, with a short inflexed point.—Hedges and banks. P. IV.—VI. *Wild Chervil.* E. S. I.

[*C. sativum* (Lam.); st. hairy above the joinings only, umbels lateral sessile, l. tripinnate, leaflets ovate pinnatifid, fr. linear smooth about twice as long as its beak.—*Anth. Cerefolium* Hoffm. *E. B.* 1268.—St. 1—3 feet high, slender, striate, much branched. Partial involucre of 3 unilateral linear-lanceolate leaflets. Peduncles downy.—Waste ground. Probably an escape from cultivation. A. V. VI. *Garden-Chervil.*] E. S.

2. *C. Anthris'cus* (Lam.); st. glabrous, umbels lateral stalked, l. tripinnate, leaflets pinnatifid, fr. ovate with hooked bristles about twice as long as its glabrous beak.—*E. B.* 818. *A. vulgaris* Pers.—St. erect, 2 feet high, branched. L. slightly hairy. Umbels on rather short stalks. Partial involucre of few ciliate leaflets.—Waste places. A. V. VI. E. S. I.

\*\* *Ridges 5, equal, blunt ; stripes solitary.*

3. *C. tem'ulum* (L.); st. thickened beneath the joinings rough, l. bipinnate, leaflets ovate-oblong pinnatifid with rather acute mucronate segments, pet. glabrous, styles equalling the stylopode.

—*E. B.* 1521.—*St.* 3—4 feet high, round, solid, spotted, rough below, hairy near the top. Umbels at first nodding. Pet. deeply obcordate.—Hedge-banks. *P. VI. VII. Rough Chervil.*  
E. S. I.

### 35. MYR'RHIS Scop. Sweet Cicely.

1. *M. odorata* (Scop.); l. downy beneath, leaflets of the partial involucre lanceolate-acuminate.—*E. B.* 697.—*St.* 2—3 feet high, round, leafy, hollow. L. very large, tripinnate. Leaflets ovate-lanceolate pinnatifid. Umbel terminal. Fl. many, white. Fr. large, nearly an inch long, dark brown. Whole plant highly aromatic. Pastures in hilly districts. *P. V. VI. E. S. I.*

## Tribe XII. Smyrnieæ.

### 36. ECHINOPH'ORA Linn.

1. *E. spinosa* (L.); l. pinnate, leaflets pinnatifid with spinous awlshaped entire segments.—*E. B.* 2413.—Formerly found on the Lancashire and Kentish shores. *P. VII. E.*

### 37. CONI'UM Linn. Hemlock.

1. *C. maculatum* (L.); leaflets of the partial involucre unilateral ovate-lanceolate with an attenuate point shorter than the umbels.—*E. B.* 1191.—*St.* 3—5 feet high, erect, round, hollow, glaucous, spotted with purple, branched. L. tripinnate; leaflets lanceolate, pinnatifid with acute cut segments. Readily distinguished by its foetid smell, spotted stem, unilateral partial involucre, and wavy crenate ridges of the fruit. Highly poisonous.—Hedge-banks and waste places. *B. VI. VII. E. S. I.*

### 38. PHYSOSPER'MUM Cusson.

1. *P. cornubien'se* (DC.); radical l. triternate, leaflets wedge-shaped cut or deeply 3-lobed with acute segments, stem-l. ternate lanceolate entire.—*E. B.* 683.—*St.* 1—3 feet high, erect, round, striate, bearing a few small ternate leaves with linear-lanceolate segments, the uppermost represented by a barren lanceolate acute sheath. Umbels terminal and axillary. Carp. longer than broad; the coat loose. Seed free.—Devon and Cornwall, rare. *P. VII. VIII. E.*

### 39. SMYR'NIUM Linn. Alexanders.

1. *S. Olusatrum* (L.); st. terete, stem-l. ternate stalked serrate.—*E. B.* 230.—*St.* 3—4 feet high, stout, branched, leafy,

furrowed. Radical l. very large, 3—4 times ternate; all with large membranous sheaths and large ovate shining cut and serrate leaflets. Fl. greenish yellow in rounded often dense umbels. Fr. nearly black, aromatic.—Waste ground and near ruins. B. V. VI. E. S. I.

### Suborder III. *Cœlospermæ*. Tribe XIII. *Coriandrea*.

#### 40. CORIAN'DRUM *Linn.* Coriander.

[1. *C. sativum* (L.).—*E. B.* 67.—St. 12—18 in. high, leafy, round, striate. L. bipinnate, cut, with broad roundish or wedge-shaped segments; upper l. more divided with linear segments. Fl. white.—Waste places, scarcely naturalized. A. VI.] E. S.

### Order XXXVIII. HEDERACEÆ.

Cal. 4—5-toothed, adnate to the ovary. Pet. 5—16, rarely wanting; valvate in the bud. Stam. as many as the pet. and alternate with them or twice as many, inserted below the margin of an epigynous disk. Ovary with 2 or more cells. Styles as many as the cells. Fr. succulent or dry, of several cells each with 1 pendulous seed. Albumen fleshy. Embryo minute (not so in our *Hedera*).—L. alternate without stipules.

1. HEDERA. Cal. superior, limb of 5 teeth. Pet. 5—10, not cohering at the apex. Stam. 5—10. Styles 5—10, connivent or combined into one. Berry 5-celled and 5-seeded, crowned with the calyx.

#### 1. HED'ERA *Linn.* Ivy.

1. *H. Hélix* (L.); l. coriaceous ovate or cordate and 5-lobed, lobes angular, umbels simple downy erect.—*E. B.* 1267.—Climbing by means of rootlike fibres. L. of the flowering branches ovate-oblong, acute, entire. Berries black. Embryo like that of *Cornaceæ*.—Rocks, old walls, hedges. Sh. X. XI. E. S. I.

### Order XXXIX. CORNACEÆ.

Cal. 4-lobed, adnate to the ovary. Pet. 4, oblong, broad at the base, inserted at the top of the calyx-tube; valvate in the bud. Stam. 4. Ovary 2-celled. Style filiform. Fruit a drupe, crowned with the remains of the calyx. Seed pendulous, solitary. Embryo in the axis of fleshy albumen and as long as it.—Leaves opposite, exstipulate.



1. CORNUS. Calyx-limb superior, of 4 teeth. Pet. 4. Stam. 4. Style 1. Drupe with a 2-celled and 2-seeded nut.

1. COR'NUS *Linn.*

1. *C. sanguinea* (L.); *arborescent*, branches straight, l. ovate cuspidate green on both sides, cymes flat without an involucre.—*E. B.* 249. *St.* 52. 3.—Shrub 5—6 feet high. Old bark reddish. Fl. many, white, in terminal cymes. Fr. dark purple. L. mostly opposite, strongly veined, acutely cuspidate, rounded below.—Hedges and thickets. Sh. VI. *Dog-wood.* E. I.

2. *C. suécica* (L.); *herbaceous*, l. all opposite sessile ovate, fl. umbellate shorter than the 4-leaved *petal-like involucre*.—*E. B.* 310. *St.* 52. 1.—Flowering shoots about 6 in. high, annual, springing from the procumbent or subterranean creeping woody leafless stems. Fl. dark purple with yellow stamens, in a small solitary terminal umbel with an inv. of 4 ovate yellow l. tipped with purple. Fr. red.—Moist alpine pastures. P. VII. E. S.

Division III. COROLLIFLORÆ.

(Orders XL.—LXIII.\*)

Pet. united. Stamens epipetalous; except in *Ericaceæ* and part of *Plantaginaceæ*, which have hypogynous, and *Campanulaceæ*, which has epigynous stamens.

Order XL. LORANTHACEÆ.

Cal. adnate to the ovary, with two bracts at its base; limb entire or lobed. Cor. of 4—8 more or less united petals. Stam. as many as and opposite to the petals, with which the filaments more or less combine; anth. sometimes adnate to the petals. Ovary 1, 1-celled with 1 pendulous ovule. Style filiform or 0. Stigma capitate. Fr. succulent. Albumen fleshy.—Parasitical plants with entire mostly opposite leaves and no stipules. Connects this Division with Div. Calycifloræ and Monochlamydeæ.

1. VISCUM. Dioecious. Male: Cal. 0. Pet. 4, ovate, fleshy, united at the base. Anth. adnate to the petals, many-celled. Fem.: Cal. an obscure entire superior margin. Pet. 4, erect, somewhat triangular, minute. Stigma sessile, blunt. Berry 1-seeded, crowned with the calyx.

\* Orders XL.—XLVI. and Tribe 3 of XLVII. are often regarded as Calycifloral. Their ovary is inferior. Order XL. is sometimes placed in Monochlamydeæ.

1. *VIS'cum* Linn. Mistletoe.

1. *V. al'bum* (L.) ; st. repeatedly forked, branches terete, l. ovate-lanceolate blunt, fl. in the forks of the stem sessile clustered.—*E. B.* 1470.—Evergreen, yellow, succulent. Spreads by runners under living bark of tree. Male fl. about 3 together, female about 5, yellowish. Berries white, pellucid, globular, viscid. L. of male plant broader.—Parasitical on various trees. P. III. IV. E.

## Order XLI. CAPRIFOLIACEÆ.

Cal. adnate to the ovary, usually with bracts at the base ; limb 4—5-lobed. Cor. regular or irregular, 4—5-cleft. Stam. 4 or 5, free, on the corolla, and alternate with the lobes. Ovary 3—5-celled. Stigmas 1—3. Fruit not bursting, 1- or many-celled, usually fleshy. Albumen fleshy.—L. opposite. Stip. rare.

1. *ADOXA*. Cal.  $\frac{1}{2}$ -inferior, 2—3-cleft. *Cor. rotate*, 4—5-lobed. *Stam.* 8—10, *in pairs alternate with the lobes* of cor. ; *anth.* 1-celled. *Styles* 5—10. *Fr.* 4—5-celled ; cells 1-seeded.—L. ternate, lobed.
2. *SAMBUCUS*. Cal.-limb 5-cleft. *Cor. rotate*, 5-lobed. *Stam.* 5. *Stigmas* 3, sessile. *Fr.* 3—4-seeded.—L. pinnate.
3. *VIBURNUM*. Cal.-limb 5-cleft. *Cor. bell- or funnel-shaped*, 5-lobed. *Stam.* 5. *Stigmas* 3, sessile. *Fruit* 1-seeded.—L. simple.
4. *LONICERA*. Cal.-limb small, 5-cleft. *Cor. tubular* or funnelshaped, limb 5-fid or irregular. *Stam.* 5. *Style fili-* form. *Stigma* capitate. *Fruit* 1—3-celled, few-seeded.
5. *LINNÆA*. *Cal.-limb* 5-cleft, *with lanceolate subulate equal deciduous segments*. *Cor.* turbinate bell-shaped, 5-lobed. *Stam.* 4, rarely 5, 2 longer. *Style* filiform ; *stigma* capitate. *Fr.* dry, 3-celled ; 2 cells barren, 1 single-seeded.—Two large and 2 minute bracts at the base of the fruit.

1. *ADOX'A* Linn. Moschatel.

1. *A. Moschatel'ina* (L.).—*E. B.* 453.—Rhizome white, fleshy, toothed, soboliferous. St. solitary, erect, simple, 3—4 in. high, with 2 opposite leaves, and a head of 4 whorled and 1 terminal flowers. Root-l. ternate, 3-lobed. Stam. often more or less united in pairs, showing their number to be normally 4.

Odour musky. Terminal fl. usually divided in fours, the others in fives, but the numbers vary.—Woods and shady hedge-banks. P. IV. V. E. S. I.

## 2. SAMBU'CUS Linn. Elder.

1. *S. Ebu'ulus* (L.); *herbaceous*, st. furrowed, *stip. leaflike* ovate serrate, l. pinnate, leaflets lanceolate serrate, cyme with 3 principal branches.—*E. B.* 475.—St. 2—4 feet high. Cymes terminal. Fl. white, reddish externally; anth. purple. Fr. reddish black.—Hedge-banks. P. VIII. *Dwarf Elder. Danewort.* E. S. I.

2. *S. nigra* (L.); *arborescent*, *stip. inconspicuous* or wanting, l. pinnate, leaflets ovate cuspidate serrate, cymes with 5 principal branches.—*E. B.* 476.—A small tree. Cymes large, terminal. Fl. cream-coloured. Fr. black, rarely green or white.—\*3. *laciniata*; l. 2—3-pinnate, lts. laciniate.—*γ. rotundifolia* (DC.); lts. usually 3 orbicular.—Woods and hedges. *γ. Isle of Wight.* T. VI. *Elder.* E. S. I.

## 3. VIBUR'NUM Linn. Guelder-rose.

1. *V. Lantána* (L.); l. *oblong* with a cordate base *finely denticulate-serrate* downy beneath, pubescence stellate.—*E. B.* 331.—A shrub with round mealy branches. Young shoots, petioles, and undersides of the l. densely, upperside more sparingly, covered with stellate down. Cymes terminal. Fl. white, not radiant; fr. black.—Hedges and thickets on a calcareous soil. T. V. *Mealy Guelder-rose. Wayfaring tree.* E.

2. *V. Op'ulus* (L.); l. 3—5-lobed, lobes acuminate and dentate, petioles with glands.—*E. B.* 332. *St.* 27. 6.—Branches glabrous, tetragonal when young. L. slightly downy beneath, with linear adnate stipules (?). Cymes large, with linear bracts; fl. white, inner ones fertile, outer barren and radiant. Fr. red.—Hedges and thickets. T. VI. VII. *Common Guelder-rose.* E. S. I.

## 4. LONICE'RA Linn. Honeysuckle.

†1. *L. Caprifólium* (L.); fl. ringent whorled terminal *sessile*, l. deciduous glabrous on both sides blunt, *upper l. connate-perfoliate*, style glabrous.—*E. B.* 799.—St. twining. Fl. white or purplish. Fr. orange. Upper pairs of leaves connate, the rest distinct.—Thickets. Sh. V. VI. E. S. ?

2. *L. Perichly'menum* (L.); fl. ringent in terminal stalked

clusters, *l. all distinct* deciduous oval, st. twining.—*E. B.* 800.—Fl. pale yellow, externally red. Fr. red. L. sometimes downy beneath, often lobed when young.—Woods and hedges. Sh. VI.—IX. *Honeysuckle. Woodbine.* E. S. I.

3. *L. Xylos'teum* (L.); *peduncles 2-flowered* downy as long as the flowers, calyx-limb deciduous, berries slightly connected at the base, *l. oval* downy, st. erect.—*E. B.* 916.—Shrub. Fl. pale yellow. L., bracts, cal., cor. externally, filaments, and style downy. Fr. scarlet.—Thickets. Native in Sussex. (*Borr.*) Sh. V. E.

#### 5. LINNÆ'A Gronov.

1. *L. boreâlis* (Gron.).—*E. B.* 433.—St. trailing and creeping. L. opposite, broadly ovate, stalked, dark green above, paler beneath. Peduncles long, erect, 2-flowered, from short lateral branches with 2—4 leaves. Fl. drooping, flesh-coloured, purple within.—Woods, chiefly of fir, in the North. P. VII. E. S.

### Order XLII. RUBIACEÆ.

Cal. superior, entire or lobed. Cor. regular, 4—6-lobed. Stam. 4—5, alternate with the lobes of the corolla. Ovary 1, 2-celled, with solitary erect ovules. Style 1, often bifid. Stigmas 2. Fr. a didymous indehiscent pericarp. Embryo straight in horny albumen.—St. herbaceous, square. Many raphides.

1. SHERARDIA. Cor. funnelshaped. Fr. crowned with the deeply 6-toothed calyx, dry.—Fl. umbellate, involucrate.
2. ASPERULA. Cor. funnelshaped. Fr. dry. Limb of the calyx inconspicuous.
3. GALIUM. Cor. rotate, 4-fid. Fr. dry. Limb of cal. inconspicuous.
4. RUBIA. Cor. rotate, 5- (or rarely 4-)fid. Fr. succulent, 2-lobed. Limb of cal. inconspicuous.

#### 1. SHERARDIA Linn. Field-Madder.

1. *S. arven'sis* (L.).—*E. B.* 891.—St. mostly decumbent, branched, leafy. L. 6 in a whorl, obovate-lanceolate, acute. Fl. lilac, in a small sessile terminal umbel with 7—8 inv.-leaves. Cal.-segm. 4, 2 bifid.—Fields. A. V.—VII. E. S. I.

#### 2. ASPERULA Linn.

1. *A. cyan'chica* (L.); *l. 4 in a whorl*, linear, uppermost l.

very unequal, fl. corymbose, bracts lanceolate mucronate, cor. rough.—*E. B.* 33.—Root fusiform. Stems many, diffuse or ascending, branched. Lowest l. very obovate; intern. obovate-lanceolate; uppermost lanceolate-attenuate. Fl. generally lilac. Fr. wrinkled and tubercled.—Dry banks in limestone districts. P. VI. VII. *Quinancy-wort.* E. I.

2. *A. odorata* (L.); l. 6—9 in a whorl lanceolate, margins rough with forward prickles, fl. in stalked terminal cymose corymbs, fr. hispid.—*E. B.* 755.—St. erect, about 6 in. high. Fl. white. L. broad, lower in sixes. Whole plant fragrant.—Woods. P. V. VI. *Woodruff.* E. S. I.

[*A. arven'sis* (L.); l. 6—10 in a whorl linear-lanceolate blunt, fl. clustered terminal surrounded by long ciliate bracts, fr. glabrous.—*E. B. S.* 2792.—Like *Sherardia arvensis*. Fl. bright blue.—Introduced. A. VI.] E.

[*A. taurina* (L.); l. 4 in a whorl elliptic acuminate 3-veined, fl. corymbose, cor.-tube very long, fr. rather rough.—*Sy. E. B.* 662.—Cadeby, Leices. Casterton, Westm.] E.

### 3. GA'LIIUM Linn. Bedstraw.

#### A. Leaves 3-veined.

\* *Fl. in a terminal panicle, perfect, white; fr.-stalks erect.*

1. *G. boreale* (L.); l. 4 in a whorl lanceolate, st. erect paniced, fruitstalks patent, fr. covered with hooked bristles.—*E. B.* 105.—St. about 18 in. high; branches many, leafy. Fl. in compact panicles.—Moist rocky places. P. VII. VIII. E. S. I.

\*\* *Fl. axillary, yellow; lateral fl. imperfect; fr.-stalks deflexed.*

2. *G. cruciatum* (With.); l. 4 in a whorl elliptic-oblong hairy, fl. corymbose bracteate, terminal fl. fertile, lateral fl. mostly male, fr. smooth.—*E. B.* 143.—St. simple above, 1—2 feet high, hairy. Fl. small, about 8 together in small corymbs, falling short of the leaves.—Hedges and thickets. P. V. VI. *Crosswort.* E. S.

#### B. Leaves 1-veined. Root annual.

\* *Flowers axillary; lateral fl. imperfect.*

[*G. saccharatum* (All.).—*E. B.* 2173.—Accidental. A. VI.—VIII.]

\*\* *Fl. in axillary cymes, all perfect, white or greenish.*

3. *G. tricorne* (With.); l. 6—8 in a whorl linear-lanceolate with marginal backward prickles, st. rough with deflexed prickles, cymes 3-flowered, fr. granular on reflexed peduncles.—*E. B.* 1641.—St. procumbent, spreading. Fl. small, all 3 appearing perfect, the middle one usually alone fertile, cream-coloured. Fr. large, a double globe, covered with small granulations.—Dry calcareous fields. A. VI.—IX. E.

4. *G. Aparine* (L.); l. 6—8 in a whorl linear-lanceolate with marginal backward prickles, st. rough with deflexed prickles, cymes few- (about 3-) flowered, fruitstalks divaricate straight, fruit covered with short hooked bristles.—*E. B.* 816.—St. straggling amongst bushes, 3—4 feet long. The marginal prickles near the extremity of the l. point forwards, the rest backwards. Fl. small, pale. Ped., or rather flowering branches, with several l. at the primary divisions. Fr. large.—Common. A. VI.—VIII. Goose-grass. Cleavers. E. S. I.

5. *G. spúrium* (L.); l. 6—8 in a whorl linear-lanceolate with marginal backward prickles, st. rough with deflexed prickles, cymes with 3—9 flowers, fruitstalks divaricate straight.—Closely resembling the preceding; distinguished by its more numerous green flowers, floral leaves solitary ("or in pairs"), fruit of about half the size.—*a*; fr. smooth. *E. B.* 1871.—*β. G. Vailantii* (DC.); fr. hispid. *E. B. S.* 2943.—Fields. *α.* Forfar; *β.* Saffron Walden and Chesterford, Essex. A. VII. E. S.

6. *G. anglicum* (Huds.); l. about 6 in a whorl linear-lanceolate bristle-pointed with marginal forward prickles, st. rough with deflexed prickles, cymes small forked with divaricate bifid branches.—*E. B.* 384.—St. 6—8 in. high, spreading, slender, brittle. L. usually 6 in a whorl, the lowermost sometimes in fours. Branches of the small panicles often spreading nearly at right angles with their stalk. Fr. granular, nearly black. [*G. parisiense* (L.) has hooked prickles on the fruit and can hardly be a form of this species.]—Old walls and dry sandy places. A. VI. VII. E.

C. Leaves 1-veined. Root perennial. Fl. in terminal panicles, white (except in *G. verum*).

\* *Fruit not granular. No downward prickles on the stem.*

7. *G. erectum* (Huds.); l. about 8 in a whorl lanceolate mucronate the margins rough with forward prickles, midrib slender, branches of the pyramidal panicle all ascending, fruitstalks divaricate, fr. oval smooth, pet. taper-pointed.—*E. B.* 2067.—St.

glabrous or hairy, erect, not much branched. L. lanceolate, scarcely at all obovate, those of the main st. erect, patent; veins not translucent; margins with 2 rows of prickles pointing forwards. Fl. white.—*a*; l. lanceolate.—[*β. G. cinereum* (Sm.); l. 6—8 in a whorl linear. *E. B. S.* 2783. Perhaps a distinct species, *G. diffusum* (Hook.), but a doubtful native.]—*G. aristatum* (Sm.) has l. in sixes, but is probably a state of this species. *E. B. S.* 2784.—Banks and pastures. P. VI. and IX. E. I.

8. *G. Mollígo* (L.); l. about 8 in a whorl lanceolate-obovate or obovate-oblong cuspidate the margins rough with forward prickles, branches of the broad pan. spreading lower ones horizontal or deflexed, fruitstalks divaricate, fr. glabrous.—*E. B.* 1673.—St. ascending, square, thickened at the joinings, glabrous or hairy. L. slightly translucent, veined, hardly separated at the base, those of the main st. horizontal or declining. Pan. large. Fl. small, white. Styles nearly free.—*β. G. insubricum* (Gaud.); l. about 6 in a whorl obovate abruptly cuspidate, pan.-branches few-flowered terminating in trichotomous umbels, floral l. large, bracts large usually solitary.—*γ. Bakeri* (Syme); l. 6—8 in a whorl linear-lanceolate, pan. with few-flowered ascending branches. It may be distinct.—Hedges and thickets. P. VII. VIII. E. S. I.

9. *G. vérum* (L.); l. about 8 in a whorl linear-setaceous with revolute margins channelled above downy beneath, panicles many small densely flowered subterminal, fruitstalks patent, fruit smooth, pet. blunt apiculate.—*E. B.* 667.—St. erect, slightly branched, somewhat woody, with many whorls of narrow deflexed leaves. Fl. golden yellow, rarely green or straw-coloured, (*G. decolorans* Gren.?) usually in many small dense panicles collected into a kind of terminal spike. St. and upper surface of the l. sometimes downy or rough. On loose sands the st. are much more branched and the fl. sometimes solitary.—Dry and sandy places. P. VII. VIII. E. S. I.

\*\* Fruit granular, not hairy. St. without downward prickles.

10. *G. saxatile* (L.); l. about 6 in a whorl obovate pointed flat, midrib slender, panicles corymbose small, fl.- and fr.-stalks erect-patent, pet. acute. *E. B.* 815.—St. many, procumbent, much branched. L. suddenly narrowed to a point, smooth with a few marginal forward prickles; lower l. roundly obovate. It turns black in drying.—Heaths. P. VII. VIII. E. S. I.

11. *G. sylvestre* (Poll.); l. 6—8 in a whorl linear or linear-lanceolate mucronate with revolute edges, midrib slender prominent, panicles few-flowered, fl.- and fr.-st. erect-patent, fr. faintly granular, pet. acute.—*G. pusillum* Sm. *E. B.* 74.—St. many,

slender, square, diffuse, ascending. L. often nearly glabrous or with *marginal hairs* (not prickles) spreading or backward. Lower part of stem and leaves sometimes densely covered with patent hairs. Panicle very variable in size. Fr. very minutely granular.—*a. G. montanum* (Vill.); l. linear-lanceolate, pan. with short ascending branches and few-flowered cymes.—*β. G. nitidulum* (Thuil.); l. linear, pan. with somewhat spreading branches and compact cymes.—Limestone hills, rare. P. VI. VII. E. S. I.

\*\*\* *Stem rough with downward prickles.*

12. *G. uliginosum* (L.); l. 6—8 in a whorl linear-lanceolate bristle-pointed the margins rough like the angles of the stem with backward prickles, panicles small axillary few-flowered trichotomous, the branches patent 3-fid, fruitstalks divaricate straight, fruit granular.—*E. B.* 1972.—Stems slender, brittle, about a foot high, weak. L. usually 6 in a whorl, discoloured at the tip, shortly acuminate. Fr. dark brown.—Wet places. P. VII. VIII. E. S.

13. *G. palustre* (L.); l. 4 in a whorl broadly linear broader upwards blunt, midrib slender, panicle diffuse, fr.-st. straight, spreading at right angles, fr. smooth.—St. 1—2 feet high, slender, usually branched. Fl. small, white. St. and branches nearly smooth. L. narrow, lowest usually in sixes, upper in fours of which 2 are smaller.—On dry ground the l. are broader, those of the barren shoots often obovate: then known by its blunt l. and downward prickly stems.—*G. Witheringii* (Sm.) differs only by having rough edges to the leaves.—A very strong form, much larger in all respects, is the *G. elongatum* (Presl). *E. B.* 1857.—Wet places by ditches and rivers. P. VI. VII. E. S. I.

#### 4. RU'BIA Linn. Madder.

1. *R. peregrina* (L.); l. 4—6 in a whorl elliptic or lanceolate shining smooth above without veins the margin and keel rough with reflexed bristles.—*E. B.* 851.—Old st. terete; shoots spreading, square. L. rigid persistent. Fl. in paniced cymes. Cor. rotate, 5-cleft; lobes oval, suddenly narrowed into a slender point.—Stony and sandy thickets in the South. P. VI.—VIII. E. I.

### Order XLIII. VALERIANACEÆ.

Cal. superior; limb various, toothed, or inconspicuous, or involute and ultimately resembling a pappus. Cor. tubular,



3—5-lobed, unequal or irregular often spurred or gibbous at the base. Stam. 1—3, inserted in the tube, free, fewer than the cor.-lobes. Ovary with 1 perfect fertile cell and often 2 abortive cells; ovule solitary, pendulous. Fr. dry.—No stipules.

1. CENTRANTHUS. Cor. 5-lobed, *with a spur*. Stam. 1. Fr. 1-celled, indehiscent, crowned with the limb of the calyx expanded into a feathery pappus.
2. VALERIANA. Cor. 5-lobed, gibbous but *without a spur*. Stam. 3. Fr. 1-celled, indehiscent, crowned with the limb of the calyx expanded into a feathery pappus.
3. VALERIANELLA. Cor. 5-lobed, without a spur. Stam. 3. Fr. 3-celled, indehiscent, *crowned with the erect unequally toothed limb of the calyx*, 2 of the cells usually empty inflated or filiform.

### 1. CENTRANTHUS *Cand.*

1. *C. rúber* (DC.); l. lanceolate stalked, upper l. ovate-lanceolate sessile, spur much shorter than the cor.-tube twice as long as the germen.—*Valeriana* Sm. *E. B.* 1531.—St. 1—2 feet high. Fl. purple or white.—Chalk-pits and old walls. P. VI.—IX. *Red Valerian.* E. I.

\*2. *C. Calcitrápa* (Dufr.); radical l. ovate entire, stem-l. pinnatifid, spur very short.—*Sy. E. B.* 665.—Eltham, Kent. E.

### 2. VALERIA'NA *Linn.* Valerian.

1. *V. officinális* (L.); l. *all pinnate*, lts. 9—21 lanceolate dentate-serrate terminal one not larger than the others, st. sulcate solitary, fr. glabrous ovate-oblong.—*R.* xii. 727. *St.* 9.—*With suckers*, not stoles. *St.* 2—4 ft. high. Fl. flesh-coloured. Radical l. on long stalks.—*a*; lts. usually 9—11 near together, their anterior edge nearly entire, the posterior edge strongly toothed.—*β.* *V. sambucifólia* (Mikan); lts. of rt.-l. ovate-acute, of st.-l. oblong-lanceolate, all toothed on both edges. *E. B.* 698. Term. lt. of rt.-l. often slightly the largest. Stoles long.—Ditches, marshes and damp places. P. VI. VII. E. S. I.

\*2. *V. pyrenáica* (L.); l. *heartshaped* serrate stalked, upper l. with 1—2 pairs of small lanceolate leaflets.—*E. B.* 1591.—*St.* 2—3 feet high, furrowed. Fl. light rose-coloured.—Woods, rare. P. VI. VII. E. S.

3. *V. dioica* (L.); fl. imperfectly dioecious, *root-l. ovate* stalked, *stem-l. pinnatifid* with a large terminal lobe, fr. glabrous.—*E. B.*

628. *St.* 9.—*St.* a foot or more in height, simple. Fl. flesh-coloured; rather large with protruded stam. when barren; or small with included stamens and forming a closer corymb when fertile. Stoloniferous.—Boggy places. P. V. VI. E. S.

## 2. VALERIANELLA Moench.

\* *Fruit with 2 barren cells, fertile cell corky on the back.*

1. *V. olitória* (Moench); fr. compressed oblique, barren cells without furrows, dissepiment incomplete, bracts ciliate.—*E. B.* 811. *St.* 2. 3. *R.* xii. 708.—About 6 in. high. L. ovate-spathulate, upper ones narrower. Fl. in *terminal dense cymes* with oblong linear opposite bracts. Fr. 3-celled; 1 cell fertile with its back formed of a thick gibbous spongy mass usually traversed by one furrow, separated by a groove on each side from 2 barren slightly confluent cells each having a slender rib on its side and their junction marked by a slight furrow.—Corn-fields and banks. A. V. VI. *Corn-Salad.* E. S. I.

\*\* *Fruit with 2 barren conspicuous cells, fertile cell not corky.*

‡2. *V. carinata* (Loisel.); fr. oblong bootshaped crowned with 1 straight tooth, cells nearly equal each with a single rib on the back, barren cells contiguous in their whole length and with a deep furrow between them, fl. in dense cymes.—*E. B. S.* 2810. *R.* xii. 708.—About 6 in. high. Fl. pale blue. Root-l. spathulate, st.-l. oblong. Bracts ciliate. Fr.-section crescent-shaped.—Hedge-banks, rare. A. IV.—VI. *Lambs' Lettuce.* E. S.

3. *V. Auric'ula* (DC.); fr. subglobose crowned with 1 erect membranous leaf, barren cells larger than the fertile ones inflated contiguous having a narrow furrow between them, fl. scattered.—*E. B. S.* 2809. *V. dentata* DC., *R.* xii. 709.—About a foot high. Fl. distant, in the forks of a repeatedly forked cyme. Lower l. obovate attenuate downwards, upper l. oblong. Bracts ciliate. Section of the fruit nearly round. Crown of one oblong-blunt obliquely truncate tooth, sometimes with a minute tooth on each side; or of 3 acute teeth, of which one is much the longest and often 3-pointed.—Cultivated land. A. VII. VIII. E. S. I.

\*\*\* *Barren cells 0, or reduced to a rib.*

4. *V. dentata* (Deitr.); fr. oval crowned with the small oblique unequally 4-toothed calyx flat in front with a space enclosed between 2 elevated curved ribs convex behind, cyme lax spreading its branches long divaricate.—*E. B.* 1370. *R.* xii. 710. *V.*

*Morisonii* DC.—Teeth of the crown spreading or all incurved except the largest. Fr. smooth or hairy. Fl. corymbose.—Corn-fields and banks. A. VI. VII. E. S. I.

[*V. eriocar'pa* (Desv.); fr. pilose crowned with the large toothed open *nearly regular rather obliquely truncate net-veined calyx*; otherwise like *V. dentata*.—*Sy. E. B.* 673. *R.* xii. 712. *Coss. Atl.* 24 E.—Between Hanley and Barnard Green, Worcester. The Welsh plant is *V. dentata*. A. VI.] E.

#### Order XLIV. DIPSACACEÆ.

Fl. capitate, involucrate. Cal. superior, surrounded by an involucl (or free outer calyx) which closely invests the ripe fruit. Cor. 4—5-fid with unequal lobes. Stam. 4, inserted in the tube, free; anth. not cohering. Style 1. Stigma simple. Ovary 1-celled, with a pendulous ovule. Fr. crowned with the pappus-like calyx. Embryo in fleshy albumen.—No stipules.

1. DIPSACUS. Involucl forming a thickened margin to the ovary. Calyx cupshaped. Cor. 4-fid. *Receptacle with spinous exserted scales* shorter than the involucre. Fr. with 4 sides and 8 little depressions.
2. KNAUTIA. Involucl terminating in 4 small teeth. Calyx cupshaped with radiant teeth. *Receptacle hairy; scales 0*. Fr. with 4 sides and 4 little depressions.
3. SCABIOSA. Involucl membranous or minute. *Calyx of 4 or 5 bristles. Receptacle with sunken scales*. Fr. nearly cylindrical with 8 excavations.

#### 1. DIP'SACUS Linn. Teasel.

1. *D. sylves'tris* (L.); l. opposite simple sessile, stem-l. connate, *scales of the receptacle straight at the end exceeding the flowers, involucre curved upwards*.—*E. B.* 1032.—St. 5—6 feet high, prickly, leafy, branched. Heads of pale lilac fl. large, conical, overtopped by the slender ascending involucre.—Hedges and roadsides. B. VIII. IX. *Wild Teasel*. E. S. I.

[*D. Fullónum* (L.); *scales of the receptacle hooked at the end equalling the fl., invol. reflexed*, otherwise like *D. sylvestris*.—*E. B.* 2080.—Scarcely naturalized. B. VIII. IX. *Teasel*.] E.

2. *D. pilósus* (L.); l. stalked with a lt. at the base on each side, *scales of the receptacle obovate-cuspidate straight, involucre deflexed*.—*E. B.* 877.—St. 3—4 feet high, slender, branched,

rough, leafy. *Heads* of white fl. small, *globose*, exceeding the involucre.—Moist shady places. B. VIII. *Shepherd's Rod*. E.

### 2. KNAUT'IA *Coult.*

1. *K. arven'sis* (Coult.); lower l. simple, stem-l. pinnatifid, st. bristly, calyx with about 8 awned teeth.—*E. B.* 659.—St. 2—3 feet high, slightly branched, with a few leaves. Radical l. many, sometimes pinnately lobed. Fl. purple, in large convex long-stalked heads; outer usually unequal and radiant. Inv. bluntish.—Sometimes l. all simple narrowly lanceolate entire or superficially crenate.—Fields. P. VII.—IX. *Field Scabious*.

E. S. I.

### 3. SCABIO'SA *Linn.* Scabious.

1. *S. succisa* (L.); root-stock abrupt, heads of fl. and fr. nearly globose, *involucre hairy 4-fid herbaceous*, cor. 4-cleft, l. oblong entire, upper l. narrower mostly entire.—*E. B.* 878. *Succisa pratensis* Moench.—St. 1—3 feet high, rarely branched. Radical l. many, stem-l. usually few. Fl. all alike, purplish blue, rarely white. St. and both sides of the l. hairy or glabrous.—Meadows and pastures. P. VII.—X. *Devil's bit*. E. S. I.

2. *S. Columbária* (L.); heads of fr. globose, *involucre membranous notched furrowed throughout*, no distinct base, cor. 5-cleft radiant, radical l. oblong stalked crenate entire or lyrate, uppermost l. pinnatifid with linear segments.—*E. B.* 1311.—St. 12—18 in. high. Radical l. blunt, or, rarely, lanceolate and acute, on long stalks; upper l. rarely entire, linear. Fl. purplish, anth. yellow.—On a calcareous soil. P. VII. VIII. E. S.

[*S. marítima* (L.); involucre and its base furrowed, cor. 5-cleft; has been found near St. Ouen's Bay, Jersey.]

## Order XLV. COMPOSITÆ.

Fl. surrounded by an involucre formed of scales (phyllaries), collected together in a head looking like a single flower. Cal. superior; limb inconspicuous or forming a toothed bristly or feathery pappus. Cor. tubular or ligulate; both kinds in the same head or only one of them. Stam. 5, inserted in the tube; filaments free; anthers united into a tube surrounding the style, often with tails at their base. Fr. indehiscent, dry, with an erect seed without albumen.—No stipules.

The following arrangement of the genera is different from that used in the arrangement of the species. In cases of difficulty they may both be used with advantage.

## Suborder I. CORYMBIFERÆ.

Flowers of the disk tubular and perfect; marginal flowers often ligulate and female or neuter. Style not swollen below its branches.—Juice watery.

A. *Pappus more or less hairlike.*

\* Anthers without bristles at their base.

† *Leaves opposite, cauline.*

- (1.) 1. EUPATORIUM. Heads few-flowered. Phyll. imbricate, oblong. Receptacle naked. *Fl. all tubular-funnel-shaped, perfect (reddish purple).* Anth. included. Branches of the style exerted, cylindrical, blunt.

†† *Leaves radical.*

- (2.) 2. PETASITES. Heads many-flowered. Fem. fl. filiform, obliquely truncate or shortly ligulate, in many rows in the fem. heads, none or in 1 row in the male heads. Male fl. tubular, few and central in the fem. heads, occupying the whole disk in the male heads. Receptacle naked. Phyll. in one row.—*Subdiœcious.* Heads panicled.
- (3.) 3. TUSSILAGO. Heads many-flowered. Fl. of ray narrowly ligulate, fem. in many rows; of disk male, tubular, 5-cleft. Receptacle naked. Phyll. in 1 row, with membranous margins.—Heads solitary.

††† *Leaves alternate, cauline.*

- (4.) 22. SENECIO. Fl. of ray in 1 row, ligulate, fem., rarely 0; of disk perfect, tubular. Inv. cylindrical or conical, of 1 row of equal phyll. not membranous at the margin, with or without smaller scales at its base.
- (5.) 21. DORONICUM. Fl. of ray in 1 row, ligulate, fem.; of disk perfect, tubular. Inv. hemispherical, of 2 or 3 rows of equal phyll. Pappus wanting in the ray.
- (6.) 8. LINOSYRIS. Heads not radiant. Fl. all perfect, tubular (yellow). Receptacle naked, pitted; pits with elevated dentate margins in our plants. Phyll. imbricate. Fr. compressed, silky, without a beak.
- (7.) 4. ASTER. *Fl. of ray fem., ligulate, in 1 row; of the disk perfect, tubular.* Receptacle naked, pitted. Phyll. imbricate and a few scales on the peduncle. Pappus in many rows. Fr. compressed, without a beak.

(8.) 5. ERIGERON. *Fl. of ray* fem., ligulate, in many rows; of the disk mostly perfect, tubular. Receptacle naked. Phyll. imbricate. Pappus in many rows. Fr. compressed, without a beak.

(9.) 7. SOLIDAGO. Pappus in 1 row. Fr. terete. (Fl. all yellow.) Otherwise like *Aster*.

\*\* Anthers with 2 bristles at their base.

† *Receptacle without scales. Fr. cylindrical or tetragonal.*

(10.) 9. INULA. *Fl. of ray* fem., ligulate, rarely subtubular; of disk perfect, tubular. Receptacle naked. Phyll. imbricate in many rows. *Pappus hairlike, uniform, in 1 row.*

(11.) 10. PULICARIA. Phyll. laxly imbricate in few rows. Pappus in 2 rows, outer short cuplike membranous toothed rarely wanting, inner hairlike. Otherwise like *Inula*.

†† *Receptacle without scales or scaly only at the margin. Fr. cylindrical or compressed. Pappus hairlike.*

(12.) 12. GNAPHALIUM. *Fl. all tubular*; outer fem.; central perfect. Receptacle flat, not scaly. Inv. hemispherical, imbricate; phyll. equalling the fl. but not intermixed with them.—Cor. of the fem. fl. often inconspicuous.

(13.) 11. FILAGO. Outer fl. fem., filiform, in several rows; *outermost ones intermixed with the inner phyllaries*; central fl. few, perfect, tubular. *Receptacle conical, scaly at the margin.* Inv. subconical, imbricate; phyll. lanceolate, longer than the flowers.

(14.) 13. ANTENNARIA. Heads subdioecious. *Male fl. tubular*; style almost simple; *pappus clavate*. Fem. fl. filiform. Receptacle convex, not scaly. Inv. hemispherical, imbricate; phyll. coloured at the end.

B. *Pappus 0, or membranous.*

† *Receptacle without scales. Heads radiant. Fl. of the ray fem., ligulate, in 1 row*; of the disk herm., tubular.

(15.) 6. BELLIS. Phyll. in 2 rows, equal, blunt. Receptacle conical. *Fr. compressed. Pappus 0.*

(16.) 17. CHRYSANTHEMUM. Receptacle flat or concave. *Fr. of disk terete, without wings*; of the ray slightly angular or somewhat winged. Pappus 0 or of 3 minute teeth.

(17.) 16. MATRICARIA. Receptacle at length conical. Fr.

angular, not winged. Pappus 0, or a slight membranous border.

†† Receptacle without scales. Heads discoidal.

(18.) 19. ARTEMISIA. Fl. of disk perfect; of the ray fem., slender, in 1 row; or all herm. and tubular. Involucre roundish. Phyll. imbricate. Receptacle naked or hairy. Fr. obovate, with a *small epigynous disk*, without pappus.

(19.) 20. TANACETUM. Fl. as in *Artemisia*. Involucre hemispherical. Phyll. imbricate. Receptacle naked. Fr. oblong, angular, with a *large epigynous disk* (as broad as the fruit), crowned with a slight membranous border.

††† Receptacle scaly. Pappus scale-like.

(20.) 24. GALINSOGA. Heads radiant. Fl. of ray fem., ligulate, in one row; of the disk perfect, tubular. Receptacle conical. Phyll. 4—5, broad, ciliate, in one row. Fr. prismatic, with a pappus of oblong scales.

†††† Receptacle scaly throughout. Pappus 0.

(21.) 15. ANTHEMIS. Heads radiant. Fl. of the ray fem., or neuter, ligulate, in 1 row; of the disk perfect, tubular. Receptacle convex or conical. Phyll. imbricate, of few rows. Fr. terete, or bluntly tetragonal, without pappus, but with a more or less prominent margin.

[ANACYCLUS. Fr. compressed, winged at the edges. Otherwise like *Anthemis*.]

(22.) 14. ACHILLEA. Heads radiant. Fl. of the ray fem., ligulate, short; of the disk perfect, tubular; *tube plane-compressed, 2-winged*. Receptacle nearly flat, afterwards often narrow and lengthened. Inv. ovate or oblong. Phyll. imbricate. Fr. *compressed*, without pappus.

(23.) 18. DIOTIS. Heads discoidal. Fl. all perfect, tubular; tube compressed, with 2 auricles at the base. Receptacle convex, with concave downy-topped scales. Inv. bellshaped. Phyll. imbricate. Fr. compressed, *crowned with the persistent auricled tube of the cor.*; pappus 0.

C. Pappus of 2—5 stiff bristles. Receptacle scaly throughout.

(24.) 23. BIDENS. Heads discoidal, sometimes radiant. Fl. (of the ray neuter, ligulate;) of the disk herm., tubular. Receptacle flat. Phyll. in 2 rows, outer row spreading.

Branches of the style surmounted by short cones. Fr. compressed, angular, rough at the edges; the angles ending in 2—5 hispid bristles.

### Suborder II. CYNAROCEPHALÆ.

Flowers all tubular. Style usually thickish below its branches, which often combine into a more or less perfect cylinder. Involucre imbricate.

a. Basal scar of fr. transverse, rounded. Anthers tailed.

25. SAUSSUREA. Fl. all perfect. *Phyll.* unarmed. Receptacle scaly. Pappus in two rows, outer of short rough bristles, inner feathery.

26. CARLINA. Fl. all perfect. *Outer phyll.* lax, leaflike spinous; inner linear, membranous, coloured and resembling a ray. Receptacle with cleft scales. Pappus in one row branched and feathery, united in a ring below.

27. ARCTIUM. Involucre globose. *Phyll.* ending in hooked points. Receptacle flat, with rigid subulate scales. Fr. compressed, oblong. Pappus short, hairlike, distinct.

b. Fr. attached obliquely. Anthers scarcely tailed.

[\* *Pappus in many rows of different lengths; inner row longest, longer than the fruit.*

28. SERRATULA. Heads dicecious by abortion. *Phyll.* sharp, unarmed. Scales of the receptacle split longitudinally into linear bristles. Fr. compressed, not beaked; basal scar oblique. Pappus persistent.

\*\* *Pappus in many rows, unequal; second row longest, equal to or shorter than the fruit; rarely none.*

29. CENTAUREA. Anthers with papillose filaments. Receptacle chaffy. Fr. attached laterally above the base to the receptacle. Pappus hairlike, rarely 0.

c. Fr. with a transverse basal scar. Pappus in many rows, equal, long. Anth. scarcely tailed.

30. ONOPORDUM. Receptacle honeycombed. Fr. 4-ribbed. Pappus rough. Otherwise like *Carduus*.

31. CARDUUS. *Phyll.* simple, spinous, pointed. Receptacle with fimbriate scales. Fr. compressed, oblong. Pappus long, hairlike or feathery, united into a ring at the base and deciduous.—Includes *Cnicus* Linn. and *Circium* DC.



*d.* Fr. with a transverse basal scar. Pappus in many rows. Filaments monadelphous, glabrous. Anth. scarcely tailed.

32. *SILYBUM*. Phyll. leaflike at the base, narrowed into a long spreading spinous point. Receptacle scaly. Fr. compressed, *its terminal scar surrounded by a papillose ring*. Pappus hair- or scale-like, united into a ring at the base, deciduous.

Suborder III. CICHORIACEÆ or LIGULIFLORÆ.

Flowers all ligulate and perfect.—Juice milky.

\* *Pappus 0. Receptacle without scales.*

33. *LAPSANA*. Heads 8—12-flowered. Phyll. in 1 row, erect, with 4—5 short bracts at their base. Fr. compressed, striate, deciduous, not enveloped in the phyllaries.

\*\* *Pappus like a crown, or of many entire broad scales. Receptacle without scales.*

34. *ARNOSERIS*. Heads many-flowered. Phyll. in 1 row, about 12, keeled, linear-lanceolate, *at length converging*, a few small bracts at their base. Fr. angular, crowned with a short elevated entire margin.

35. *CICHORIUM*. Head many-flowered. Phyll. in 2 rows, outer of about 5, lax, shortish; inner of 8—10, longer, converging, *at length reflexed*. Receptacle sometimes slightly pilose. Fr. obovate, compressed striate. Pappus of 2 rows of minute erect chaffy scales.

\*\*\* *Pappus feathery. Receptacle scaly.*

36. *HYPOCHÆRIS*. Heads many-flowered. Phyll. oblong, imbricate. Fr. glabrous, muricate, often beaked. Pappus in 2 rows, outer short and setaceous, inner long and feathery; or in 1 row and feathery.

\*\*\*\* *Pappus feathery, or on the exterior fruit scaly. Receptacle without scales.*

37. *THRINCIA*. Inv. oblong. Phyll. in 1 row, with a few additional at the base. Fr. beaked. Pappus in 2 rows; outer setaceous, deciduous; inner longer, feathery, dilated at the base. *Marginal row of fruits enveloped in the phyllaries, scarcely beaked, and with a short crown-like pappus.*

38. *LEONTODON*. Inv. subimbricate; exterior phyll. much

smaller, in 1—3 rows. *Fr. uniform*, slightly beaked. *Pappus of all the fr. in 2 rows*; outer setaceous, persistent; inner longer, feathery, dilated at the base; or in 1 row, feathery.

39. TRAGOPOGON. Inv. simple, of 8—10 equal phyll. connected at the base. *Fr. longitudinally striate*, with a long beak; basal scar lateral. *Pappus in many rows, feathery, interwoven in the ray.*

40. PICRIS. Phyll. in 1 row, equal, with unequal linear often spreading ones at the base. *Fr. terete, transversely striate, constricted or slightly beaked above.* *Pappus in 2 rows, feathery; external row rather hairlike.*

41. HELMINTHIA. Phyll. in 1 row, equal, with equal subulate adpressed ones at the base, and surrounded by 3—5 leaflike loose bracts. *Fr. compressed, transversely rugose, rounded at the end and with a slender beak longer than the fruit.* *Pappus in several rows, feathery.*

\*\*\*\*\* *Pappus filiform, deciduous, never feathery nor dilated at the base. Receptacle generally without scales. Fruit compressed.*

42. LACTUCA. Heads few-flowered. Phyll. with a membranous margin, imbricate in 2—4 rows: outer row shorter. *Fr. plane compressed, contracted and prolonged into a filiform beak which is neither crowned nor muricate.*

43. TARAXACUM. Heads many-flowered. *Inv. double*; inner phyll. in 1 row, erect; outer few, short, lax or adpressed, imbricate. *Fr. subcompressed, muricate and suddenly contracted above, prolonged into a filiform beak.*

44. SONCHUS. Heads many-flowered. Phyll. imbricate in 2 or 3 rows, unequal. *Fr. plane compressed, truncate, not beaked.*

45. MUGGEDIUM. Heads many-flowered. *Inv. double*; inner phyll. in 1 row; outer short, lax, imbricate. *Fr. compressed, constricted above, and ending in a ciliate disk.* Outer rows of the pappus rigid and brittle.

\*\*\*\*\* *Fruit terete, ribbed. Otherwise like the preceding section.*

46. CREPIS. Heads many-flowered. *Inv. double*; inner phyll. in 1 row; outer short, lax. *Fr. terete, narrowed upwards or beaked.* *Pappus soft.*

47. HIERACIUM. Heads many-flowered. Phyll. imbricate,

many, oblong. Fr. truncate, not beaked, with a very short crenulate margin. Pappus brittle.

[*Anomalous Genus.* Order AMBROSIACEÆ Link.]

- [48. XANTHIUM. Heads monœcious.—Male: inv. of 1 row of free phyll., many-flowered. Receptacle scaly. Cor. funnelshaped, 5-cleft. Anth. free. Stigma blunt, entire. —Fem. fl. 2, enclosed within the inv. which ends in 1—2 beaks, is covered with hooked spines, and hardens over the fruit. Cor. 0. Stam. 0. Stigmas 2, diverging, linear. Fr. compressed, each occupying a cell in the involucre.]

Suborder I. *TUBULIFLORÆ*. Flowers all tubular, regular, with 4—5 teeth; or the outer ligulate.

Tribe I. *Eupatoriæ*.

Fl. all tubular, regular, perfect. Anth. not tailed. Style-branches blunt, terete, subclavate.

1. EUPATO'RIMUM *Linn.* Hemp-Agrimony.

1. *E. cannab'inum* (L.); l. in 3 or 5 deep lanceolate serrate segments the middle one longest.—*E. B.* 428.—St. herbaceous, erect, striate scabrous, 2—3 feet high. Heads in a fastigiate corymb, 5—6-flowered. Phyll. about 10; 5 exterior short, blunt. Florets reddish purple. L. downy. Herb slightly aromatic.—Banks of streams. P. VIII. IX. E. S. I.

Tribe II. *Tussilagineæ*.

Fl. of ray female, filiform or ligulate; of disk male, tubular. Anth. not tailed. Style-branches connate or short, with conical tips.

2. PETASI'TES *Gaert.* Butterbur.

1. *P. vulgaris* (Desf.); l. roundish-cordate unequally toothed downy beneath with approximate basal lobes, stigmas of the submale fl. short ovate, female fl. truncate obliquely.—*R.* xvi. 901. *Tussilago Petasites* L. *E. B.* 430, 431.—Soboliferous. Panicle long and lax in the female plant, ovoid and dense in the male. Fl. appearing before the l., on stout erect stalks which are clothed with concave tumid petioles either leafless or with a small limb. L. very large, radical, ultimately often 3 feet broad, glabrous above.—Swamps. P. IV. E. S. I.

[*P. frágrans* (Presl), *Sy. E. B.* 781, which has shortly ligulate female fl., is established in some places in the South.—*P. albus* (Gaert.), *Sy. E. B.* 782, with white or cream-coloured fl., and much smaller deeply scalloped l., is established in Scotland.]

### 3. TUSSILA'GO *Linn.* Coltsfoot.

1. *T. Far'fara* (L.).—*E. B.* 429. *R.* xvi. 904.—Soboliferous. Fl. appearing before the l., in bright yellow solitary heads, erect in blossom and seed, drooping before and after flowering; their stalks clothed with scalelike smooth bracts. L. roundish-cordate, angular, toothed, downy beneath.—Moist chalky and clay soils. P. III. IV. E. S. I.

## Tribe III. *Asteroideæ.*

Fl. of ray female or neuter, ligulate (or in *Linosyris* 0); of disk tubular. Anth. not tailed. Style-branches flattened with a subconical tip. Leaves alternate.

### 4. (7.) AS'TER *Linn.* Starwort.

1. *A. Tripólium* (L.); st. glabrous corymbose, l. linear-lanceolate fleshy smooth, involucre imbricate, phyll. blunt membranous the inner ones longer.—*E. B.* 87. *R.* xvi. 907.—St. 1—2 feet high, erect, hollow, leafy, many-flowered. Heads large; disk yellow; rays bright blue, often wanting.—Muddy salt marshes. P. VIII. IX. E. S. I.

[*A. sáignus* (W.) is established in Wicken Fen, Cambridge-shire, and *A. longifolius* at and about Seggieden, Perth; but they are probably both escapes from gardens. Other species are occasionally found.]

### 5. (8.) ERIG'ERON *Linn.* Fleabane.

\*1. *E. canadensis* (L.); st. much branched hairy *panicled many-headed*, l. linear-lanceolate ciliate.—*E. B.* 1019.—St. erect, 1—2 feet high. Heads many, small, yellowish. Inv. cylindrical, scarcely shorter than the fl. of the ray, finally spreading.—Waste ground rare. A. VIII. IX. E.

2. *E. ácris* (L.); st. *corymbose*, branches alternate 1-headed, l. linear-lanceolate entire spreading, lower l. narrowed below, ray erect scarcely longer than the disk, inner female fl. filiform many.—*E. B.* 1158. *R.* xvi. 917.—St. erect, 6—18 in. high, simple below, corymbosely branched above, often several from

one root. Fl. yellow, the ray pale blue.—Dry gravelly places and walls. B. VII. VIII. *Blue Fleabane*. E. S. I.

3. *E. alpinus* (L.); st. mostly with a single head, l. lanceolate, lower l. narrowed below, ray spreading twice as long as the disk, inner female fl. tubular filiform many.—*E. B.* 464. *R.* xvi. 914.—St. 4—8 in. high, usually ending in a solitary head with a yellow disk and light-purple ray. Inv. hairy.— $\beta$ . *E. uniflorus* (Sm. not L.) has a shorter and more erect ray and a rather more hairy involucre. *E. B.* 2416.—Highland mountains. P. VII. VIII. S.

6. (15.) *BEL' LIS* Linn. Daisy.

1. *B. perennis* (L.); l. obovate-spathulate crenate-dentate.—*E. B.* 424.—St. a short procumbent rhizome producing l. only at its end. Stalks simple, each bearing a single head. Sometimes all the fl. are ligulate; rarely all are tubular.—Banks and pastures. P. III.—X. E. S. I.

7. (9.) *SOLIDA'GO* Linn. Golden Rod.

1. *S. Virgaúrea* (L.); st. erect slightly angular, l. lanceolate narrowed at both ends, lower l. elliptic stalked serrate, raceme erect simple or compound, phyll. lanceolate acute, fr. downy.—*E. B.* 301. *R.* xvi. 911.—St. usually 1—3 feet high, leafy, nearly simple, ending in a long cluster of yellow heads.— $\beta$ . *angustifolia* (Koch); l. all lanceolate.— $\gamma$ . *S. cambrica* (Huds.); st. 2—6 in. high, l. ovate-lanceolate, heads larger.—Woods and thickets.  $\gamma$  on mountains. P. VII.—IX. E. S. I.

8. (6.) *LINOSY'RIS* Cand. Goldilocks.

1. *L. vulgaris* (Cass.); herbaceous, l. linear glabrous entire, heads corymbose, inv. lax.—*Chrysocoma Linosyris* L. *E. B.* 2505.—St. 12—18 in. high, simple, leafy. L. single-ribbed, smooth or rough, very many, more or less dotted. Fl. yellow.—Limestone cliffs, rare. P. VIII. IX. E.

Tribe IV. *Inuleæ*.

Heads never dicecious. Fl. of ray female or neuter, ligulate, in one row, or wanting; of disk perfect, tubular. Anth. with slender tails. Style-branches broadened upwards and rounded. L. alternate.

9. (10.) *I'NULA* Linn.

†1. *I. Helénium* (L.); outer phyll. ovate, inner obovate, l. unequally dentate downy beneath cordate-ovate acute clasping,

root-l. stalked elliptic-oblong, fr. quadrangular glabrous.—*Sy. E. B.* 766. *R.* xvi. 921.—St. 3—4 feet high, round, furrowed, solid, leafy, branched above. Heads few together or solitary, terminal, very large; fl. bright yellow, those of the ray ligulate. Phyll. reflexed.—Moist pastures. P. VII. VIII. *Elecampane.*  
E. I.

2. *I. salicina* (L.); outer phyll. oblong-lanceolate, inner linear, l. lanceolate  $\frac{1}{2}$ -clasping tuberculate-serrate scabrous-ciliate glabrous above the edge revolute, st. and underside of l. pilose-hairy.—*Sy. E. B.* 768. *J. of B.* iv. t. 43.—L. with crisped hairs on the veins beneath as also the st., disk of l. beneath pilose. Heads terminal, solitary or 3—5 in a corymb. Rays yellow.—Shore of Lough Derg, Co. Galway. P. VII. VIII. ? I.

3. *I. Conyza* (DC.); outer phyll. lanceolate, inner linear acute, l. ovate-lanceolate downy denticulate, lower l. narrowed into a haft, fl. of the ray tubular-ligulate, fr. terete.—*R.* xvi. 923. *Conyza squarrosa* L. *E. B.* 1195.—St. 1—2 feet high, leafy. Heads corymbose. Phyll. reflexed, leaflike. Fl. yellow, those of the ray deeply divided on the inner side.—Calcareous soils. P. VII.—IX. *Ploughman's Spikenard.* E.

4. *I. crithmoides* (L.); phyll. linear taper-pointed l. fleshy linear blunt or with 3 points, fr. terete.—*E. B.* 68.—St. about a foot high, slightly branched near the top, each branch ending in a solitary head with an orange-coloured disk and yellow rays.—On rocks and in muddy salt marshes by the sea. P. VII. VIII. *Golden Samphire.* E. S. I.

#### 10. (11.) PULICARIA Gaert.

1. *P. vulgaris* (Gaert.); l. lanceolate wavy narrow at the base and somewhat clasping, st. much branched downy, heads lateral and terminal hemispherical with very short rays.—*Inula Pulicaria* L. *E. B.* 1196.—St. 6—12 in. high, leafy. Heads small; fl. yellow. Fr. terete. Outer pappus of small distinct scales.—Moist sandy heaths. A. VIII. IX. E.

2. *P. dysenterica* (Gaert.); l. oblong cordate at the base clasping downy beneath, st. paniced woolly, heads axillary and terminal corymbose, rays much exceeding the disk.—*Inula* L. *E. B.* 1115.—Creeping, floccose. St. 12—18 in. high, leafy. Heads larger than in Sp. I, bright yellow; fr. angular. Outer pappus cuplike, crenulate.—Damp places. P. VIII. IX. E. S. I.

#### Tribe V. Gnaphaliæ.

Heads with fem. marginal fl., or dicecious. Fl. all tubular. Anth. with slender tails. Style-branches broadened upwards and rounded. Pappus hairlike, rarely 0.

## 11. (13.) FILA'GO Linn. Cudweed.

\* *Phyll. not spreading with the ripe fruit, in 2 rows.*

1. *F. german'ica* (L.); cottony, st. proliferous at the summit, l. lanceolate wavy acute, heads obscurely 5-angled half-sunk in wool forming axillary and terminal clusters not surrounded and overtopped by l., phyll. longitudinally folded linear cuspidate with glabrous points.—*E. B.* 946. *F. canescens* Jord.—Grey. St. erect or ascending, 4—12 in. long, usually simple below, bearing a solitary terminal cluster of heads, afterwards producing from just below it 2 erect branches which are again proliferous. Heads 20—40 in each cluster; furrows obscure; l. nearest to the cluster much narrowed upwards and acute. Tips of phyll. yellow, rarely reddish.—Dry fields. A. VII. VIII. E. S. I.

2. *F. apiculáta* (G. E. Sm.); cottony, st. proliferous at the summit, l. all oblong blunt apiculate, heads prominently 5-angled half-sunk in wool forming lateral axillary and terminal clusters surrounded and overtopped by 1—2 blunt l., phyll. boatshaped cuspidate with glabrous points.—*F. lutescens* Jord! *Pl. nov. Fr.* iii. t. 7.—Greenish. Smelling like Tansy. St. mostly erect, with short erect branches below. Heads larger than in Sp. 1, 10—20 in a cluster; furrows deep; cluster often seeming lateral from only 1 branch being produced just below it; l. nearest to the cluster scarcely narrowed upwards, blunt apiculate. Tips of phyll. purple.—Sandy places. A. VII. VIII. E.

3. *F. spathuláta* (Presl); silky, st. proliferous, l. oblong-obovate, heads prominently 5-angled not deeply sunk in wool forming axillary and terminal clusters overtopped by 2—3 acute l., phyll. cuspidate boatshaped with glabrous points.—*F. Jussæi Atl. Fl. Par.* t. 26.—Whitish. St. usually branched from near its base; branches mostly horizontal. Heads larger than those of Sp. 1, 8—15 in each cluster; furrows very deep. Tips of phyll. yellow.—Dry fields. A. VII. VIII. E.

\*\* *Phyll. at length spreading.*

4. *F. min'ima* (Fr.); st. forked, l. linear-lanceolate acute flat adpressed, heads pyramidal in lateral and terminal clusters longer than the leaves, phyll. blunty cottony with glabrous points.—*E. B.* 1157. *Gnaphalium arvense* Willd.—St. slender, erect, 2—6 in. high, branched, the branches forked; or prostrate and spreading. Fl. yellowish in very small heads. Whole plant cottony, greyish.—Dry sandy and gravelly places. A. VI.—IX. E. S. I.

5. *F. gallica* (L.); st. forked, l. linear acute, heads conical in axillary and terminal clusters shorter than the leaves, outer phyll. cottony with bluntish glabrous points gibbous at the base and enclosing the marginal fr.—*E. B.* 2369.—St. 6—8 in. high slender. L. narrowing upwards from the base, upright, afterwards revolute.—Dry gravelly places, very rare. Berechurch, Essex. Bayford, Herts. A. VII.—IX. E.

## 12. (12.) GNAPHA'LIUM Linn.

†1. *G. luteo-âlbum* (L.); st. simple branched at the base slightly corymbose above, heads densely clustered leafless, l. linear-oblong wavy woolly on both sides half clasping, lower l. broader at the end and blunt, upper l. narrowing and acute.—*E. B.* 1002.—Woolly. St. 3—12 in. high, decumbent below, then erect or ascending. Heads collected at the extremity of the stem; inv. straw-coloured; fl. tinged with red.—Sandy fields, now lost in E. Jersey and Guernsey. A. VII. VIII. E.

2. *G. uliginosum* (L.); st. diffuse much branched, heads in terminal dense clusters shorter than the leaves, l. linear-lanceolate cottony on both sides.—*E. B.* 1194.—St. 3—5 in. high, much branched, decumbent or ascending. Heads collected at the extremity of the st. and branches; inv. yellowish brown. Fr. glabrous or hairy.—Wet sandy places. A. VII. VIII.

E. S. I.

3. *G. sylvaticum* (L.); st. simple nearly erect, heads in axillary clusters forming an interrupted leafy spike, l. acute linear-lanceolate, st.-l. narrower.—*R.* xvi. 58. *G. rectum* Sm. *E. B.* 124.—St. 3—24 in. high, upper half constituting the spike. Upper l. very narrow.—Woods and heaths. P. VII.—IX. E. S. I.

4. *G. norvégicum* (Gunn.); st. simple nearly erect, heads in a close terminal leafy spike, l. silky or cottony on both sides lanceolate, st.-l. acuminate mucronate broad.—*R.* xvi. 58. *G. sylvaticum* Sm. *E. B.* 913.—St. 6—12 in. high, spike distinctly terminal. St.-l. broad. Fl. longer in proportion to the inv. Phyll. dark brown.—Highland mountains. P. VIII. S.

5. *G. supinum* (L.); caespitose, st. decumbent, flowering st. erect, heads 1—5 distant, l. linear downy on both sides mostly radical.—*E. B.* 1193.—Height 2—3 in. Caespitose, very leafy at the root. Flowering st. with few leaves, which are downy on both sides. Heads sessile forming a sort of capitate spike, or stalked subracemose, or solitary.—Highland mountains. P. VII. S.

## 13. (14.) ANTENNA'RIA R. Br.

1. *A. dioica* (Gaert.); shoots procumbent, flowering st. simple erect, corymb dense terminal, phyll. oblong dilated upwards



blunt coloured, root-*l. obovate spatulate* glabrous above cottony beneath, stem-*l. nearly equal linear-lanceolate* adpressed.—*Gnaphalium* Sm. *E. B.* 267.—St. prostrate, woody, ending in a tuft of many *l.* and producing prostrate leafy stoles. Flowering st. 4—8 in. high, quite simple, cottony. Heads 4—5, erect, slightly stalked. Phyll. white or rose-colour.—*β. A. hyperborea* (D. Don); *l.* cottony on both sides. *E. B. S.* 2640.—Mountain heaths. P. VI. VII. *Cat's-foot.* E. S. I.

\*2. *A. margaritacea* (R. Br.); *st. erect* corymbosely branched above leafy, *l. linear-lanceolate* acute cottony below, heads in level-topped corymbs.—*E. B.* 2018.—St. 2—3 feet high, cottony. *L.* alternate, slightly cottony above, densely beneath. Inv. white. Fl. yellowish.—Moist meadows, rare. Established by rivers in Monm., Glam., and Merionethshires. P. VIII. E.

### Tribe VI. *Anthemideæ.*

Heads usually radiant, fl. of ray fem. or neuter, ligulate or slender and tubular; of disk tubular. Anth. not tailed. Style-branches truncate. Pappus often wanting or crownlike, rarely formed of scales or slender hairs.

#### 14. (22.) ACHILLE'A Linn. Yarrow.

1. *A. Ptar'mica* (L.); *l. shining linear-lanceolate* attenuate acute glabrous smooth uniformly and finely serrate, teeth adpressed mucronate minutely scabrous at the margin, ray 8—12-flowered equalling the involucre, corymb compound.—*E. B.* 757. *R.* xvi. 1024.—St. about 2 feet high, slightly branched above, erect, leafy, angular, smooth. Phyll. with a dark-brown membranous margin. Limb of the radiant florets longer than broad, white. Disk broad, white. *L.* sometimes very narrow; lower teeth not deeper than the others.—Moist meadows and thickets. P. VII. VIII. *Sneezewort.* E. S. I.

[*A. decolorans* (Schrad.); *l.* opaque bluntish downy thickly dotted coarsely and doubly serrate with spreading teeth lacinate and radiating at the base, ray 5- or 6-flowered about equalling the involucre.—*A. serrata* Sm. *E. B.* 2531.—Not known except in gardens. P. IX.]

[*A. tomentosa* (L.); *l.* with a linear-lanceolate outline pinnatifid woolly, lobes crowded linear acute, trifid in the lowermost leaves, 2—3-fid in the intermediate, uppermost simple, corymb repeatedly compound, ray equalling about half the involucre.—*E. B.* 2532.—St. 10—12 in. high, decumbent at the base, woolly, simple. Phyll. woolly, edged with brown. Disk and rays golden yellow.—Scarcely naturalized. P. VII. VIII.]

2. *A. Millefolium* (L.); l. with a lanceolate outline bipinnatifid woolly or nearly glabrous, lobes cut with linear segments, rachis entire or subdentate with entire teeth, corymb dense, rays equaling about half the involucre.—*E. B.* 758. *R.* xvi. 1024.—St. erect, 6—18 in. high, nearly glabrous or woolly. Phyll. nearly glabrous with a brown margin. Heads small. *Fl.* white, occasionally reddish or purple.—Pastures and waste ground. P. VI.—VIII. *Yarrow. Millefoil.* E. S. I.

[*A. tanacetifolia* (All.); stem-l. with an oblong outline bipinnatifid woolly or nearly glabrous, rachis winged serrate, pinnæ pinnules and teeth of the rachis incise-serrate, corymb dense, rays equaling about half the involucre.—*Sy. E. B.* 728. *R.* xvi. 1027.—St. 3—4 feet high, erect, woolly. Root-l. stalked, with a lanceolate outline; rachis with large incise-serrate teeth especially just under the pinnæ; pinnæ and pinnules broad.—An escape. P. VII.] E.

15. (21.) AN'THEMIS *Linn.* Chamomile.

\* *Scales of the receptacle lanceolate or oblong abruptly ending in an acute rigid point.*

[*A. tinctoria* (L.); receptacle hemispherical, fr. tetragonal crowned with a membranous undivided border, l. bipinnatifid downy beneath, segments parallel decurrent serrate.—*E. B.* 1472.—St. 1—2 feet high, much branched, cottony. Heads on long stalks, solitary, terminal; disk and rays bright yellow. Scales not protruding.—Fields. Not a native. B. ? VII. VIII.]

1. *A. arven'sis* (L.); receptacle conical, fr. tetragonal, l. bipinnatifid hairy, segments linear-lanceolate.—*E. B.* 602. *R.* xvi. 1004.—St. 1—2 feet high, striate, downy, much branched. Segments of the l. parallel and at length converging. Heads on long stalks, solitary, terminal; disk convex, bright yellow; ray white, always having styles. Scales just appearing above the fl. of the disk, lanceolate. Outer fr. crowned with a tumid plicate-rugose ring, inner with an acute margin.—Borders of cultivated fields, rare. A. VI. VII. *Corn-Chamomile.* E. S. I.

[*A. ang'lica* (Spr.); "receptacle flat," fr. crowned with a very narrow entire border, l. pinnatifid somewhat hairy, lobes incise-serrate acute bristle-pointed rather fleshy.—*A. maritima* Sm. *E. B.* 2370.—Probably a maritime form of *A. arvensis*.—Seashore. Sunderland. Not recently found. A. VII.] E.

\*\* *Scales of the receptacle linear setaceous acute.* MARUTA Cass.

2. *A. Cot'ula* (L.); receptacle long conical, fr. terete tubercular-striate crowned with a crenulate margin surrounding a

slightly convex disk, l. bipinnatifid nearly glabrous, lobes linear acute mostly entire.—*E. B.* 1772. *R.* xvi. 1000.—St. 1—2 feet high, branched, angular, furrowed. Heads solitary on long terminal stalks; scales confined to the central part of the receptacle; disk yellow; ray white, usually without styles. Cor.-tube 2-winged. Phyll. blunt, with white membranous margins. Fetid and acrid.—Fields and waste places. A. VII.—IX. E. S. I.

\*\*\* *Scales of the receptacle thin membranous blunt.*

3. *A. nob'ilis* (L.); receptacle conical, fr. subtrigonus, smooth, l. bipinnate, leaflets linear-subulate slightly downy rather fleshy acute.—*E. B.* 980. *St.* 27. 15.—*St. procumbent*, 1 foot long, much branched. Heads solitary, terminal; disk yellow; ray white. Pleasantly aromatic. Gravelly and sandy places. P. VII. VIII. *Chamomile*. E. S. I.

[*Anacy'chus radiátus* (Lois.); l. bipinnatifid rather woolly, lobes linear-subulate, outer phyll. hairy on the back, inner blunt dilated diaphanous above.—Head solitary; ray yellow.—Dunboy Quay, Bearhaven, Cork. Accidental. A. VII.] I.

16. (17.) MATRICA'RIA Linn. Feverfew.

‡1. *M. Parthénium* (L.); l. stalked pinnate, lts. ovate or oblong pinnatifid, lobes cut, st. branched, heads corymbose, phyll. linear blunt, receptacle convex, fr. crowned with a short jagged membrane.—*Pyrethrum* Sm. *E. B.* 1231.—St. erect, 2 feet high, branched, furrowed, paniced. Heads in small corymbs terminating the stem and branches; disk yellow; ray white.—Waste places, not very common. P. VII. VIII. *Feverfew*. E. S.

2. *M. inodóra* (L.); l. sessile pinnatifid with many capillary pointed segments, st. branched, heads solitary, phyll. lanceolate blunt, receptacle ovate, fr. rugose and with 2 glandular spots on the external face just below the elevated entire border.—*Pyrethrum* Sm. *E. B.* 676.—St. erect, 12—18 in. high, smooth, angular. L. in very narrow mostly alternate segments. Heads solitary, ending the branches; margin of phyll. cut and fuscous; ray white; disk yellow. Base of the invol. turbinate afterwards truncate; recept. hemispherical afterwards conical, much longer than broad. Fr. with 3 prominent smooth ribs.— $\beta$ . *sabina*; l.-segm. short fleshy linear bluntish convex above, principal ribs keeled beneath, inv. and recept. as in typical plant. *E. B.* 979. L. with short crowded mostly opposite segments.— $\gamma$ . *M. maritima* (L.); l.-segm. long, base of inv. subumbilicate, recept. as broad as long. Fr. larger. Margin of phyll. broadly and darkly coloured.—Fields and waste places.  $\beta$ . Sea-shore.  $\gamma$ . Sea-shore in the north of Scotland. A. VII. VIII. E. S. I.

3. *M. Chamomilla* (L.); l. bipinnate smooth, segments capillary simple or divided, heads solitary or subcorymbose, *receptacle hollow conical*, phyll. linear blunt.—*E. B.* 1232.—St. erect, 1 foot high, branched. Heads on long naked stalks or forming an irregular corymb; disk yellow; ray white.—Cultivated and waste ground. A. VI. VII. *Wild Chamomile.* E. I.

17. (16.) CHRYSAN'THEMUM Linn.

1. *C. Leucanthemum* (L.); lower l. obovate stalked, stem-l. oblong blunt cut sessile pinnatifid at the base, *phyll.* lanceolate blunt with a narrow dark purple membranous margin, fl. of ray white, fr. without a border.—*E. B.* 601. St. 2. ll.—St. erect, 1—2 feet high, simple, striate. Lower l. narrowing into a winged and auricled stalk. Heads solitary, terminal, large; disk yellow.—Fields. P. VI.—VIII. *Ox-eye.* E. S. I.

†2. *C. seg'etum* (L.); l. glabrous toothed dilated outwards and lobed, upper l. clasping, *phyll.* ovate blunt, with a broad membranous margin, fl. of ray yellow.—*E. B.* 540.—St. a foot high, alternately branched, angular. L. incise-serrate or lobed in the upper part, simply toothed below. Heads solitary, terminal.—Corn-fields. A. VI.—VIII. *Corn-Marigold.* E. S. I.

18. (23.) DIO'TIS Desf. Cotton-weed.

1. *D. maritima* (Cass.).—*E. B.* 141.—Densely cottony and white. St. about a foot long, decumbent below, densely leafy, corymbose above. L. sessile, oblong, blunt, flat, crenate, persistent. Heads in terminal corymbose tufts. Inv. cottony. Fl. yellow.—Sandy sea-shores, rare. P. VIII. IX. E. I.

19. (18.) ARTEMIS'IA Linn. Wormwood.

\* *Receptacle pilose.*

1. *A. Absinthium* (L.); heads drooping hemispherical, fl. not all perfect, l. silky in many deep lanceolate blunt segments, outer phyll. linear silky, inner roundish scarious.—*E. B.* 1230.—St. bushy, 1—2 feet high. Heads in erect leafy panicles. Floral l. simple. Fl. dull yellow, the outer row female.—Waste ground. P. VII. VIII. *Wormwood.* E. S. I.

\*\* *Receptacle naked.*

2. *A. campes'tris* (L.); heads drooping ovate glabrous, fl. not

all perfect, *l. silky* with many linear-lanceolate mucronate segments, stem-*l.* once or twice pinnate with linear segments, st. wandlike procumbent before flowering, *phyll. ovate glabrous with a scarious margin.*—*E. B.* 338.—Barren st. cæspitose. Flowering st. slender, 1—2 feet long, ascending when the flowers appear, leafy, smooth. Fl. yellow, those of disk sterile; inv. purplish.—Sandy heaths in Norf. and Suff., rare. P. VIII. IX. E.

3. *A. vulgáris* (L.); heads ovate, fl. not all perfect, *l. woolly* and white beneath pinnatifid with lanceolate acuminate cut and serrate segments, *phyll. woolly.*—*E. B.* 978.—St. 2—3 feet high, erect, leafy. Clusters leafy, nearly simple, erect. Fl. few, reddish or brownish yellow.—*a*; *l.*-segm. oblong, racemes open.—*β.* *A. coarctata* (Forcell.); *l.*-segm. linear-lanceolate, racemes subspicate condensed.—Waste ground. P. VII.—IX. *Mugwort.*  
E. S. I.

4. *A. marit'ima* (L.); heads oblong, *florets few all perfect, l. downy pinnatifid with linear blunt segments,* *phyll. oblong outer woolly inner scarious.*—*E. B.* 1706.—St. procumbent or ascending, woolly, much branched. Fl. reddish yellow. Racemes drooping.—*β.* *A. gallica* (Willd.); racemes more dense, heads erect. *E. B.* 1001.—Salt marshes. P. VIII. IX. E. S. I.

[*A. cærules'cens* (L.); heads oblong, fls. few all perfect, *l. hoary lanceolate undivided pinnatifid* on the barren shoots.—*E. B.* 2426.—Not a native. P. VIII. IX.]

## 20. (19.) TANACE'TUM Linn. Tansy.

1. *T. vulgáre* (L.); *l. bipinnatifid, lts. serrate.*—*E. B.* 1229.—St. 2—3 feet high, leafy. Heads in a terminal corymb. Fl. golden yellow. Fr. with an entire crown. St. 2—3 feet high.—Waysides. P. VIII. E. S. I.

## Tribe VII. Senecioneæ.

Fl. of ray fem. ligulate; of disk tubular. Anth. tailless. Style-branches truncate or slightly rounded. Pappus hairlike.

## 21. (5.) DORONICUM Linn. Leopard's-bane.

†1. *D. Pardalian'ches* (L.); *l. cordate denticulate, lowermost l. on long stalks, intermediate with clasping auricles at the base of*

*the haft*, uppermost sessile clasping.—*E. B. S.* 2654.—St. 2—3 feet high, erect, solitary, hollow, hairy. L. hairy, minutely toothed, soft, blunt, the uppermost acute. Lowest petioles not auricled. Heads several, phyll. lanceolate-subulate. Fl. yellow. The earlier heads overtopped by the later ones. Fr. oblong, furrowed, of disk hairy, of ray glabrous.—Damp and hilly woods and pastures, rare. P. V.—VII. E. S.

†2. *D. plantagin'eum* (L. ?); *l. ovate* denticulate, radical on long stalks rounded or subcordate produced at the base, *stem-l. sessile* clasping the lowermost with a winged and auricled stalk.—*Sy. E. B.* 762.—Crown of the root woolly. St. 2—3 feet high. Stem-l. narrowed in their lower half but sessile, uppermost with a long taper point. Heads usually solitary, or, if more, the lateral ones not overtopping the terminal one. Phyll. subulate. Fr. of ray glabrous. Fl. yellow.—Damp places, rare. P. VI. VII. E. S.

## 22. (4.) SENE'CIO Linn. Ragwort.

A. Involucre with small scales at its base.

\* *Fl. all tubular, or marginal ones ligulate but mostly revolute.*

1. *S. vulgáris* (L.); *l. half-clasping* pinnatifid, segments distant oblong blunt and together with the rachis and auricles acutely and unequally toothed, lower *l. narrowed* into a stalk, heads in clustered racemes, *outer phyll. very short* adpressed with black points, ray 0.—*E. B.* 747.—Smooth or woolly. Not viscid. St. 6—12 in. high, branching. Heads small, involucre oblong-conical, glabrous; fl. yellow; fr. silky.—Rarely there is a single row of ligulate minute revolute marginal flowers. A form with long prominent rays has been supposed to be *S. crassifolius* (W.).—Common. A. I.—XII. *Groundsel.* E. S. I.

2. *S. sylvat'icus* (L.); *l. deeply pinnatifid* downy, segments oblong unequally toothed, *heads corymbose*, involucre downy, *outer phyll. very short glabrous* (or slightly downy), ray small revolute, fr. silky.—*E. B.* 748.—Slightly viscid. St. 1—2 feet high, erect, more or less branched, hairy. L. narrower than in Sp. 3. Inv. conical; fl. yellow.—*S. lividus* (Sm.) is a slight var. with the upper *l. more distinctly auricled and clasping*.—Dry and gravelly places. A. VII.—IX. E. S. I.

3. *S. viscósus* (L.); *l. deeply pinnatifid viscid glandular-hairy*, segments oblong unequally toothed and lobed, heads in an irregular corymb, *involucre viscid, outer phyll. half the length of the*

inner hairy, ray small, fr. glabrous.—*E. B.* 32. *Very viscid.* St. 1—2 feet high, much branched, spreading. Heads on long stalks; inv. cylindrical; fl. yellow.—Waste ground, rare. *A. VII.—IX.* E. S. I.

\*\* *Heads with spreading rays. Leaves pinnatifid.*

[\**S. squalidus* (L.); l. pinnatifid glabrous, segments linear or oblong distant toothed irregular, heads loosely corymbose, involucre glabrous, outer phyll. few small, fr. silky.—*E. B.* 600. *S. chrysanthemifolius* DC.—St. much branched, leafy, smooth. L. sessile, often auricled, deeply and irregularly lobed. Heads few, broad. Outer phyll. very small, sometimes very few. Many awlshaped scattered bracts below the heads. Fl. yellow.—Walls. Oxford. Bidford, Devon. Cork. *A. VI.—X.*] E. I.

4. *S. erucifolius* (L.); l. pinnatifid margins somewhat revolute cottony beneath, lower l. stalked, segments linear the lowermost smallest entire and clasping the stem, outer phyll. half as long as the inner, ribs of all the fr. silky.—*S. tenuifolius* Sm. *E. B.* 574.—Creeping slightly. St. erect, 2 feet high, angular, furrowed, somewhat cottony, simple. Lower l. oblong-ovate, deeply pinnatifid, cottony especially beneath; segments often linear. Fr. each having a persistent pappus. Fl. yellow. When the l. are divided into very narrow segments it is *S. tenuifolius* Jacq.—Calcareous soils. *P. VII. VIII.* E. S. I.

5. *S. Jacobæ'a* (L.); l. glabrous, lower l. oblong-ovate attenuate below lyrate-pinnatifid stalked, stem-l. sessile bipinnatifid, segments spreading oblong deeply and irregularly toothed and cut, lowermost much divided clasping, outer phyll. scattered few, fr. hairy those of the ray glabrous.—*E. B.* 1130.—Root fleshy. St. 2—3 feet high, smooth, striate, branched, leafy. Corymb with erect branches. Fr. of the ray with deciduous pappus. Fl. yellow. Ray sometimes wanting.—Waste ground. *P. VII.—IX.* *Ragwort.* E. S. I.

6. *S. aquat'icus* (Huds.); l. glabrous, lower l. stalked crenate or dentate obovate or oblong slightly prolonged at the base undivided or sublyrate blunt, upper l. lyrate or pinnately cut, segments oblong or linear, st. round corymbosely branched, fr. all glabrous.—*E. B.* 1131.—St. erect, 1—4 feet high, simple or branched in the upper half, branches ascending. Terminal lobe of the lower l. rounded below and narrowed into its stalk. Fl. yellow.—A larger much branched form has l. all lyrate, term. lobe truncate or subcordate below, segm. subspathulate.—In marshy places. *P. VII. VIII.* E. S. I.

\*\*\* *Heads with spreading rays. Leaves undivided.*

7. *S. paludósus* (L.); l. sessile long lanceolate tapering *sharply serrate cottony beneath*, st. straight hollow, corymbs terminal.—*E. B.* 650.—St. 4—6 feet high, somewhat woolly. Fl. yellow; those of the ray narrow, 13—16.—Fen ditches, very rare. P. V.—VII. E.

8. *S. saracen'icus* (L.); l. sessile lanceolate acute *glabrous irregularly serrate* especially the uppermost, st. straight solid, corymbs terminal, ray of 6—7 flowers.—*E. B.* 2211. *S. salicetorum* Godr.—St. 3—5 feet high, smooth. L. broad. Corymb many-headed. Fl. yellow.—There may be a second plant here with l. broadest below their middle very finely toothed, uppermost nearly entire. Its l. are much prolonged and rather glaucous. It closely resembles *S. Doria* Jacq.—Watery places, local. P. VIII. E. S. I.

B. Involucre without scales at its base. Heads with a spreading ray. Leaves nearly entire. CINERARIA L.

9. *S. palus'tris* (DC.); shaggy, st. *much branched* and corymbose above, l. *broadly lanceolate half-clasping*, lower l. *sinuate-dentate*.—*E. B.* 151.—St. 3 feet high, thick, hollow, leafy. Heads erect. Fl. bright yellow.—Fen ditches, now become very scarce. P. VI. VII. E.

10. *S. campes'tris* (DC.); shaggy, st. *simple, rt.-l. oblong* nearly entire narrowed below, *stem-l. lanceolate*, heads corymbose, involucre woolly below nearly glabrous in the upper half, fr. hispid.—*E. B.* 152.—St. 6—8 in. high, with small st.-leaves. Heads erect, 1—6, in a simple corymb. Involucre often almost glabrous, pale. Fl. yellow. In very wet seasons it is often thrice as large throughout.—Chalk Downs. P. ? VI. E.

11. *S. spathulifólius* (DC.); shaggy, st. *simple, rt.-l. ovate-spathulate* arachnoid above more woolly beneath, st.-l. *ovate-oblong* narrowed into broadly winged petioles, upper sessile linear or lanceolate clasping, invol. woolly, fr. hispid.—*R.* xvi. 978. *Schultz Herb. Norm.* 690!—St. 1—3 ft. high, with large clasping st.-leaves often much enlarged at the base. Fl. yellow.—Near Holyhead. P. or B. VI. VII. E.

### Tribe VIII. *Helianthoideæ.*

Heads discoid with all the fl. similar and perfect; or with fem. radiant ligulate flowers. Anth. without tails or scarcely tailed. Style-branches truncate. Pappus of 2—4 bristles, or of oblong scales.



23. (24.) *BIDENS* Linn.

1. *B. tripartita* (L.); l. stalked 3-partite, segments lanceolate serrate, fr. obovate-cuneate usually with 2 bristles.—*E. B.* 1113. *R.* xvi. 941.—St. 1—3 feet high, with opposite branches. L. narrowed into winged footstalks, sometimes undivided, sometimes pinnate-5-fid. Heads terminal, solitary, suberect. Fl. brownish yellow.—Marshy places. A. VIII. IX. E. S. I.

2. *B. cer'nea* (L.); l. sessile connate lanceolate undivided serrate, fr. cuneate usually with 3—4 bristles.—*E. B.* 1114. *R.* xvi. 941.—St. 1—3 feet high, with opposite branches. L. simple narrowed below but not stalked. Heads terminal, solitary, drooping. Fl. brownish yellow.—Sometimes radiant marginal fl. are found.—Watery places, A. VIII. IX. E. S. I.

24. (20.) *GALINSO'GA* R. and P.

\*1. *G. parviflora* (Cav.); subglabrous, receptacle conical, pappus of 8—16 scales.—*Sy. E. B.* 765. *R.* xvi. 983.—St. 1—2 feet high; branches opposite. L. opposite, ovate, stalked. Fl. of ray few, broadly ligulate, short, white; of the disk about as long as the phyllaries, yellow.—A South-American plant escaped from Kew Gardens, and quite naturalized in many places. A. VII.—IX. E.

Tribe IX. *Cynarocephalæ*.

Fl. all tubular. Style of the perfect fl. thickened and often with a tuft of hairs below the branches, which are united or free and downy externally.

Section 1. *CARLINEÆ*. Heads many-flowered, never diœcious. Phyllaries in many rows, distinct, often spinous. Filaments distinct, naked. Fr. mostly villose. Pappus in 1—2 rows, not surrounded by an elevated margin.

25. *SAUSSU'REA* Cand.

1. *S. alpina* (DC.); l. nearly glabrous above cottony beneath, lower ones ovate-lanceolate, upper sessile lanceolate, all distantly toothed, heads few in a dense corymb, involucre subcylindrical, phyll. adpressed hairy.—*Serratula* L. *E. B.* 599.—St. 3—12 in. high, erect, downy, simple, ending in a small corymb of heads with pinkish fl. and purple anthers. Fl. scented like Heliotrope. Fr. glabrous.—In alpine situations. P. VIII. E. S. I.

26. *CARLINA* Linn.

1. *C. vulgaris* (L.); st. corymbose one- or many-headed, l. oblong-lanceolate sinuate spinous, outer phyll. bipinnatifid spinous,

inner linear-lanceolate attenuate acute ciliate in the lower half, bracts shorter than the heads.—*E. B.* 1144.—St. 6—12 in. high, usually cottony, leafy. Spines many, short. Root-l. lanceolate or linear-lanceolate. Underside of the l. and phyll. often cottony. Heads large; inner phyll. cream-coloured (a false ray); anth. yellow.—Dry sandy heaths. *B. VII.*—*X.* *E. S. I.*

### 27. *ARC'TIUM* Linn. Burdock.

1. *A. május* (Schk.); heads loosely subcorymbose long-stalked hemispherical and open in fr. glabrous (green), phyll. equalling or exceeding fl. subulate inner row shorter than the others, subcylindrical upper part of fl. more than  $\frac{1}{2}$  as long as the lower part.—*Sy. E. B.* 699.—St. 3—4 ft. high, centre and usually most of the branches ending in corymbs. L. broadly cordate-ovate, blunt; petioles solid with prominent angles, deeply furrowed. Heads very large, a few of the lower sometimes with short stalks. Fl.-heads not umbilicate. Fr. yellowish, ultimately dark brown, irregularly rugose.—A form with more spherical and webbed heads (*A. tomentosum* Bab.<sup>1</sup>) is common near Cambridge and is the *L. major* v. *subtomentosa* Lange!—Waste places. *B. VIII.* *E. S. I.*

2. *A. nemorósum* (Lej.); heads racemose subsessile ovate and contracted at the mouth in fr. slightly webbed, phyll. equalling or exceeding the fl. subulate inner row lanceolate shorter than the others, subcylindrical upper part of fl. as long as the lower part.—*Sy. E. B.* 701. *A. intermedium* Bab.—St. 2—4 feet high. L. convolute, cordate, oblong-ovate, petioles hollow, rather angular, nearly flat above. Heads all nearly sessile, less than in *Sp. 1.*, three usually placed close together at the end of each branch, ovate-prolonged when young, not umbilicate. Most of the phyll. ascending.—Local? *B. VIII.* *E. S.*

3. *A. minus* (Schk.); heads racemose shortly stalked globular slightly contracted at the mouth in fr. slightly webbed (greenish), phyll. falling short of the fl. subulate inner row equalling the others and gradually subulate, subcylindrical upper part of fl. about as long as the lower part.—*Sy. E. B.* 702. *Fl. Dan.* 2662. *R. xv.* 811.—Smaller than either of the preceding. Central st. mostly nodding and as well as the branches having scattered small heads; term. head solitary. L. deeply cordate-prolonged; petioles hollow, slightly angular, nearly round, scarcely furrowed. Fl.-heads not umbilicate. Fr. fuscous with black blotches.—Waste places. *B. VIII.* *E. S. I.*

4. *A. intermedium* (Lange); heads subracemose stalked hemispherical and open in fr. much webbed (greenish), phyll. equalling the fl. subulate inner row about equalling the others and

<sup>1</sup> The true *A. tomentosum* is not a native of England.

gradually subulate, subcylindrical upper part of fl. as long as the lower part.—*Sy. E. B.* 700. *Fl. Dan.* 2663. *R.* xv. 812. *A. pubens* Bab.—St. about 3 feet high. L. deeply cordate-prolonged; petioles hollow, scarcely angular, slightly furrowed. Stalks of the heads rather long, those of the lower heads longest. Heads usually with much wool, twice as large as those of *A. minus*. Fr. dark brown with a few paler spots towards the top.—Waste places. B. VIII. E. I

Section 2. *SERRATULÆ*. Heads many-flowered; flowers all tubular, perfect or diœcious, the external row sometimes female. Involucre of many rows of distinct phyllaries. Filaments distinct. *Pappus in many rows of different lengths, inner row longest, hairlike or feathery, surrounded by a margin.*

### 28. SERRAT'ULA Linn. Saw-wort.

1. *S. tinctoria* (L.); l. with bristly serratures pinnatifid or lyrate, heads oblong, phyll. ovate adpressed, inner ones linear coloured.—*Sy. E. B.* 708. *St.* 3. 16—St. 2—3 feet high, straight, erect, angular, branched above. L. rarely entire. Fl. purple.— $\alpha$ ; heads stalked in a lax corymb.— $\beta$ . *S. monticola* (Bor.); heads subsessile few large close together. *E. B.* 38.—Groves and thickets. P. VIII. E.

Section 3. *CENTAUREÆ*. Heads many-flowered, discoidal; outer row of fl. usually barren, enlarged, irregular. Phyllaries in many rows. Filaments distinct. *Pappus in many rows of different lengths, second row longest, setaceo-pilose, placed within the margin which surrounds the epigynous disk, rarely 0.*

### 29. CENTAU'REA Linn. Knapweed.

\* *Phyll. with a scarious pectinate not decurrent appendage.*

[*C. Jacea* (L.); *phyll.-appendages* erect rounded, pappus 0, l. linear-lanceolate lower l. broader and toothed.—*E. B.* 1678.—Outermost phyll. with deeply pinnatifid appendages, few innermost entire, the rest irregularly jagged. Heads radiant.—Not native. P. VIII. IX.]

1. *C. nigra* (L.); *phyll.-appendages* patent or erect lanceolate or ovate-lanceolate pectinate their teeth subulate, pappus 0 or short deciduous, l. lanceolate or ovate-lanceolate acute.—St.  $\frac{1}{2}$ —2 ft. high. L. green; lower narrowed into long stalks, entire or sinuate-dentate; upper sessile. Heads globose. Few innermost phyll.-appendages torn. Fr. oblong, downy.— $\alpha$ . *C. nigra* (L.); *phyll.-appendages* erect or patent ovate-lanceolate usually quite covering the phyll. their teeth long. *E. B.* 278. *Mart. Fl. Rust.* 130. *R.* xv. 761. St. usually with long 1-headed

branches. Pappus 0 or short. Heads sometimes radiant. Autumnal forms have erect-patent branches, ending in solitary heads; vernal have almost divaricate branches.— $\beta$ . *C. decipiens* (Thuill.); phyll.-appendages erect lanceolate or ovate-lanceolate usually not wholly covering the phyll. their teeth short. St. usually simple, 1-headed. L. broader than those of  $\alpha$ . Teeth often scarcely longer than the breadth of the brown appendage; 3 inner rows of phyll. usually protruding. Pappus 0. Heads usually (perhaps always) radiant. *Sy. E. B.* 707. *C. nigrescens* (Bab.). The plant when seen is easily distinguishable from the radiant form of *C. nigra*, although hardly to be separated by characters.—Meadows and pastures.  $\beta$ . South of England, rare. P. VI.—IX. E. S. I.

\*\* *Phyll. lanceolate, their upper half with a somewhat scarious deeply toothed or fringed decurrent margin.*

2. *C. Cy'anus* (L.); phyll. erect adpressed deeply toothed, pappus as long as the fruit, *l. linear-lanceolate*, the lowermost toothed or pinnatifid.—*E. B.* 277.—St. 1—3 feet high, loosely cottony, leafy. L. slightly cottony above, densely beneath. Involucre greenish yellow; phyll. often tinged with purple in their upper half, margins brown decurrent with whitish teeth. Heads with large radiant *blue flowers*, disk purple.—Corn-fields. A. VI.—VIII. *Corn Bluebottle.* E. S. I.

3. *C. Scabiōsa* (L.); phyll. erect adpressed, the triangular-ovate black pectinate appendages not covering the inv., teeth ascending setaceous short, pappus as long as the fruit, *l. pinnatifid* roughish, segments lobed with hard points.—*E. B.* 56.—St. 2—3 feet high, rough, furrowed. L. hispid, lobes of the upper ones entire. Heads on long naked stalks, solitary. Involucres usually rather woolly; phyll. pale, with dark acute membranous pectinate decurrent appendages; teeth paler, short, not longer than  $\frac{1}{2}$  the width of the phyllary. Fl. purple, outer row radiant or 0. Rarely the inv. is quite covered by the appendages.—Fields and hedges. P. VII.—IX. *Great Knapweed. Matfellow.* E. S. I.

[*C. paniculāta* (L.); phyll. erect adpressed rigid with subulate teeth and a short term. rigid point innermost narrow long toothed at the end, pappus much shorter than the fruit, lower *l. pinnatifid* with linear segments.—*R.* xv. 780.—St. about a foot high, paniced above, rough, rather cottony. Heads cylindrical-oblong. Fl. purplish.—Quenvais and St. Ouen's Bay, Jersey. B. VII.]

\*\*\* *Phyll. horny at the end, spines palmate or pinnate.*

†4. *C. solstitialis* (L.); phyll. woolly palmately spinous, central

*spine* of the intermediate ones *very long* needle-shaped, inner phyll. with a roundish scarious appendage, *heads terminal* solitary, st. winged with the decurrent bases of the linear-lanceolate *entire* hoary leaves, root-l. lyrate—*E. B.* 243.—St. 1—2 feet high, branched, spreading. Involucres sometimes glabrous. Fl. yellow.—Cultivated land, probably introduced. A. VII.—IX. *Yellow Star-Thistle.* E.

5. *C. Calcitrapa* (L.); *phyll. glabrous* palmately spinous, *central spine strong channelled*, innermost phyll. with a scarious blunt appendage, *heads lateral* sessile solitary, pappus 0, 1. deeply *pinnatifid*, lobes of the root-l. lanceolate toothed, of the stem-l. linear.—*E. B.* 125.—St. furrowed, slightly hairy, branched, spreading, about a foot high. Fl. purplish.—Gravelly and sandy places. A. VII. VIII. *Common Star-Thistle.* E.

[*C. aspera* (L.); *phyll. palmately spinous, spines nearly equal* 3—5, innermost phyll. with a scarious blunt lanceolate or slightly spatulate appendage, *heads terminal* solitary, pappus of all the fr. in several rows, *l. linear* coarsely toothed narrowed below sessile rough, lower l. (and those of the primary stem?) broader incise-dentate with clasping auricles.—*C. Isnardi* L. *E. B.* 2256.—St. procumbent; branches long slender simple leafy. L. slightly toothed or entire. Fl. purple.—Typical *C. aspera* has its upper leaves sessile but not clasping.—Guernsey, P. VII. VIII.]

[*C. Salman'tica* (L.) and *C. leucophæa* (Jord.) are said to have been found in Jersey with *C. paniculata*; but I have not seen specimens.]

Section 4. *CARDUINEÆ*. Heads many-flowered; flowers all tubular. Involucre in many rows of distinct spinous phyllaries. *Filaments distinct. Pappus in many rows, not surrounded by a prominent margin.*

### 30. ONOPOR'DUM Linn. Cotton Thistle.

1. *O. Acan'thium* (L.); st. erect many-headed, l. elliptic-oblong woolly on both sides sinuate spinous decurrent, outer phyll. lanceolate-subulate recurved and spreading.—*E. B.* 977.—St. 4—5 feet high, woolly, with broad spinous wings, branched. Inv. nearly globose, large, somewhat cottony; phyll. fringed with minute spinous teeth. Fl. purple.—Waste ground in South-east. B. VIII. E.

31. *CARDUUS* Linn. Thistle.

[Many hybrids apparently occur in this genus.]

\* *Pappus rough*. *CARDUUS* Sm., DC., Koch.

1. *C. nutans* (L.); l. decurrent spinous lanceolate sinuate, heads solitary drooping hemispherical, phyll. lanceolate cottony outer ones reflexed.—*E. B.* 1112.—St. 2 feet high, erect, angular, furrowed, cottony, winged. L. hairy on both sides, with woolly veins beneath, pinnatifid with 3-lobed wavy spinous-ciliate segments ending in strong spines. Heads large; fl. crimson; unopened anth. purple. Inv. hemispherical, internal phyll. contracted above the base and then lanceolate.—Waste ground. B. V.—VIII. *Musk-Thistle*. E. S. I.

2. *C. crispus* (L.); l. decurrent spinous-ciliate lanceolate glabrous or cottony beneath deeply pinnatifid, lobes trifid and dentate, heads roundish, phyll. linear-subulate erect or ascending.—*E. B.* 973.—St. about 3 feet high. Phyll. not contracted above their base.— $\alpha$ . *C. crispus* (L.); l. lanceolate usually cottony beneath, heads small clustered subglobular (or ovoid *C. polyanthemus* Koch), phyll. ending in a weak spine, ped. winged to the top, central tubercle of fr. not angular.— $\beta$ . *C. acanthoides* (L.); l. broadly lanceolate less downy beneath, head twice as large solitary or rarely 2 or 3 together subglobular, phyll. strongly spinous erect, ped. often naked at the top, central tubercle of fr. 5-angled.—Dry banks and waste places.  $\beta$  is the less common plant. B. ? VI.—VIII. E. S. I.

3. *C. tenuiflorus* (Curt.); l. decurrent sinuate spinous broadly lanceolate cottony beneath, segments ovate lobed, heads many crowded sessile subcylindrical, phyll. ovate-lanceolate attenuate.—*E. B.* 412. *C. pycnocephalus* Hook.—St. about 3 feet high, slightly branched, with broad deeply lobed spinous wings. L. deeply sinuate or pinnatifid. Involucres nearly glabrous. Fl. pink.—Sandy places near the sea. B. ? VI.—VIII. E. S. I.

\*\* *Pappus feathery*. *CNICUS* Linn., Sm. *CIRCUM* Koch, DC.

† Leaves spinous-hairy above, flowers purple.

4. *C. lanceolatus* (L.); l. decurrent white and cottony beneath pinnatifid, lobes bifid with lanceolate entire segments each terminated by a strong spine, involucres ovate shaggy, phyll. lanceolate spinous spreading.—*E. B.* 107.—St. 3—4 feet high, erect, furrowed, hairy, with strong spinous wings. Heads terminal, solitary or 2 or 3 together, large.—Waste ground. B. VII. VIII. *Spear Thistle*. E. S. I.

5. *C. erioph'orus* (L.); *l.* half-clasping not decurrent white and cottony beneath deeply pinnatifid, lobes bifid the segments lanceolate entire alternately pointing upwards and downwards and each terminated by a strong spine, *involucres* globose shaggy, *phyll.* lanceolate with a long spinous-tipped reflexed point.—*E. B.* 386.—St. 3—4 feet high, much branched, furrowed, hairy. Root-*l.* 1—2 feet long, linear with long divergent lobes which form double rows in a very regular manner. Stem-*l.* similar but smaller. Heads very large; *inv.* covered with a dense white web. A remarkably conspicuous plant.—Waste ground on a limestone soil. B. VIII. *Woolly-headed Thistle.* E.

†† Leaves not spinous-hairy above.

a. *Limb of the cor.* 5-parted to its base. BREEA Less.

6. *C. arven'sis* (Curt.); heads subdicocious, *l. subsessile oblong-lanceolate* pinnatifid spinous wavy, *inv.* ovoid subglabrous, *phyll.* broadly lanceolate adpressed terminating in a short spreading spine, rhizome creeping.—*E. B.* 975.—St. erect, 3—4 feet high, leafy, angular, corymbose above. *L.* very spinous, sessile or very slightly decurrent, varying greatly in width.—†*β. setosus*; *l.* lanceolate flat entire or slightly lobed. *Cir. setosum* M. B.—Fields and roads. *β.* Culross and Kirkwall, S. P. VII. *β.* IX. *Creeping Thistle.* E. S. I.

b. *Limb of the cor.* 5-parted to its middle.

7. *C. palus'tris* (L.); *l. decurrent* lanceolate deeply pinnatifid spinous, *inv.* ovoid crowded, *phyll.* ovate-lanceolate adpressed mucronate.—*E. B.* 974.—St. solitary, erect, 3—5 feet high, wand-like, with wavy spinous wings throughout, slightly branched. *Heads in a terminal cluster, small.* Fl. purple or white. Underside of the *l.* usually cottony. *Inv.* with a slight web.—Wet meadows. A. VII. VIII. E. S. I.

8. *C. pratens'is* (Huds.); *l. mostly radical lanceolate wavy or lobed* pilose above cottony beneath fringed with minute prickles, *stem-l.* not decurrent few clasping, *inv.* globose solitary terminal slightly cottony, *phyll.* lanceolate-attenuate adpressed mucronate, rhizome creeping.—*E. B.* 177. *Cir. anglicum* Lam.—St. 1—2 feet high, cottony, usually quite simple and single-headed, leafless in the upper half with a few scaly bracts, springing singly from the rhizome. *L.* broad, soft, sinuate-dentate, rarely with small 2—3-fid lobes, not pinnatifid, fringed with small but unequal prickles, lower *l.* stalked. Occasionally there are 2 or 3 fl. on a stem.—*C. Forsteri* (Sm.) is probably a hybrid between this and *C. palustris*. It has *l.* slightly decurrent lanceolate all pinnatifid spinous cottony beneath, *st.* paniced, *inv.* ovoid slightly cottony, root cæspitose producing several stems. *Sy. E. B.* 695.—Boggy meadows. P. VI. VIII. E. I.

9. *C. tuberosus* (L.); l. lanceolate deeply pinnatifid pilose above hairy or slightly cottony beneath fringed with minute prickles, stem-l. sessile not decurrent, lobes 2—3-fid, inv. ovoid terminal 1—3 together slightly cottony, phyll. lanceolate mucronate adpressed, root of elliptic tapering fleshy fibres.—*E. B.* 2562. *Cir. bulbosum* DC., Koch.—Not stoloniferous. St. 2 feet high, erect, round, hairy, leafless above the middle with a few minute bracts. Lower l. stalked, stem-l. nearly or quite sessile. May be a hybrid between Sp. 2 and 10.—*C. Woodwardii* (Wats.) much resembles this, and may be a hybrid between *C. acaulis* and *C. pratensis*. *Sy. E. B.* 696.—Greatridge Wood near Boyton, and at Avebury, Wilts. P. VIII. IX. E.

10. *C. acaulis* (L.); l. glabrous radical lanceolate pinnatifid, lobes subtrifid spinous, inv. ovoid glabrous nearly sessile mostly solitary, outer phyll. ovate inner ones gradually longer adpressed, root with filiform fibres.—*E. B.* 161. St. 24. 16.—St. usually almost wanting; sometimes 3—12 in. long, leafy, woolly. L. all stalked, glabrous except a few hairs on the ribs beneath. Heads very large, fl. crimson.— $\beta$ . *C. dubius* (Willd.); st. much branched woolly a foot or more in height. *Willd. Fl. Berol.* f. 11. Perhaps a hybrid between this and *C. arvensis*.—Dry calcareous pastures.  $\beta$ . Saffron Walden, Essex. *Mr. G. S. Gibson.* P. VII.—IX. *Ground-Thistle.* E.

11. *C. heterophyllus* (L.); l. clasping not decurrent glabrous above white and downy beneath lanceolate serrate fringed with minute prickles, root-l. with long stalks clasping at the base, heads ovoid truncate below slightly downy, phyll. ovate or lanceolate acuminate adpressed.—*E. B.* 675.—Creeping. St. 3—4 feet high, furrowed, cottony, slightly branched above. Heads large and handsome. L. very large, undivided or lacinate.—*C. Carolorum* (Jenn. in *Edin. Bot. Tr.* ix. 257) seems to be a hybrid of *C. heterophyllus* and *C. palustris*.—Moist mountain-pastures. P. VII. VIII. E. S.

Section 5. *SILYBEÆ*. *Filaments monadelphous. Pappus in many rows.*

### 32. *SILYBUM* Gaert. Milk-Thistle.

†1. *S. maritimum* (Gaert.).—*Carduus* Sm. *E. B.* 976.—St. 3—4 feet high, ribbed and furrowed. L. very large, oblong-lanceolate, wavy, clasping; radical l. pinnatifid, usually variegated with green and milk-white. Heads large, globose. Phyll. closely adpressed below, leaflike, with a long terminal recurved spine. Fl. purple; tube very long.—Waste places. B. VI. VII. E. S. I.



Suborder II. *LIGULIFLORÆ*. Fl. all perfect, ligulate.—Style cylindrical above; branches long, blunt, equally pubescent. Stigmatic lines prominent, narrow, terminating below the middle of the branches.—Juice milky.

A. *Receptacle naked. Pappus 0.*

33. *LAP'SANA* Linn. Nipplewort.

1. *L. comminis* (L.); l. dentate or lobed stalked, lower l. lyrate, involucre glabrous angular, st. paniced.—*E. B.* 844.—St. and l. hispid or nearly glabrous. St. 1—3 feet high, branched above. Heads small, with yellow fl. in terminal panicles with small subulate bracts at the subdivisions. Inv. of fr. erect.—Waste and cultivated ground. A. VII. VIII. E. S. I.

B. *Receptacle naked. Pappus like a crown, of many entire broad scales.*

34. *ARNOS'ERIS* Gaert. Swine's Succory.

1. *A. pusilla* (Gaert.).—*E. B.* 95.—St. 3—8 in. high, swelling and hollow upwards, leafless, with a minute bract at the base of each branch. Each branch overtopping its predecessor and gradually thickening up to the solitary small terminal head of yellow flowers. Inv. connivent over the fr. when its phyll. become remarkably keeled. Receptacle honeycombed towards the margins. Fr. small, obovate, attenuate below, 5-angled. L. radical, oblong, toothed.—Gravelly and sandy fields, rare. A. VI.—VIII. E. S.

35. *CICHO'RIMUM* Linn. Succory. Chicory.

1. *C. In'tybus* (L.); lower l. rucinate hispid on the keel, upper l. oblong or lanceolate clasping entire, heads axillary in pairs nearly sessile.—*E. B.* 539. St. 6. 15.—St. 2—3 feet high, bristly, alternately branched. Heads many, fl. bright blue, handsome. Floral l. lanceolate from a broad clasping base.—Waste places on a gravelly or chalky soil. P. VII. VIII. E. S. I.

C. *Receptacle scaly. Pappus feathery.*

36. *HYPOCHÆ'RIS* Linn. Cat's-ear.

\* *Pappus with an outer row of shorter bristles.*

1. *H. glabra* (L.); st. branched leafless glabrous, l. oblong, inv. glabrous equalling the flowers.—*E. B.* 575.—St. 3—10 in. high, scaly; primary stem simple, leafless, but lateral branched

stems prostrate leafy. L. spreading in a circle on the ground, glabrous, except a few scattered hairs. Outer row of fruits destitute of a beak; the rest with a long beak.— $\beta$ . *H. Balbisii* (Lois); all the fruits with long beaks.—Sandy and gravelly places.  $\beta$ . In Kent and Salop. A. VII. VIII. E. S.

2. *H. radicata* (L.); st. branched leafless glabrous, l. runcinate blunt, *inv. falling short of the flowers*.—*E. B.* 831.—St. about a foot high, scaly, each branch terminating in a rather large solitary head. L. spreading upon the ground, rough. Stalks slightly thickened beneath the heads. *Fr. all beaked*.—Waste ground. P. ? VII. E. S. I.

\*\* *Pappus in one row.* ACHYROPHORUS Scop.

3. *H. maculata* (L.); st. simple or slightly branched almost leafless, l. obovate-oblong undivided toothed pilose, phyll. bristly on the back.—*E. B.* 225.—St. about a foot high, stout, slightly hairy. L. often all radical. Heads large, fl. deep yellow.—Chalky and limestone hills, rare. P. VII. VIII. E.

D. *Receptacle without scales. Pappus feathery or on the exterior fruits scaly.*

### 37. THRIN' CIA Roth.

1. *T. hirta* (DC.); l. lanceolate sinuate-dentate or entire hispid or hairy with forked or simple hairs, stalks simple pilose below.—*E. B.* 555.—L. all radical, sometimes nearly or quite entire, occasionally runcinate. Stalks often purplish, quite simple, longer than the leaves, somewhat hairy in their lower half. Phyll. downy on the margins at the apex or hairy.—Gravelly places and fields; also in the Fens. P. VI.—IX. E. S. I.

### 38. LEON'TODON Linn. Hawkbit.

\* *Pappus feathery and with an outer row of bristles.*

1. *L. hispidum* (L.); l. radical oblong-lanceolate runcinate hispid with forked hairs, stalks simple naked or with 1 or 2 minute scales thickened upwards hispid.—*E. B.* 554.—L. with regular spreading or reflexed narrow teeth. Stalks green, erect, longer than the leaves. Head drooping in bud, afterwards erect. *Inv. nearly always hairy. Fl. glandular at the end. Fr. muricate*.—Meadows and pastures. P. VI.—IX. E. S. I.

\*\* *Pappus in one row, feathery.* OPORINIA Don, DC.

2. *L. autumnale* (L.); l. radical linear-lanceolate toothed or pinnatifid nearly glabrous, stalk branched scaly and thickened upwards.—*E. B.* 830.—L. all radical tapering at the base, often with long linear spreading segments, usually somewhat hairy particularly on the midrib beneath. Inv. nearly always hairy. Pappus brownish.— $\beta$ . *O. pratensis* (Less.); l. glabrous, stalk mostly simple, inv. shaggy with greenish black hairs. *Apargia Taraxaci* Sm. *E. B.* 1109.— $\gamma$ . *sordida*; l. hairy, st. branched, inv. as in  $\beta$ , plant very large.—Meadows and pastures.  $\beta$ . Mountains.  $\gamma$ . Highland glens. P. VIII. E. S. I.

### 39. TRAGOPOGON Linn. Goat's-beard.

1. *T. minor* (Fries); inv. about twice as long as the flowers, ped. slightly thickened at the very top, l. tapering from a dilated base to a long slender acute point.—*Sy. E. B.* 799.—St. 2 feet high, branched, erect. L. clasping the stem. Phyll. 8, in 2 rows. *Fl.* yellow, truncate, 5-toothed. Anth. dark brown. Marginal fr. angular, striate; angles squamously toothed; interstices tubercled.—Meadows and pastures. B. ? VI. VII.

E. S. I.

2. *T. pratensis* (L.); inv. equalling or shorter than the flowers, ped. slightly thickened at the very top, l. tapering from a dilated base to a long linear acute point keeled.—St.  $1\frac{1}{2}$ —2 feet high, branched, erect. L. clasping the stem. Phyll. 8, in 2 rows. *Fl.* yellow, truncate, 5-toothed. Anth. yellow. There are 2 forms of this plant:—(a) inv. equalling the fl., marginal fr. obscurely striate and rough throughout (*Sy. E. B.* 798); (b) inv. rather shorter than the fl., marginal fr. (in my specimens) yellow slightly furrowed and quite smooth (*E. B.* 434).—Meadows and pastures, less frequent than the preceding. B. ? VI.

E. S. I.

\*3. *T. porrifolius* (L.); inv. longer than the flowers, ped. much thickened upwards, l. tapering slightly dilated just above the base.—*E. B.* 638.—St. 3—4 feet high, erect, branched. L. slightly broader just above the base, then gradually narrowing to an acute point. Heads twice as large as in the two preceding. Inv. usually  $\frac{1}{2}$  longer than the fl., sometimes only equalling or even falling short of them. *Fl.* purple. Marginal fr. with scalelike tubercles throughout but particularly on the ribs.—Moist meadows. B. VI. *Salsify.* E.

### 40. PTERIS Linn.

1. *P. hieracioides* (L.); st. rough with forked and hooked

bristles, l. linear or lanceolate dentate or sinuate, upper l. somewhat clasping, heads solitary terminating the stem and branches, outer phyll. lax oblong bristly on the keel glabrous on the margin, fr. constricted just below the pappus.—*E. B.* 196.—St. 1—3 feet high, divaricately branched above, irregularly corymbose, very rough. Florets yellow.—*P. arvalis* (Jord.) with branches ascending and heads in an umbellate corymb passes gradually into the type.—Dry banks. B. VII.—IX. E.

#### 41. HELMIN'THIA *Juss.* Ox-tongue.

1. *H. echioides* (Gaert.); st. erect hispid with rigid 3-fid and hooked hairs from tubercular bases, phyll. 5 ovate-cordate.—*E. B.* 972.—St. 2—3 feet high, branched, covered, as well as the leaves and involucre, with strong prickles springing from white tubercles and with 3 minute hooks at the apex (glochidate). L. clasping.—Dry banks. A. VII.—IX. E. I.

*E. Receptacle generally without scales. Pappus filiform, deciduous, never feathery, nor dilated at the base. Fruit compressed.*

#### 42. LACTU'CA *Linn.* Lettuce.

\* *Beak long, white. Leaves with a bristly keel.*

1. *L. salig'na* (L.); upper l. linear entire acuminate with a sagittate base, lower l. pinnatifid, *beak twice as long as the fruit.*—*E. B.* 707.—St. 2 feet high, slender, wavy, slightly branched. Heads in small alternate tufts forming long clusters. Flowers yellow.—Chalky places and near the sea. B. VII. VIII. E.

2. *L. virósa* (L.); upper l. horizontal oblong auricled and clasping mucronate-dentate or sinuate, *beak equalling the black fruit.*—*E. B.* 1957.—St. scabrous, 2—4 feet high, leafy, branched above, paniced. Heads scattered, with many heartshaped acute bracts. Plant full of acrid milky juice.—Dry banks. B. VII. VIII. *Acrid Lettuce.* E. S.

3. *L. Scariola* (L.); upper l. upright arrowshaped at the base and clasping sinuate, *beak equalling the pale fruit.*—*E. B.* 268.—St. slightly scabrous below, 2—5 feet high, leafy, paniced. Heads scattered, with many heartshaped acute bracts. Juice rather less acrid than in Sp. 2.—Waste places, rare. B. VII. VIII. *Prickly Lettuce.* E.

\*\* *Beak short. Leaves with a smooth keel.*

4. *L. murális* (DC.); florets 5, l. lyrate-runcinate angled and

toothed clasping, terminal lobe largest, *beak much shorter than the fruit*, heads paniced.—*Prenanthes* L. *E. B.* 457.—St. erect, a foot high, smooth, round, hollow. Flowers bright yellow. Fruit black.—Banks and old walls. P. VII. E. I.

#### 43. TARAX'ACUM *Juss.* Dandelion.

1. *T. officinale* (Wigg.); l. runcinate toothed, fr. linear-obovate blunt and muricate at the top longitudinally striate with a long beak.—Stalks single-headed, radical, hollow. Fl. yellow. L. all radical, very variable, glabrous or slightly hispid.—I. Outer phyll. reflexed or patent.—*a. Leontodon Taraxacum* (L.); *outer phyll. linear deflexed inner ones simple at tip*, fr. yellow its upper half muricate, crown of the root glabrous or woolly, l. runcinate broad.—*E. B.* 510.—*β. T. lævigatum* (DC.); *outer phyll. erect-patent ovate, inner gibbous or appendaged at tip*, fr. reddish yellow muricate at the top, beak with a thickened and coloured base, l. runcinate-pinnatifid with unequal teeth.—*γ. T. erythrospermum* (DC.); *outer phyll. lanceolate adpressed or patent, inner gibbous or appendaged at tip*, fr. bright red muricate at the top, beak with a thickened and coloured base, l. runcinate-pinnatifid with unequal teeth and intermediate smaller ones. Lowermost l. sometimes obovate and dentate (or runcinate when it becomes *T. obovatum* DC.). *Sy. E. B.* 803.—II. Outer phyll. adpressed.—*δ. L. palustre* (Sm.); *outer phyll. ovate-acuminate, inner simple at tip*, fr. pale yellow or brown muricate at the top, l. oblong and entire sinuate-dentate or runcinate; or *outer phyll. ovate-lanceolate, or (L. leptocephalum* R.) lanceolate. *E. B.* 553.—Very common, *γ* in dry places, *δ* in bogs or damp places. P. III.—X. E. S. I.

#### 44. SON'CHUS *Linn.* Sowthistle.

1. *S. oleráceus* (L.); l. undivided or pinnatifid toothed clasping, *auricles spreading arrowshaped, fr. transversely rugose and longitudinally ribbed*, st. branched, heads subumbellate, inv. glabrous.—*E. B.* 843.—St. 2—3 feet high. L. flattish, lower stalked. Fl. yellow.—Common. A. VI.—VIII. *Sowthistle.*  
E. S. I.

2. *S. as'per* (Hoffm.); l. undivided or pinnatifid sharply toothed clasping, *auricles rounded, fr. longitudinally ribbed not transversely rugose*, st. branched, heads subumbellate, inv. glabrous.—*E. B. S.* 2765, 2766.—St. 2—3 feet high. L. crisped, lower stalked. Fl. yellow.—Common. A. VI.—VIII. *Sowthistle.*  
E. S. I.

3. *S. arven'sis* (L.); l. lanceolate runcinate sharply toothed

cordate at the base, uppermost l. entire, *st. simple*, heads corymbose, *inv. and ped. glandular-hairy*, fr. with transversely rugose ribs, creeping.—*E. B.* 674.—St. 3—4 feet high, leafy. L. long, acute. Heads large, fl. yellow.—Fields and waste ground, also in fens. P. VIII. IX. *Corn-Sowthistle.* E. S. I.

4. *S. palustris* (L.); l. linear-lanceolate all acutely arrow-shaped denticulate, lower l. long with 2—4 linear-lanceolate lobes, *st. simple*, heads corymbose, *inv. and ped. glandular-hairy*, fr. with finely rugose ribs, *no stoles*.—*E. B.* 935.—St. 4—6 feet high, leafy. Fl. lemon-coloured.—Marshes, very rare. P. VII. VIII. E.

#### 45. MULGE'DIUM Cass.

1. *M. alpinum* (Less.); l. glabrous lyrate arrowshaped at the base, terminal lobe large triangular-hastate acute, *st. simple*, heads racemose, bracts ped. and *inv. glandular-hairy*, fr. oblong not attenuate with many ribs.—*Sonchus cæruleus* Sm. *E. B.* 2425.—St. 3 feet high, glabrous below, leafy. L. gradually smaller upwards, cordate-acute on the barren shoots. Heads small, many. Fl. blue.—Clova Mountains. P. VIII. S.

*F. Receptacle generally without scales. Pappus filiform, never feathery nor dilated at the base. Fruit terete, ribbed.*

#### 46. CRE'PIS Linn. Hawk's-beard.

\* *Fruit with a long subulate beak. BARKHAUSIA Moench.*

1. *C. taraxacifolia* (Thuil.); l. rough runcinate-pinnatifid, buds erect, *inv.* bristly and downy covering half the pappus, outer phyll. lanceolate with a membranous margin, bracts herbaceous, *fr. all equally beaked*.—*E. B. S.* 2929.—St. 1—2 feet<sup>\*</sup> high, hispid, angular, furrowed, purple below, at length branched, corymbose. L. mostly radical lyrate-runcinate with backward teeth or deeply pinnatifid with the terminal lobe large. Stem-l. few, sessile, clasping, deeply pinnatifid and toothed. Fl. yellow, purple beneath. Fr. narrowing very gradually into a setaceous beak of about its own length, ribs rough.—Limestone districts. B. VI. VII. E.

2. *C. fœtida* (L.); l. hairy runcinate-pinnatifid, unopened buds nodding, *inv.* hairy and downy, outer phyll. lanceolate acute downy, *marginal fr. slightly beaked shorter than the inv., central fr. with long beaks equalling inv., pappus protruding*.—*E. B.* 406.—St. 6—12 in. high, hairy, round, branched. L.

mostly radical; stem-l. few, small, lanceolate, deeply toothed at the base, sessile. Heads solitary, terminal, on long simple stalks. Midrib of the phyll. at length much thickened and hardened. Ribs of the fr. rough.—Chalky places, rare. B. VI. VII. E.

[*C. setosa* (Hall.); l. runcinate-dentate or lyrate-runcinate, st.-l. sagittate entire or incise-dentate below, buds erect, *inv.* not quite covering the pappus, margin of the outer lanceolate acute phyll. and the bracts and the back of the inner phyll. and the ped. hispid with rigid simple bristles.—*E. B. S.* 2945.—Plant 1—2 feet high. Stem-l. large, rather strapshaped, clasping.—Fields. Introduced with seed. A. VII. VIII.] E.

\*\* *Fruit narrowed upwards or obscurely beaked. Pappus silky.*

[*C. pulchra* (L.), *E. B.* 2325, was probably an error.]

3. *C. virens* (L.); *outer phyll. adpressed linear, inner ones glabrous within*, l. lanceolate remotely dentate runcinate or pinnatifid, *uppermost l. linear-arrowshaped clasping with flat margins*, st. subcorymbose, *fr. shorter than the pappus* oblong slightly narrowed upwards with smooth ribs.—*C. tectorum* Sm., *E. B.* 1111 (not Linn.).—Very variable. St. 1—3 feet high. Fl. yellow. [*C. tectorum* (L.) has revolute-margined upper l., phyll. downy within, fr. somewhat beaked with scarious ribs; Mr. Brotherston found it as an escape near Kelso.]—Common. A. VI.—IX. E. S. I.

4. *C. bien'nis* (L.); *outer phyll. oblong-linear lax, inner downy within*, l. runcinate-pinnatifid hispid, uppermost l. lanceolate clasping dentate-pinnatifid, st. subcorymbose, fr. oblong slightly narrowed upwards with nearly smooth ribs and about as long as the pappus.—*E. B.* 149.—St. 1—3 feet high, hispid, nearly leafless above, corymbosely branched. Heads large; fl. yellow. *L. radical and extending halfway up the stem.*—Chalky places, rare? B. VI. VII. E. I.

5. *C. hieracioides* (W. & K.); *phyll. lanceolate-attenuate, outer ones very short adpressed, l. entire* nearly glabrous oblong blunt, lower ones narrowed into a footstalk, *upper l. sessile and somewhat clasping*, st. corymbose, ped. and *inv.* glandular-hairy, fr. much striate slightly narrowed upwards as long as the pappus which is shorter than the involucre.—*C. succisæfolia* Tausch. *Hieracium molle* Sm. *E. B.* 2210.—St. 2—3 feet high, erect, simple below; l. few. Heads few; fl. yellow.—Woods in the North. P. VII. VIII. E. S.

\*\*\* *Fruit not beaked, cylindrical. Pappus stiff, brittle.*  
ARACIUM Moen.

6. *C. paludosa* (Moench); phyll. lanceolate much attenuate glandular-pilose, outer ones short, *l. ovate-oblong taper-pointed runcinate-dentate* narrowed into a footstalk glabrous, *upper l. ovate-lanceolate cordate* and clasping acute entire or dentate, st. subcorymbose, fr. striate scarcely narrowed upwards.—*Hieracium* L. *E. B.* 1094.—St. 2 feet high, leafy, simple, angular. L. large. Fl. yellow.—Damp woods and shady places. P. VII.—IX. E. S. I.

47. HIERA'CIUM, Linn.<sup>1</sup> Hawkweed.

i. *PILOSELLOIDEA*. Stoloniferous. St. scapelike. Fr. minute, crenulate at top, striate. Hairs of pappus equal, very slender.

1. *H. Pilosella* (L.); stoles slender leafy rooting, *scape 1-headed leafless*, *l. oblong or lanceolate hairy* on both sides whitish and densely floccose beneath, *inv. ovate* below ultimately conical, inner phyll. acute, styles yellow.—*E. B.* 1093.—Stoles many, sometimes flowering. Fl. pale yellow; outer striped with red or purple externally.—β. *H. Peleterianum* (Mer.); stoles short thick, *l. st. and inv. with long silky hairs*, heads large, phyll. all lanceolate. *R. xix.* 1468.—Dry banks. P. V.—VIII. E. S. I.

\*2. *H. pratense* (Fr.); stoles leafy, st. pilose floccose and setose above corymbose at top, *l. narrowly obovate-lanceolate green* on both sides pilose slightly floccose beneath, phyll. linear blunt, styles bright yellow.—*H. collinum* Fr. *J. of B.* vi. t. 86. *Edin. Bot. Tr. x. t. 1.*—Stoles not very long. Corymbs dense. Phyll. dark green hairy setose, with a pale margin and tip. Fl. yellow.—Near Selkirk and Edinburgh. P. VI. VII. S.

[*H. aurantiacum* (L.); stoles often wanting, *st. slightly hairy densely corymbose at top* more hairy and setose above, *l. obovate-lanceolate green* and hairy on both sides *not floccose beneath*, phyll. blunt, styles brown.—*E. B.* 1469.—St. 15—20 in. high. Phyll. dark. Fl. orange.—In the North, scarcely naturalized.] P. VI.—VIII. I.

[*H. præaltum*, *H. glomeratum*, and *H. stoloniferum* have been found as escapes in the South of Scotland.]

<sup>1</sup> A *seta* is a gland-tipped hair.—In this genus I have availed myself largely of the 'British Hieracia' by my friend Mr. James Backhouse.



ii. *PULMONAREA*. Radical rosettes in autumn which produce the persistent root-l. of the next year. Phyll. interruptedly or irregularly imbricate. Fr. short, truncate, not crenulate, striate. Hairs of pappus unequal, rigid.

\* *Alpiniformes*. Inv. shaggy or silky. Outer phyll. lax; inner acuminate or acute. Fl. hairy externally, more or less pilose at the tips. St.-l. 1 or few, or leaflike bracts.—*H. alpinum* L.

3. *H. melanocephalum* (Tausch.); deep green, st. 1-headed hairy floccose, rt.-l. lanceolate or ovate spathulate narrowed into petioles, st.-l. 1 or few sessile, buds slightly nodding, inv. hemispherical shaggy with long soft black-based hairs setose, phyll. few broad acuminate lax, styles yellow.—*H. alpinum* Backh., *Sy. E. B.* 827.—St. 4—8 in. high. Usually 1. st.-leaf. Head large, always solitary; fl. bright yellow, with short hairs at the top.— $\alpha$ ; rt.-l. ovate-spathulate.— $\beta$ . *insigne*; rt.-l. lanceolate with a few large teeth, heads very large.—Lofty mountains of Scotland. P. VII. VIII. S.

4. *H. holosericeum* (Backh.); green, st. 1-headed shaggy silky floccose, rt.-l. lanceolate-spathulate or linear-lanceolate blunt hairy on both sides narrowed into winged petioles, st.-l. few small sessile, buds nodding, inv. turbinate shaggy with long silky white black-based hairs, outer phyll. very lax leaflike blunt, inner phyll. adpressed linear acute, styles yellow.—*H. alpinum* Sm. *E. B.* 1110.—St. 3—9 in. high. Usually 1—2 st.-leaves. Head solitary; outer phyll. very broad, often scattered below the head. L. entire or sometimes slightly denticulate. Fl. densely hairy at the tips, bright yellow.—Lofty mountains. Cumberland. Grampian and Breadalbane Mountains. P. VII. VIII. E. S.

5. *H. eximium* (Backh.); green, st. usually 1-headed hairy floccose, rt.-l. lanceolate acute sharply toothed hairy on both sides narrowed into broadly winged petioles, st.-l. small slender, buds nodding, inv. truncate below shaggy with rather silky black-based hairs, phyll. many linear-attenuate, outer phyll. small lax.—*Sy. E. B.* 825. *H. villosum* Sm., *E. B.* 2379, not L.—St. 6—15 in. high; l. few. Rarely more than 1 large head. L. sometimes entire, usually with large teeth. Flowers twice in cultivation, (1) with one head, (2) with many nearly parallel branches and many heads.— $\alpha$ ; st. long, rt.-l. lanceolate dentate, styles livid.— $\beta$ . *tenellum*; st. short slender, rt.-l. lanceolate or linear-lanceolate, styles yellow. Considered a distinct species by Dr. Grenier.—Lofty mountains. Clova. Braemar. P. VII. VIII. S.

6. *H. calenduliflorum* (Backh.); deep green, st. usually 1-headed hairy floccose, *rt.-l. broadly ovate-spathulate blunt* apiculate sharply toothed hairy on both sides narrowing into broadly winged petioles, *st.-l. very small slender*, buds nodding, *inv. truncate below shaggy with rather silky black-based hairs*, *phyll. many linear-attenuate*, outer *phyll. lax*, *styles livid.*—*Sy. E. B.* 824.—*St.* 6—14 in. high. *St.-l. few.* Head most usually solitary, very large. Youngest l. sometimes rather acute. Original l. nearly circular. Nearly allied to *H. eximium*.—Lofty mountains. Lochnagar. Clova. P. VII. VIII. S.

\*\* *Nigricantes.* *Inv. villose or hairy.* *Phyll. adpressed, or few outermost lax.* *Fl. nearly hairless externally, minutely pilose or subglabrous at the tips.*—*H. nigrescens* Hook.

† *St. short with 1 or few heads.*

7. *H. gracilentum* (Backh.); green, st. usually 1-headed hairy floccose, *rt.-l. lanceolate or oblong-lanceolate* narrowed into petioles, *st.-l. few large narrow-lanceolate attenuate below*, *inv. ventricose rounded below shaggy with black soft hairs setose*, *phyll. few broad acuminate adpressed with floccose tips* outermost lax, *styles livid.*—*Sy. E. B.* 828. *H. alpinum* var. *melanocephalum* Fries, not Bab., nor *H. melanocephalum* Tausch.—*St.* 6—10 in. high, hairy and floccose, with straight simple diverging branches and many heads when under cultivation; *st.-l. usually 2 or more.* L. broad, narrowed gradually below; original l. roundish. Petioles sometimes slightly winged. Heads usually solitary, rather large; fl. bright yellow, nearly or quite glabrous on the back.—Lofty mountains. P. VII. VIII. S.

8. *H. globosum* (Backh.); *glaucous or green*, st. few-headed floccose, *rt.-l. obovate or ovate-lanceolate subacute glabrous above* entire or dentate narrowed into petioles, *inv. rounded below ultimately globose* greenish-black with short black-based hairs, *phyll. many attenuate acute adpressed*, *styles yellow.*—*Sy. E. B.* 829.—*St.* 6—12 in. high, usually 1-headed, rarely branching even from near the base, leafless or with few narrow bractlike leaves. Original l. blunt. Innermost l. sometimes acute. Petioles short sometimes slightly winged. *Buds globose.* Fl. bright yellow.—Cairngorm Mountains. P. VII. VIII. S.

†† *St. frequently lengthened and branched with several heads.*

9. *H. nigrescens* (Willd.); green, *rt.-l. ovate or lanceolate coarsely and irregularly dentate* in their lower half hairy with slender petioles, st. simple or branched with few heads and few

lanceolate or oblong shortly stalked l., *inv. ovate below*, dark green with soft black-based hairs setose, ped. densely floccose, phyll. acuminate, outer ones blunt, *styles brownish*.—*Sy. E. B.* 832.—Variable in height. Fl. deep brilliant yellow, pilose at the tips. Phyll. overtopping the buds. L. rarely nearly glabrous above. Rt.-l. suddenly narrowed below; original l. roundish. St.-l. usually 1, narrowed at both ends. St. floccose, branching much in cultivation.—Highland mountains. P. VII. VIII. S.

10. *H. lingulatum* (Backh.); green, *rt.-l.* lanceolate or oblong apiculate *denticulate* or dentate hairy with short petioles, st. simple or branched with few heads and few lanceolate or lanceolate-attenuate acute sessile l., *inv. broad becoming truncate below very dark* with soft black-based hairs slightly floccose, phyll. attenuate acute incumbent, *style livid*.—*Sy. E. B.* 834. *H. saxifragum* Bab.—St. 15—24 in. high. Fl. bright yellow. Phyll. greenish black, overtopping the buds. L. *coarsely hairy above*, entire towards the end. Original l. broad and rounded. St. with scattered black-based hairs throughout, a little floccose. St. and *inv.* nearly without setæ.—Mountain Glens. Clova. Braemar. Ben Lawers. P. VII. VIII. S.

11. *H. senes'cens* (Backh.); green, *rt.-l.* elliptic-lanceolate denticulate or remotely dentate hairy with slender petioles, st. simple with few heads and 1 or 2 *linear-lanceolate l. with slender stalks*, *inv. ovate or turbinate below* with short black-based hairs and many setæ slightly floccose, phyll. acute adpressed (?), styles yellow.—*Sy. E. B.* 833.—St. 15—18 in. high. Fl. golden. Phyll. of the buds with a tuft of white down at the tips. L. with scattered hairs on both sides, entire at both ends. Original l. broader. Lowest ped. above the middle of the stem.—Grassy slopes and edges of streams in mountain districts. Clova. Braemar. Ben Voirlich, Dumbartonshire. P. VII. VIII. S.

12. *H. chrysan'thum* (Backh.); green, *rt.-l.* ovate acute at both ends *sharply irregularly and deeply toothed* hairy with long slender petioles, st. simple or branched with few heads and linear-lanceolate or subulate stalked st.-l., *inv. rounded below* floccose with short black-based hairs and setæ, *phyll.* many *linear-attenuate* outermost small and lax, styles dull yellow.—*Sy. E. B.* 830.—St. 9—15 in. high. Fl. golden. Phyll. greenish black, overtopping the drooping buds, incumbent. L. with scattered hairs on both sides, entire towards the end. Original l. blunt, roundish. St. often with only 1 head, with subplumose hairs below, floccose setose and with black-based hairs above, sometimes branching throughout. Hairs on st. and *inv.* with

dull-reddish tips. Buds sometimes pilose at the tip.—*β. microcephalum*; st. simple with 1 or few nearly erect heads, l. dentate or nearly entire, inv. urceolate, styles rather livid. *H. atratum* Bab.—Lofty mountains in Scotland. *β.* Cumberland and Loch-nagar. P. VII. VIII. E. S.

††† *St. usually lengthened.*—*H. Lawsoni* Hook.

13. *H. anglicum* (Fr.); glaucous green, rt.-l. ovate-lanceolate apiculate or acuminate denticulate with long shaggy winged *petioles*, st. with few l. and few heads, *st.-l. clasping* ovate acuminate, inv. ventricose hairy setose, phyll. acuminate, *styles livid.*—*H. cerinthoides* Backh.—St. 12—18 in. high, branching in cultivation. Original rt.-l. roundish. Rhizome short. Fl. pale yellow.—*α. H. anglicum* (Fr.); st. with long arcuate ascending *lateral ped.*, *st.-l. usually* solitary sessile scarcely clasping, phyll. all adpressed. *Sy. E. B. 836.*—*β. H. decipiens* Syme; st. scarcely branched with 1 or few terminal heads, *st.-l. few clasping*, outer phyll. lax. *Sy. E. B. 837.* *H. cerinthoides* Bab.—*γ. acutifolium* (Backh.); *very glaucous*, st. much branched, l. very acute or acuminate sharply and coarsely toothed nearly glabrous, phyll. adpressed.—*α* and *γ. H. vogesiacum* Gren.—Mountains. *α.* Teesdale. Craven. Lake district. Scottish Highlands. *β.* Grampians. Ben Bulbin, Sligo. *γ.* Cairntoul and by rivers Clunie and Dee in Scotland. Aberdeen Links. P. VII. VIII. E. S. I.

14. *H. iricum* (Fr.); glaucous green, rt.-l. ovate or oblong-lanceolate acute with short shaggy winged *petioles*, st. leafy corymbose at top, *st.-l. clasping broadly ovate* taper-pointed denticulate or dentate, lower ones narrowed below, ped. and inv. hairy setose floccose, inv. truncate below, phyll. blunt, *styles livid.*—*H. Lapeyrousi* Bab. *E. B. S. 2915.*—St. 1—3 ft. high, very leafy, rigid, not branching, 1-headed on barren soil. Ped. rigid, diverging. Original rt.-l. blunt. L. often purplish at the end. *St.-l. decreasing* successively upwards, upper l. broad and rounded below. Phyll. broad, narrowed upwards.—Chiefly in mountain districts. P. VII. VIII. E. S. I.

\*\*\* *Pallidiformes.* Inv. more or less hairy. Phyll. adpressed. Fl. hairless externally, nearly or quite glabrous at the tips.

† *Styles yellow, rarely with minute fuscous hairs.*

15. *H. pal'idum* (Fr.); glaucous, rt.-l. ovate-lanceolate acute apiculate-dentate or dentate below with slender *petioles*, *st. few-headed* with about 1 sessile or stalked l., inv. ovate below ventri-

cose constricted above floccose slightly hairy and setose, phyll. acute or cuspidate, styles yellow.—*Sy. E. B.* 840.—St. 6—18 in. high. Rhizome long, prostrate. *L. fringed with long hairs* which sometimes spread over the surfaces, obscurely floccose beneath. St.-l. nearly or quite sessile; if more than 1, then the lower is stalked. Original rt.-l. rounded at both ends. Fl. very slightly pilose at the tips, bright yellow.—Mountains. P. VI. VII. and IX. E. S. I.

16. *H. cineras'cens* (Jord.); ash-colour or glaucous, rt.-l. oval or ovate-lanceolate rounded at both ends entire or denticulate, petioles shaggy, st.-l. 0 or 1, st. few-headed with straight-based ped., inv. rounded below floccose setose slightly hairy, *phyll. acuminate or cuspidate*, styles yellow.—*Sy. E. B.* 841. *H. lasiophyllum* Bab. *H. pallidum* var. *crinigerum* Fr. !—St. 8—20 in. high. Very hairy. Branches slender, long. Young heads roundish. Outer phyll. sometimes bluntish. Fl. bright yellow. L. rosulate, persistent, broad, apiculate, usually with many scattered bulbous hairs on each side. Original l. roundish. St.-l. rare, lanceolate, acute at both ends.—Mountains. P. VII. VIII. E. S. I.

17. *H. Gibsóni* (Backh.); glaucous green, rt.-l. ovate blunt apiculate rather truncate or cordate below denticulate, petioles slender, st. *leafless* rigid simple or forked with *straight-based ped.*, *inv. truncate below* floccose hairy setose, *phyll. broad and blunt* incumbent, styles yellow.—*Sy. E. B.* 842.—St. 6—18 in. high. Fl. bright yellow. Phyll. not cuspidate. L. rosulate, persistent, very broad, blotched with purple above.—On limestone rocks near Settle, Yorkshire. Co. Sligo and Clare. P. VII. VIII. E. I.

18. *H. argen'teum* (Fr.); *very glaucous*, rt.-l. lanceolate acute *denticulate towards the middle* or entire glabrous above with short petioles, st. simple or branched with few sessile or stalked l., branches long straight, inv. broadly ventricose becoming subtruncate below slightly setose hairy and floccose, *phyll. blunt* adpressed, styles yellow.—*Sy. E. B.* 843. *H. pallidum*  $\beta$ . *persicifolium* Bab.—St. 1—2 feet high. Buds cylindrical. Inner phyll. subacute, pale-edged. Fl. bright yellow. L. with scattered hairs beneath and at the edges. Petioles usually short. Original l. bluntish. Upper st.-l. small narrow.—Mountain districts of England, Scotland, and Wales. P. VII. VIII. E. S.

19. *H. nit'idum* (Backh.); *dark green*, rt.-l. lanceolate acute *coarsely and sharply toothed* glabrous above with short petioles, st.-l. 1 or 0, st. few-headed, ped. scaly, inv. ventricose setose and

hairy slightly floccose, phyll. acuminate or bluntish, styles yellow.—*Sy. E. B.* 844.—St. 15—24 in. high, nearly hairless. Heads large; fl. bright yellow. L. slightly hairy beneath; teeth very large, not directed downwards. Petioles usually short. Original l. roundish.—By mountain-streams in Aberdeenshire. P. VII. VIII. S.

20. *H. aggregatum* (Backh.); deep green, rt.-l. ovate blunt coarsely dentate below floccose beneath when young, st.-l. l lanceolate narrowed at both ends, *st. corymbose* and floccose at the top, *ped. aggregate*, inv. subtruncate below loosely floccose setose nearly hairless, *phyll. blunt*.—*Sy. E. B.* 845. *H. bifidum* Koch?—St. 12—20 in. high, usually simple. Ped. forming a close corymb as in *H. umbellatum*. Heads many. Buds cylindrical. Fl. bright yellow or orange. Styles yellow. L. rosulate, persistent; original l. nearly round; innermost acutish. Petioles rather winged, slightly hairy. St.-l. sessile. Phyll. adpressed.—By mountain-streams in the Grampian Mountains. P. VII. VIII. S.

†† *Styles more or less livid.*

21. *H. murorum* (L.); green, rt.-l. ovate often cordate and with large patent or descending teeth below, petioles shaggy, st.-l. 0 or 1 stalked and placed high, heads subcorymbose, *ped. arcuate* ascending, inv. thinly clothed floccose setose with few hairs, phyll. acuminate erect overtopping the glabrous-tipped buds, styles livid.—*Sy. E. B.* 846.—St. 12—18 in. high. Young heads cylindrical. L. usually thin. St.-l. ovate-acuminate, often rounded or cordate below. Styles rarely yellowish.—Woods and rocks. P. VI. VII. E. S. I.

22. *H. cæ'sium* (Fr.); cæsious or dull green, rt.-l. ovate or lanceolate rounded or narrowed below irregularly dentate-serrate with patent or ascending teeth, petioles slender, st.-l. 0 or 1 placed low, st. few-headed with *straight-based ped.*, inv. rounded below floccose hairy slightly setose, *phyll. bluntish*, inner ones acute incumbent, styles slightly livid.—*H. murorum* Sm. *E. B.* 2082. *Fl. Dan.* 2598.—St. 12—18 in. high. Young heads roundish. Fl. bright yellow. Phyll. not cuspidate; inner very slender, pointed. Rootstock long. L. nearly glabrous above, coriaceous, not fringed with coarse hairs; innermost acute. St.-l. not stalked, narrowed below.—*H. obtusifolium* (Backh.); yellowish green, original rt.-l. ovate rounded at both ends entire, petioles densely villose, st.-l. stalked ovate-lanceolate, inv. ventricose becoming truncate below, phyll. acuminate, buds pilose at the tip; is probably a variety and is found at Clova.—Mountains. P. VII. VIII. E. S. I.

23. *H. flocculosum* (Backh.); ashy-green, *rt.-l.* ovate or oval blunt or acutish with small teeth below *floccose on both sides* narrowed into long petioles, *st.-l.* few large  $\frac{1}{2}$ -clasping, *st.* subcorymbose *floccose* throughout, ped. straight-based, inv. ovate below floccose setose hairy, phyll. acuminate incumbent, styles rather livid.—*Sy. E. B.* 848. *H. stelligerum* Backh. (not. Froel.).—*St.*  $1\frac{1}{2}$ —2 ft. high. Fl. bright yellow. L. rosulate, persistent; original spatulate; innermost acutish. Petioles rather slender and woolly. *St.-l.* stalked; uppermost sessile.—Alpine streams on Grampian Mountains. P. VII. VIII. S.

24. *H. vulgatum* (Fr.); green or glaucous, l. oblong or lanceolate often with patent or *forward teeth* on lower half, *rt.-l.* narrowed into petiole, *st.-l.* often many all or uppermost sessile, heads panicled or subcorymbose, inv. and *straight ascending ped.* floccose setose with few hairs, *phyll. equally attenuate* acutish incumbent, styles livid.—*H. sylvaticum* Sm. *E. B.* 2031.—Very variable, 1—3 feet high. The more common plant has panicled heads cylindrical when young, glaucescent l. purplish beneath, persistent *rt.-l.*, few *st.-l.*, greenish inv.; but sometimes the *st.* is very leafy, heads panicled, *rt.-l.* evanescent.—*H. maculatum* (Sm.) has few broad purple-based *st.-l.*, and imperfectly corymbose heads. The spots on the leaves are not constant: in shade they are absent.—*Sy. E. B.* 849. Many other forms are named, but they defy definition.—Woods, banks, walls. P. VII.—IX. E. S. I.

\*\*\*\* *Pseudo-accipitrinae.* Inv. subglabrous. *Rt.-l.* often forming a winter rosette but rarely present with the flowers.

25. *H. gothicum* (Fr.); l. ovate-lanceolate denticulate or dentate at the middle, *rt.-l.* shortly stalked usually evanescent, lower *st.-l.* often stalked and aggregate, upper *st.-l.* sessile, heads few subcorymbose, ped. rigid erect slightly floccose, inv. roundly ventricose subglabrous or slightly hairy, *phyll. dark green broad blunt* incumbent, style yellow with fuscous hairs.—*Sy. E. B.* 851.—Very variable, 1—4 ft. high, rigid, rarely much branched. *Rt.-l.* very rarely persistent. L. all similar, firm, narrowed at both ends, sometimes very narrow. Heads large, cylindrical when young. Inv. with few brownish hairs and very slightly floccose or subglabrous.—*H. latifolium* (Backh.); *st.* branched, l. broadly lanceolate, style nearly wholly yellow.—Heathy or grassy subalpine places. P. VII.—IX. E. S. I.

26. *H. Dewári* (Sy.); l. oblong-lanceolate *denticulate*, lowest oval with winged stalks, *upper l. ovate-lanceolate*  $\frac{1}{2}$ -clasping, ped. slightly floccose and setose scarcely hairy, heads in lax panicle, invol. subcylindrical obconic slightly hairy and setose,

phyll. dark green few the outer short adpressed inner with pale edges, *style fuscous*.—*Edin. Bot. Trans.* xiii. 211, t. 5.—Bright green. St. 1—3 ft. high. Root-l. rare and persistent.—Mountain glens. P. VIII. IX. S.

iii. *ACCIPITRINA*. Forming closed buds at the base of the st. in autumn. No true rt.-leaves with the flowers. St. leafy. Phyll. in many rows. Hairs of pappus rigid, unequal.

\* *Pseudo-pulmonareæ*. Rarely forming small basal rosettes which do not outlive the winter.

27. *H. tridentatum* (Fr.); st. leafy, l. lanceolate few-toothed at the middle falsely 3-veined, rt.-l. narrowed into winged petioles, st.-l. sessile, bracts nearly linear, heads corymbose, round-based inv. and long floccose *slender ped.* with short hairs, *inv. of fr. truncate below constricted in the middle*, phyll. pale-edged subglabrous rather blunt, styles livid.—*Sy. E. B.* 852.—Rt.-l. rarely persistent. St. 2—5 ft. high, often much branched above, with leaves decreasing successively to small bracts, hairy below. Petioles and base of sessile l. villose. Teeth of l. usually large. Heads rather small. *Inv. of fr.* often an acute cone, sometimes floccose. Pappus reddish.—Banks. P. VIII. IX. E. S.

\*\* *Aphyllopodæ*. Never forming rosettes. Root-leaves fading early.

† *Flowers minutely pilose.*

28. *H. prenanthoides* (Vill.); st. leafy, l. all clasping net-veined and glaucous beneath *hairy on both sides*, lowest narrowed into winged auricled petioles, interm. pinched above their base, uppermost l. cordate-lanceolate, heads in a corymbose panicle, *ped.* short lax very floccose and *setose*, *inv.* cylindrical rather floccose *very setose*, *outer phyll. few much the shortest*, *inner all about equal* and blunt, styles with dark hairs, fr. pale.—*E. B.* 2235.—St. 1—3 ft. high rather rigid, usually hairy. Pan. leafy below. Heads small. Pappus whitish.—River-sides in the North. P. VII. VIII. E. S. I.

[*H. Juranum* Fr.; like Sp. 28 but l. fewer smaller upwards nearly oval, lowest l. abruptly contracted below, interm. l. not pinched above the base. *H. Borreri*, *Sy. E. B.* 859. *H. denticulatum* Borr. MS. not Sm.—Harehead wood near Selkirk? P. VII. VIII.]

29. *H. stric'tum* (Fr.); st. leafy, l. *sessile* oblong-lanceolate denticulate glaucous beneath, lowest lanceolate much narrowed



below *uppermost l.* rounded below *scarcely clasping*, heads irregularly corymbose, *ped.* straight floccose *scarcely setose*, inv. elliptic *thinly floccose setose* and hairy, *phyll.* *irregularly imbricate* blunt, outermost rather acute, styles with dark hairs, fr. fuscous.—*H. denticulatum* Sm. *E. B.* 2122.—St. 1—3 ft. high, slightly hairy. L. broad, not auricled, mostly with bulbous hairs beneath. Heads thicker and paler than those of *H. prenanthoides*. Phyll. pale-edged, not in two distinct ranks. Pappus reddish.—Mountain glens. P. VII. VIII. E. S. I.

†† *Flowers glabrous at the tips.*

30. *H. umbellatum* (L.); st. leafy, *l.* all linear or oblong-lanceolate narrowed below sessile net-veined, heads in an umbellate corymb, top of *ped.* floccose, turbinate-based *inv. uniformly dark green glabrous*, *phyll.* blunt with recurved points, styles yellow.—*E. B.* 1771.—St. 1—4 ft. high, rigid, hairy below. L. usually all alike, or upper rather broad and rounded at the base. Outer *phyll.* often very small, slender, acute. Inv. very rarely straw-coloured, rarely with a few white hairs. Pappus whitish.—Sandy and stony places. P. VII.—IX. E. S. I.

31. *H. crocátum* (Fr.); st. leafy, *l.* oblong or narrowly lanceolate sessile with a broad base falsely 3-veined, lower *l.* narrowed downwards, heads subcorymbose, *branches nearly simple*, *ped.* floccose, inv. very broad and blunt below subglabrous, *phyll.* blackish blunt adpressed, styles yellowish with fuscous hairs.—*Sy. E. B.* 856. *H. inuloides* Tausch.—Very variable. St.  $1\frac{1}{2}$ —4 ft. high, rigid. L. toothed or nearly entire; lower often narrowed very gradually but again slightly enlarged at their base. Inv. rarely pilose. Inner *phyll.* sometimes pale-edged.—Mountain districts. P. VII. VIII. E. S. I.

32. *H. corymbosum* (Fr.); st. very leafy, *l.* ovate or lanceolate dentate about their middle, lowest *l.* narrowed into petioles, heads in a spreading leafy branched corymbose panicle, inv. ovate becoming conical and subtruncate below not constricted with scattered hairs and setæ, *phyll.* blunt uniformly greenish black adpressed, styles yellowish.—*Sy. E. B.* 855. *H. rigidum* and *H. corymbosum* Backh., Bab.—St. 3—4 ft. high, rigid, nearly glabrous, or slightly floccose. L. sometimes floccose beneath, more or less hairy on both sides; uppermost narrowed from a broad base. Pappus tawny.—Mountain districts. P. VII.—IX. E. S. I.

33. *H. boreale* (Fr.); st. leafy, *l.* ovate or lanceolate dentate below falsely 3-veined, upper broad sessile scarcely clasping, lowest *l.* much narrowed below but scarcely stalked, heads in a rather leafy corymb or panicle, top of *ped.* floccose, ovate-based

*inv. uniformly blackish green nearly glabrous or pilose, phyll. blunt adpressed, styles livid blackish.*—*H. sabaudum* Sm. *E. B.* 349.—St. 2—4 ft. high, rigid, usually hairy and often very leafy below. Base of corymb or panicle leafy; branches nearly erect. Interm. l. narrowed to a rounded base; upper with a broad rounded or subcordate base. Phyll. turning black, scarcely at all setose or floccose, rarely with spreading tips. Pappus whitish—Banks. P. VIII. IX. E. S. I.

*Index to the Hieracia.*

- |                                |                            |                            |
|--------------------------------|----------------------------|----------------------------|
| <i>H. acutifolium</i> , 13.    | <i>Gibsoni</i> , 17.       | <i>pallidum</i> , 15.      |
| <i>aggregatum</i> , 20.        | <i>globosum</i> , 8.       | <i>pallidum</i> , 16, 18.  |
| <i>alpinum</i> , 3—7.          | <i>glomeratum</i> , 2*.    | <i>Peleterianum</i> , 1.   |
| <i>anglicum</i> , 13.          | <i>gothicum</i> , 25.      | <i>persicifolium</i> , 18. |
| <i>argenteum</i> , 18.         | <i>gracilentum</i> , 7.    | <i>Pilosella</i> , 1.      |
| <i>atratum</i> , 12.           | <i>holosericeum</i> , 4.   | <i>præaltum</i> , 2*.      |
| <i>aurantiacum</i> , 2*.       | <i>insigne</i> , 3.        | <i>pratense</i> , 2.       |
| <i>bifidum</i> , 20.           | <i>inuloides</i> , 31.     | <i>prænanthoides</i> , 28. |
| <i>boreale</i> , 33.           | <i>iricum</i> , 14.        | <i>rigidum</i> , 32.       |
| <i>Borreri</i> , 28*.          | <i>juratum</i> , 28*.      | <i>rotundatum</i> , 21.    |
| <i>cæsius</i> , 22.            | <i>Lapeyrousii</i> , 14.   | <i>sabaudum</i> , 33.      |
| <i>calenduliflorum</i> , 6.    | <i>lasiophyllum</i> , 16.  | <i>saxifragum</i> , 10.    |
| <i>cerinthoides</i> , 13.      | <i>latifolium</i> , 25.    | <i>senescens</i> , 11.     |
| <i>chrysanthum</i> , 12.       | <i>lingulatum</i> , 10.    | <i>stelligerum</i> , 23.   |
| <i>cinerascens</i> , 16.       | <i>maculatum</i> , 24.     | <i>stoloniferum</i> , 2*.  |
| <i>collinum</i> , 2.           | <i>melanocephalum</i> , 3. | <i>strictum</i> , 29.      |
| <i>corymbosum</i> , 32.        | <i>melanocephalum</i> , 7. | <i>sylvaticum</i> , 24.    |
| <i>crocatum</i> , 31.          | <i>microcephalum</i> , 12. | <i>tenellum</i> , 5.       |
| <i>decipiens</i> , 13.         | <i>murorum</i> , 21.       | <i>tridentatum</i> , 27.   |
| <i>denticulatum</i> , 28*, 29. | <i>murorum</i> , 22.       | <i>umbellatum</i> , 30.    |
| Dewari, 26.                    | <i>nigrescens</i> , 9.     | <i>villosum</i> , 5.       |
| <i>dubium</i> , 2.             | <i>nitidum</i> , 19.       | <i>vogesiacum</i> , 13.    |
| <i>eximium</i> , 5.            | <i>obtusifolium</i> , 22.  | <i>vulgatum</i> , 24.      |
| <i>flocculosum</i> , 23.       |                            |                            |

[*H. Auricula*, *H. dorrense*, *H. Oreades*, and *H. villosum* are omitted, having probably never been found in Britain. *H. amplexicaule* is not even naturalized. *H. pulmonarium* is perhaps not determinable, but may be *H. nigrescens*.]

*Anomalous Genus.* Order AMBROSIACEÆ Link.

48. XAN'THIUM Linn.

[*X. strumarium* (L.); st. without spines, lower l. heartshaped 3-lobed at the base coarsely dentate, beaks of the fr. 2 straight.—*E. B.* 2544.—Involucre of the fr. oval, downy.—Rich waste land, not naturalized. A. VIII. IX.] E.

[*X. spinosum* (L.); has also been found.]

## Order XLVI. CAMPANULACEÆ.

Cal. superior, 5-fid or entire. Cor. gamopetalous, inserted on the calyx, 5-lobed, regular or irregular. Stam. inserted with, but not adhering to the cor., alternate with its lobes; anth. distinct or cohering, 2-celled, opening longitudinally. Fr. dry, capsular, opening by lateral fissures or valves at the top, many-seeded. Embryo straight, in the axis of fleshy albumen.—No stipules.

Tribe I. *LOBELIÆ*. Cor. irregular. Anth. cohering. Style glabrous with a fringe of hairs below the stigma.

1. *LOBELIA*. Cal. 5-fid. Cor. irregular; tube split to the base at the back; limb 2-lipped, 5-parted. Anth. 5, cohering. Stigma blunt, surrounded by a cupshaped fringe. Caps. 2—3-celled, opening at the end by 2—3 valves.

Tr. II. *CAMPANULÆ*. Cor. regular. Anth. usually free. Style pubescent.

2. *JASIONE*. Cal. 5-fid. Cor. rotate with 5 long linear segments. Anth. cohering at their base. Style hairy, bifid. Caps. 2-celled, opening broadly at the end by short teeth.
3. *PHYTEUMA*. Cal. 5-parted. Cor. 2-lipped, with 5 long linear segments. Anth. free; filaments dilated at the base. Style hairy, 2—3-fid. Caps. 2—3-celled, bursting at the sides.
4. *CAMPANULA*. Cal. 5-parted. Cor. mostly bellshaped, with 5 broad and shallow segments. Anth. free; filaments dilated at the base. Stigma 3—5-fid. Caps. not long, 3—5-celled, opening by lateral pores below the cal.-limb.
5. *SPECULARIA*. Cor. rotate. Caps. linear-oblong, prismatic, opening by lateral pores between the segments of the calyx. Otherwise like *Campanula*.
6. *WAHLENBERGIA*. Caps. half superior, 3-celled, opening by 3—5 valves above the segments of the calyx. Otherwise like *Campanula*.

Tribe I. *Lobeliæ*.1. *LOBE'LIA* Linn.

1. *L. Dortman'na* (L.); l. radical linear entire of 2 parallel tubes, st. simple nearly leafless.—*E. B.* 140.—Rootstock fleshy,

with filiform runners. L. blunt, 1—2 in. long, submerged. St. 12—18 in. high, with or without 2—3 small bractlike leaves. Fl. pale lilac, distant, in a raceme, slightly raised above the water.—In lakes with a gravelly bottom. P. VII.

E. S. I.

2. *L. úrens* (L.); st. nearly upright leafy, lower l. obovate or oblong slightly toothed, upper lanceolate serrate, fl. in long terminal racemes.—*E. B.* 953.—St. 1—2 ft. high, branched, angular, roughish. Racemes erect, simple, lax. Fl. light blue.—Heath near Axminster, Devon. P. VIII. IX. E.

## Tribe II. *Campanuleæ*.

### 2. JASIO'NE *Linn.* Sheep's Scabious.

1. *J. montána* (L.); root simple, l. oblong bluntish wavy, fl. in long stalked heads.—*E. B.* 882.—St. several from the crown of the root, 6—12 in. long, simple or branched, pilose, leafy below, bare and usually glabrous above. Radical l. in a rosette. Fl. small, in terminal heads with involucre. Bracts glabrous or hairy. Cal.-segm. subulate, glabrous. Cor. light blue.—Dry places. B. VII. E. S. I.

### 3. PHYTEU'MA *Linn.*

1. *P. obiculáre* (L.); heads of fl. globose, of fr. oblong, l. crenate-serrate, lowermost cordate-ovate stalked, upper l. linear-lanceolate sessile, outer bracts ovate-lanceolate attenuate, stigmas 3.—*E. B.* 142.—St. 4—18 in. high, each with 1 terminal head of blue flowers.—Chalky downs. P. VII. E.

2. *P. spicátum* (L.); heads of fl. oblong, of fr. elongate cylindrical, lower l. cordate-ovate somewhat doubly serrate stalked, upper l. linear-lanceolate sessile, bracts linear, stigmas 2.—*E. B. S.* 2598.—St. 1—2 feet high, each with a solitary terminal head of cream-coloured flowers. Spike of fruit often 2—3 in. long.—Woods and thickets about Waldron, Sussex. P. VII. E.

### 4. CAMPAN'ULA *Linn.* Bell flower.

\* *Caps. sessile, erect; pores at the base.*

1. *C. glomeráta* (L.); l. minutely crenate-serrate, lowermost stalked ovate-lanceolate generally cordate at the base, upper l. half-clasping sessile ovate acute, fl. sessile in terminal and axil-

lary clusters.—*E. B.* 90.—St. 6—18 in. high. Bracts ovate-acuminate, shorter than the large erect flowers. Cal. hoary, with lanceolate segments. Cor. funnelshaped, large, deep blue, downy. L. often hoary beneath.—Dry calcareous pastures. P. VII. VIII. *Clustered Bell flower.* E. S.

\*\* *Caps. stalked, nodding; pores at the base.*

2. *C. latifolia* (L.); *l. ovate-lanceolate* acuminate doubly serrate hairy, lower ones stalked, upper l. nearly sessile, fl. racemose, peduncles 1-flowered, cal.-segments lanceolate-acuminate glabrous finely serrate, st. erect slightly angular.—*E. B.* 302. St. 72. 3.—St. 3—4 feet high, simple, leafy. Cor. very large, blue, glabrous, hairy within.—Woods and thickets chiefly in the North. P. VII. VIII. *Giant Bell flower.* E. S.

3. *C. Trachelium* (L.); *l. coarsely doubly serrate* hispid, lower ones cordate with long stalks, upper l. nearly sessile ovate or lanceolate-acuminate, fl. racemose, peduncles 2—3-flowered, cal.-segments triangular-lanceolate entire erect, st. erect angular.—*E. B.* 12.—St. 2—3 feet high, mostly simple, leafy. Cor. truly bellshaped, large, deep blue.—Hedges and thickets chiefly in the South. P. VII. VIII. *Nettle-leaved Bell flower.* E. I.

†4. *C. rapunculoides* (L.); *l. unequally crenate serrate* scabrous, lower ones cordate with long stalks, upper l. sessile lanceolate, fl. racemose unilateral, peduncles 1-flowered, cal.-segments linear-lanceolate entire at length reflexed, st. erect slightly angular, root creeping.—*E. B.* 1369.—St. 2 feet high, simple, leafy. Cor. pale blue.—Hedges, very rare. Near Kirkcaldy, Fifeshire. *Mr. J. B. Syme.* P. VII. VIII. E. S.

5. *C. rotundifolia* (L.); radical *l. cordate or reniform* shorter than their stalks, stem-*l. linear*, the lower ones lanceolate, fl. 1 or more racemose, cor. turbinate-campanulate.—*E. B.* 866.—St. 6—12 in. high. Radical l. soon vanishing. Cor. blue. Cal.-segments linear subulate equalling  $\frac{1}{3}$ -corolla.— $\beta$ . *lancifolia* (Koch); lower st.-l. rather broadly lanceolate, upper l. gradually smaller, fl. often solitary.— $\gamma$ . *arctica* (Lange); as  $\beta$ , but fl. much larger and more erect, cor.-lobes short. *Fl. Dan.* 2711.—Dry and hilly places.  $\beta$  on mountains.  $\gamma$  Innis Boffin, I. P. VII. VIII. *Hairbell.* E. S. I.

\*\*\* *Caps. stalked, erect; pores just below cal.-segments.*

[*C. persicifolia* (L.); l. smooth slightly serrate, root-l. obovate narrowed into a petiole, stem-l. linear-lanceolate sessile, raceme few-flowered, cal.-segments lanceolate.—*E. B. S.* 2773.

—St. 1—2 feet high. L. long, narrow, with very narrow serratures. Fl. very large, often solitary. Cal.-segments entire.—“Woods near Cullen, Banffshire, and Thorpe Arch, Yorkshire.” Not a native. P. VII.]

‡6. *C. Rapun'culus* (L.); l. crenate, root-l. elliptic-lanceolate narrowed into a petiole, stem-l. linear-lanceolate, panicle erect racemose, *cal.-segments subulate*.—*E. B.* 283.—St. 3 feet high, angular, rough. Fl. small, pale blue. Cal.-segments entire.—Sandy soil in the South. B. VII. VIII. *Rampion*. E.

7. *C. pat'ula* (L.); l. crenate, root-l. oblong-elliptic narrowed into a petiole, stem-l. linear-lanceolate, *panicles very lax, fl. on long stalks erect, cal.-segments toothed at the base subulate*.—*E. B.* 42.—St. 2 feet high, terminating in a very loose spreading panicle. Fl. purplish blue, funnelshaped, open.—Hedges and thickets. B. VII. VIII. E.

#### 5. SPECULA'RIA *Heist.*

1. *S. hybrida* (A. DC.); st. simple or branched, l. slightly crenate wavy oblong sessile, lower l. spatulate, cal. rough, its segments lanceolate longer than the cor. shorter than the ovary.—*Campanula* L. *E. B.* 375.—St. 3—12 in. high, rough with rigid minute hairs. Fl. few, terminal, solitary, small, lilac.—Corn-fields. A. VI.—IX. E. S.

#### 6. WAHLENBERG'IA *Schrad.*

1. *W. hederacea* (Reich.); l. roundish cordate angularly 5-lobed stalked alternate, st. filiform prostrate, peduncles solitary.—*Campanula* L. *E. B.* 73.—St. branched, slender, creeping greatly. Peduncles longer than the leaves. Fl. pale blue, bellshaped, narrow, at first nodding, afterwards erect. Cal.-segments subulate. Caps. nearly globose.—Damp peaty places in the South and West. P. VII. VIII. *Ivy-leaved Bell flower*. E. S. I.

### Order XLVII. ERICACEÆ<sup>1</sup>.

Cal. 4—5-parted, persistent. Cor. gamopetalous, 4—5-parted, usually regular and marcescent; or sometimes only slightly cohering below. Stam. 8—10, hypogynous. Anth. 2-celled, opening by 2 pores or slits and often with spurlike appendages at the base. Ovary surrounded by a disk or scales, free or adhering to the corolla. Fr. capsular or baccate, with several cells, many-seeded.—No stipules.

<sup>1</sup> Monotropeæ, Vacciniæ, and Pyrolaceæ are considered distinct orders by Maout and Decaisne.

\* *Anthers opening by pores.*

Tribe I. *ARBUTEÆ*. *Fr. baccate, fleshy, superior.* Disk and stam. hypogynous. Petals cohering.—L. usually broad.

1. *ARBUTUS*. Cal. 5-parted. Cor. globose or ovoid with a small contracted 5-cleft reflexed border, deciduous. Stam. 10, with flattened filaments. Anth. compressed, with 2 pores at the apex, fixed at the back beneath the apex and there furnished with 2 reflexed appendages. *Berry globose, tubercled; cells 5, many seeded.*

2. *ARCTOSTAPHYLOS*. *Fr. with 5 1-seeded cells, smooth.* Otherwise like *Arbutus*.

Tr. II. *ERICÆÆ*. *Fr. capsular, dry, superior.* Anth. 2-celled. Disk and stam. hypogynous. Testa close. Petals cohering.

3. *ANDROMEDA*. Caps. of 5 cells and 5 valves, dry. Otherwise like *Arbutus*.—L. usually broad.

4. *CALLUNA*. Cal. 4-parted, membranous, coloured, longer than the 4-cleft bellshaped persistent but fading corolla, surrounded by 4 green bracts. Stam. 8, with dilated filaments. Caps. 4-celled; *dissepiments adhering to the axis; valves opening at the dissepiments and separate from them.*

5. *ERICA*. Cal. 4-parted. Cor. bellshaped or ovoid, often ventricose, 4-toothed, persistent, fading. Stam. 8. Caps. 4-celled; *valves opening between the dissepiments and carrying a part with them.*

6. *PHYLLODOCE*. Cal. 5-parted. Cor. ovoid, deciduous; mouth contracted, 5-toothed. Stam. 10, included; *filaments slender, longer than the anthers; cells short, truncate, opening by pores at the apex. Stigma peltate, with 5 tubercles.* Caps. 5-celled with 5 valves opening at the dissepiments.

7. *DABEOCIA*. Cal. 4-cleft. Cor. ovoid, ventricose, limb 4-toothed. Stam. 8, included; *filaments flattened, shorter than the linear anthers, which are sagittate below; cells loosened and opening by oblique pores at the apex. Stigma simple, truncate.* Caps. 4-celled, with 4 valves opening at the dissepiments.

8. *AZALEA*. Cal. 5-parted. Cor. bellshaped, 5-cleft. Stam. 5, equal, shorter than the corolla; anth. roundish; *cells opening by a longitudinal fissure.* Stigma capitate. Caps.

2—3-celled with 2 or 3 bifid valves whose inflexed edges form the double partitions.

Tr. III. *VACCINIEÆ*. Fr. baccate, fleshy, inferior. Disk and stam. epigynous. Petals cohering.

9. *VACCINIUM*. Cal. entire or 4—5-toothed or lobed. Cor. 4—5-cleft or toothed. Stam. 8—10; anth. oblong, bifid at the summit. Berry globose, crowned by the persistent limb of the calyx, 4—5-celled, many-seeded.

Tr. IV. *PYROLEÆ*. Fr. capsular, dry, superior. Seeds with a loose testa. Disk 0. Stam. hypogynous. Anth. opening by pores. Petals scarcely cohering.

10. *PYROLA*. Cal. 5-parted. Cor. of 5 connivent petals. Stam. 10; anth. inverted, with 2 cells each opening by a round pore at the base. Style 5-lobed. Caps. 5-valved, opening from near the base to the top; margins of the valves connected by a web.

11. *MONESSES*. Cal. 5-parted. Cor. of 5 petals connected below. Stam. 10; anth. inverted, with 2 cells each furnished with a tubular horn opening at the end. Stigma 5-parted, radiant. Caps. 2-celled, 5-valved, opening from the top to the base with glabrous sutures.

\*\* *Anthers opening by a transverse fissure.*

Tr. V. *MONOTROPEÆ*. Fr. capsular, dry, superior. Seeds with a loose testa. Disk 0. Petals scarcely cohering.

12. *MONOTROPA*. Cal. 4—5-parted. Cor. of 4—5 petals each with a hooded honey-bearing base. Stam. 8—10. Anth. kidney-shaped, 1-celled, 2-valved. Stigma peltate. Caps. 5-celled, 5-valved, many-seeded.

### Tribe I. *Arbuteæ*.

#### 1. *AR'BUTUS* Linn. Strawberry-tree.

1. *A. Unédo* (L.); bark rough, l. elliptic-lanceolate serrate coriaceous glabrous, panicle terminal nodding, pedicels glabrous. —*E. B.* 2377.—An evergreen tree. Fl. whitish, pendulous. Fr. red.—Killarney, where it is truly wild. T. IX. X. I.

#### 2. *ARCTOSTAPH'YLOS* Adans.

1. *A. alpina* (Spr.); procumbent, l. thin wrinkled serrate



fading but persistent, clusters terminal.—*Arbutus* L. *E. B.* 2030. *St.* 6. 8.—St. woody, trailing, long. L. obovate, netted. Fl. white, hairy about the mouth. Berry smooth, black.—Dry barren spots on the Highland mountains. Sh. V. S.

2. *A. Uva-ur'si* (Spr.); procumbent, l. coriaceous obovate entire shining evergreen, clusters terminal.—*Arbutus* L. *E. B.* 714. *St.* 6. 8.—St. woody, trailing, long. L. blunt, quite entire, rigid. Fl. rose-coloured, smooth. Berry globose, scarlet, superior.—Dry stony mountain heaths. Sh. VI. E. S. I.

## Tribe II. *Ericææ*.

### 3. ANDROM'EDA Linn.

1. *A. polifolia* (L.); l. alternate lanceolate with revolute margins glaucous beneath, fl. clustered terminal.—*E. B.* 713.—St. slender, woody, prostrate below. Ped. variable in length. Fl. drooping ovate, pink, occasionally 4-fid and 8-androus. L. evergreen, acute. Peduncles 2 or 3 times as long as the flowers.—Peat bogs. Sh. V.—IX. E. S. I.

### 4. CALLU'NA Salisb. Ling.

1. *C. vulgaris* (Salisb.).—*E. B.* 1013.—A low tufted shrub. L. small, sessile, closely imbricate, patent, in 4 rows, keeled, each with 2 small spurs at the base, nearly or quite smooth. Fl. small, shortly stalked, drooping, lilac-rose-coloured or white, with the lower ped. leafy; sep. and pet. oblong erect; fl.-raceme ending in a leafy shoot. L. sometimes hoary.—The supposed *C. atlantica* is not constant to the characters recorded, even on the same bush.—Dry heaths. Sh. VI.—VIII. E. S. I.

### 5. ERICA Linn. Heath.

\* *Cor.* globose or urceolate, *stam.* included, filaments filiform flattened, *stigma* peltate. ERICA D. Don.

1. *E. Tet'ralix* (L.); l. 4 in a whorl lanceolate or linear ciliate downy above and on the midrib beneath with revolute edges, fl. in an umbellate head, sep. linear downy ciliate, anth. spurred, ovary downy.—*E. B.* 1014.—St. branched below and often especially about the middle, simple in the upper part, densely leafy below, the whorls more distant towards the top and usually leaving a leafless space next to the flowers. Tips of young shoots green. Young l. always downy above, old l. sometimes glabrous. Sep. downy and mealy. Fl. rose-coloured. Style

usually included.—*β. Watsoni*; cor. ventricose, fl. more or less racemose. *Sy. E. B.* 888. It may be a hybrid.—Boggy heaths. *β. Truro*, Cornwall. *Sh. VII. VIII.* E. S. I.

2. *E. Mackaiána* (Bab.); l. 4 in a whorl ovate ciliate the midrib beneath and upper surface glabrous with revolute edges, fl. in an umbellate head, sep. ovate lanceolate glabrous, anth. spurred, ovary glabrous.—*E. B. S.* 2900.—St. irregularly branched throughout, particularly above, densely and equally leafy quite up to the flowers. Tips of young shoots pink. L. and sepals quite without down; l. mealy beneath but the midrib bare; sep. with a small portion of meal near the apex beneath, otherwise quite bare. Fl. purplish. Style protruded.—I have seen forms of this taken for *E. ciliaris*.—Very wet moors between Roundstone and Clifden, and between Carna and Lough Sheedah, Co. Galway. *Sh. VIII. IX. Mackay's Heath.* I.

3. *E. cinérea* (L.); l. 3 in a whorl linear-lanceolate acute keeled beneath with a central furrow glabrous, fl. in dense whorled racemes, sep. linear-lanceolate smooth acute keeled, anth. spurred, ovary glabrous.—*E. B.* 1015.—St. with many upright branches. L. flat above, minutely serrulate. Fl. reddish purple.—Dry heaths. *Sh. VII. VIII. Fine-leaved Heath.* E. S. I.

4. *E. ciliaris* (L.); l. 4 in a whorl ovate ciliate with revolute edges, fl. in terminal unilateral racemes, anth. not spurred, mouth of the cor. oblique.—*E. B. S.* 2618.—St. long, straggling, ending in a long raceme of large oblong purple flowers and producing many short barren branches. Style protruded. Ovary glabrous.—Sandy heaths. Wareham, Dorset. Edgecombe Downs near Carclew, Cornwall. Mr. More, the author, and others have failed to find it near Clifden, Co. Galway. *Sh. VII. VIII.* E.

\*\* *Cor. bellshaped or shortly tubular, Stam. exerted, filaments flattened, style capitate.* GYPSOCALLIS D. DON.

5. *E. mediterránea* (L.); l. 4 in a whorl linear glabrous flat above convex with a central furrow beneath, decurrent line from the l. reaching but not extending beyond the next whorl, fl. axillary drooping racemose, cor. cylindrical-urceolate twice as long as the coloured calyx, anth. terminal not spurred opening throughout nearly their whole length.—*E. B. S.* 2774. *E. hibernica Syme.*—St. 2—5 feet high, with many upright rigid branches terminating in leafy racemes of flesh-coloured flowers but afterwards prolonged. L. many, erect-patent. Bracts above the middle of the pedicels. Stam. and style slightly exerted, style afterwards elongated. Ovary glabrous.—Mountain bogs. West of Mayo and Galway. *Sh. IV.* I.

6. *E. vagans* (L.); l. 4—5 in a whorl linear glabrous, fl. axillary crowded, cor. short bellshaped, sep. small ovate blunt, anth. lateral ovate of 2 distinct cells gibbous at the base.—*E. B.* 3. Probably *E. decipiens* St. Amand.—St. 1—2 feet high, copiously branched. Fl. usually collected in large numbers considerably below the top of the branches, cor. red or white. Anth. dark purple, not spurred. Ovary glabrous.—West of Cornwall. Sh. VII. VIII. *Cornish Heath.* E.

#### 6. PHYLLOD'OE *Salisb.*

1. *P. cærulea* (Bab.); l. linear denticulate, pet. glandular-hairy, calycine segments lanceolate acute, anth. two-thirds shorter than the glabrous filaments.—*Menziesia* Sm. *E. B.* 2469.—St. 4—5 in. high, branched, naked below, densely hairy above. Ped. terminal, all together, simple. Fl. large, pale bluish red.—Sow of Athol, Perthshire. Sh. VI. VII. S.

#### 7. DABEO'CIA *D. Don.*

1. *D. polifolia* (D. Don).—*Menziesia* Sm. *E. B.* 35.—St. bushy, 1—2 feet long, ultimately decumbent. L. ovate or elliptic, flat, with revolute edges, white and cottony beneath. Fl. large, purple, sometimes white, drooping, on short stalks, in terminal simple unilateral clusters. Anth. very large.—Conemara. Sh. VIII. *St. Dabeoc's Heath.* I.

#### 8. AZA'LEA *Linn.*

1. *A. procumbens* (L.).—*E. B.* 865. *Loiseleuria* Desv. The original and only *Azalea* of Linn.—St. woody, spreading procumbent. L. small, opposite, revolute. Fl. small, on simple stalks, terminal, collected together.—Summits of the Highland mountains. Sh. V. VI. S.

### Tribe III. *Vaccinieæ.*

#### 9. VACCIN'IUM *Linn.*

\* *Anth. with 2 dorsal horns. L. deciduous. St. erect.*

1. *V. Myrtillus* (L.); l. ovate serrate glabrous, fl. solitary, st. acutely angular.—*E. B.* 456.—St. woody, about a foot high, branching. Fl. subglobular, greenish tinged with red, nodding. Berries black.—Stony woods and heaths. Sh. V. *Bilberry.*  
E. S. I.

2. *V. uliginosum* (L.); l. obovate entire glaucous and veined beneath, fl. several together, st. rounded.—*E. B.* 581. *St.* 12.—*St.* woody. Fl. ovoid, flesh-coloured, nodding.—Berries black.—Mountain bogs. *Sh.* V. *Bog-Whortleberry.* E. S.

\*\* *Anth.* without horns on the back. *L.* evergreen.

3. *V. Vitis-idæa* (L.); l. obovate dotted beneath, margins revolute and somewhat crenate, fl. racemose terminal, cor. bellshaped.—*E. B.* 598.—*St.* woody, 6—8 in. high, straggling. *L.* like those of Box, dark green above. Fl. pink, 4-cleft. Berries red, inferior.—Mountain heaths. *Sh.* VI. VII. *Red Whortleberry. Cowberry.* E. S. I.

4. *V. Oxycoccos* (L.); l. ovate entire with revolute margins glaucous beneath, fl. terminal on long simple peduncles, cor. rotate with reflexed segments.—*E. B.* 319. *Oxycoccus palustris* DC.—*St.* procumbent, filiform, rooting. *L.* small. Fl. bright rose-colour. Cor. deeply divided, remarkably reflexed. Berries crimson.—Wet bogs. *Sh.* VI. VII. *Cranberry.* E. S. I.

[*V. macrocarpum* (Ait.); l. oblong with flat margins, fl. lateral on long simple peduncles.—Loughton Bog, Mold, Flintshire. An American plant, probably sown there.]

#### Tribe IV. *Pyroleæ.*

##### 10. PYR'OLA Linn. Winter-green.

1. *P. rotundifolia* (L.); l. nearly round entire or slightly crenate, fl. racemose, cal.-segments lanceolate acute, style bent down and curved upwards at the end longer than the ascending stam., stigma annular with 5 erect blunt points.—*E. B.* 213.—Fl. white, rather many, expanded. Style longer than the petals. Stam. all turned upwards. *L.* many.— $\beta$ . *arenaria* (Koch); l. smaller, st. with bracts throughout, cal.-segments shorter and broader. *Sy. E. B.* 896.—Damp bushy places and reedy marshes.  $\beta$ . Sand-hills near Lytham, Lancashire. P. VIII. E. S.

2. *P. média* (Sw.); l. nearly round or roundish-oval slightly crenate, fl. racemose, cal.-segments ovate-acute, stam. all regularly inflexed shorter than the nearly straight declining style, stigma annular with 5 erect points.—*E. B.* 1945.—Fl. milk-white tinged with pink, rather many, less expanded than in the preceding. Style projecting a little beyond the corolla, longer than the ovary, always nearly straight. *L.* many.—Woods in the North. P. VII. VIII. E. S. I.

3. *P. minor* (L.); l. roundish-oval crenate, fl. racemose, cal.-segments ovate-prolonged acute, *stam.* regularly inflexed *equalling the straight style, stigma without a ring* 5-lobed pointless.—*E. B.* 2543.—*St.* 13. 12.—Fl. pale pink, many, on very short pedicels, nearly closed. Style shorter than the ovary, included. L. many.—Mossy woods and thickets. P. VI. VII. E. S. I.

4. *P. secun'da* (L.); l. ovate acute serrate, *fl. in a secund raceme*, cal.-segments triangular rounded notched, *stam.* regularly inflexed equalling the *long straight style, stigma* 5-lobed without a ring or points.—*E. B.* 517. *St.* 13. 13.—Fl. white, drooping, oval-oblong, nearly closed. Style very long, exserted. L. many.—Mossy alpine woods. P. VII. E. S. I.

### 11. MONE'SES *Salisb.*

1. *M. grandiflora* (Salisb.).—*E. B.* 146. *Pyrola uniflora* L.—L. few, roundish, serrate. Fl. solitary, terminal, large, drooping, white, open, nearly an inch broad. *Stam.* shorter than the pet. and closely adpressed to them. Stigmas very large.—Woods in the north of Scotland, rare. P. VI. VII. S.

## Tribe V. *Monotropeæ.*

### 12. MONOT'ROPA *Linn.*

1. *M. Hypopitys* (L.); fl. in a drooping cluster, 8 *stam.* in terminal, 10 in lateral fl., fr. erect, bracts and fl. glabrous externally.—*E. B.* 69.—Inner side of the pet., filaments, germen and style glabrous (*Hypopitys glabra* Bernh., DC.) or hairy (*M. multiflora* Scop., DC.).—Plant 6—8 in. high, succulent, simple, clothed with ovate scales, terminating in a short cluster, dingy yellow, turning nearly black. Fl. with large scale-like bracts. Not parasitical. Clusters sometimes erect.—Woods. P. VII. VIII. *Yellow Bird's-nest.* E. S. I.

## Order XLVIII. AQUIFOLIACEÆ.

Sep. inferior, 4—9, imbricate. Cor. regular, 4—6-parted, imbricate. *Stam.* inserted upon the base of the corolla and alternate with its lobes. Disk 0. Ovary 2—6-celled; ovules solitary, pendulous, with a cupshaped seed-stalk. Fr. fleshy, not bursting; seeds stony, 2—6.—No stipules.

1. ILEX. Cal. 4—5-fid, persistent. Cor. rotate, 4—5-fid. *Stam.* 4—5. Stigmas 4—5, nearly sessile. Fr. fleshy, containing 4—5 seeds.

1. I'LEX *Linn.* Holly.

1. *I. Aquifolium* (L.); l. ovate acute spinous wavy shining, peduncles axillary short many-flowered, fl. somewhat umbellate.—*E. B.* 496. *St.* 7. 4.—A small tree. L. evergreen, often quite entire on the upper branches, edged with strong spinous teeth and terminated by a spine on the lower ones. Fl. white. Berries scarlet.—Woods and hedges. T. VI.—VIII. E. S. I.

## Order XLIX. OLEACEÆ.

Cal. gamosepalous, divided, persistent; or none. Cor. with 4—8 divisions, valvate, rarely 0. Stam. 2. Ovary free, 2-celled; ovules pendulous. Stigma entire or bifid. Fr. a berry, drupe, or capsule, often 1-seeded. Albumen dense or none.—No stipules.

1. LIGUSTRUM. Fr. fleshy, a berry containing 2 seeds. Cal. cupshaped with 4 minute teeth. Cor. funnelshaped; limb 4-cleft, spreading. Stam. 2.

2. FRAXINUS. Fr. (a samara) dry, of 1 or 2 single-seeded cells, compressed and leaflike at the end, pendulous. Cal. 0 or 4-cleft. Cor. 0, or of 4 petals. Cal. and cor. wanting in our plant.—Fl. sometimes with only stam. or with pistils only.

1. LIGUSTRUM *Linn.* Privet.

1. *L. vulgare* (L.); l. elliptic-lanceolate entire glabrous, panicles terminal compound dense.—*E. B.* 764. *St.* 14. 1.—A bushy shrub, 6—8 feet high, with straight smooth branches and opposite leaves. Fl. white. Berries globose, black, rarely yellow.—Thickets in the South of England. Sh. VI. VII. E. I.

2. FRAXINUS *Linn.* Ash.

1. *F. excelsior* (L.); l. pinnate with 4—8 pairs of nearly sessile ovate-lanceolate acuminate serrate leaflets, cal. wanting.—*E. B.* 1692. *St.* 44. 7.—A handsome tree. Usually dioecious. Fl. appearing before the l., in axillary clusters.—[*β. E. heterophyllus* (Vahl); l. simple and pinnate. *E. B.* 2476.]—Woods and hedges. T. IV. V. E. S. I.

## Order L. APOCYNACEÆ.

Cal. in 4 or 5 persistent, divisions. Cor. regular, 4—5-lobed,

deciduous, twisted in the bud. Stam. 5, filaments distinct. Anth. 2-celled, basifixed; pollen granular. Ovaries 2, 1-celled; or 1 of 2 cells. Stigma 1. Fr. 2 follicles. Seed with fleshy albumen.—Rudimentary stipules rarely seen.

1. *VINCA*. Cor. salvershaped; tube long, with 5 angles at the mouth, closed by spreading hairs and the connivent stamens; limb flat, 5-lobed. Stigma capitate with a ring at its base. Fr. of 2 erect long slender follicles or rarely one, 2 glands alternating with them.

### 1. *VIN'CA* Linn. Periwinkle.

†1. *V. minor* (L.); st. procumbent, l. lanceolate-elliptic, their margins as well as those of the small lanceolate calyx-segments glabrous.—*E. B.* 917.—St. rooting. Flowering branches erect. Fl. smaller than those of the next, blue, rarely white.—Woods and thickets. P. V. VI. *Lesser Periwinkle*. E.

[2. *V. major* (L.); st. somewhat ascending, l. ovate acute or subcordate, their margins as well as those of the long subulate calyx-segments ciliate.—*E. B.* 514.—St. at first ascending, afterwards prostrate and rooting. Flowering shoots erect. Fl. large, purplish blue.—Hedges and thickets, introduced. P. IV. V. *Greater Periwinkle*.] E.

## Order LI. GENTIANACEÆ.

Cal. inferior, divided, persistent. Cor. regular, 4—8-fid, hypogynous, marcescent; imbricate and twisted, rarely induplicate in the bud. Stam. inserted on the cor., as many as the segments. Anth. versatile. Ovary of 2 carpels with the edges slightly inflexed or meeting. Caps. many-seeded, generally two-valved.—No stipules.

Subord. I. GENTIANEÆ. Corolla twisted in the bud.  
L. opposite.

\* *Style deciduous*.

Tribe I. *CHLOREÆ*. Corolla rotate.

1. *CHLORA*. Cal. 8-parted. Cor. nearly rotate, 8-parted. Stam. 8. Style 1. Stigma 2—4-cleft. Caps. 1-celled placentas on the inflexed margin of the valves.

Tr. II. *ERYTHRÆEÆ*. Corolla funnel- or salver-shaped.

2. *ERYTHRÆA*. Cal. 4—5-fid. Cor. funnel-shaped, limb

short, 4—5-fid. *Stam.* 4—5. *Anth.* erect, at length spirally twisted. *Style* simple; stigmas 2. *Caps.* imperfectly 2-celled from the inflexed margins of the valves.

3. *CICENDIA*. *Cal.* 4-partite or lobed. *Cor.* salver-shaped, limb short, 4-fid. *Stam.* 4. *Anth.* erect, not twisted. *Stigma* capitate, undivided. *Caps.* 1- or imperfectly 2-celled.

\*\* *Style* persistent or *stigma* sessile.

Tr. III. *SWERTIÆ*. *Style* often wanting, *stigma* persistent.

4. *GENTIANA*. *Cal.* 4—5-cleft. *Cor.* funnel- or salver-shaped, limb 4—5-cleft. *Stam.* 4—5, straight. *Stigmas* 2. *Caps.* 1-celled, seeds on the inflexed margins of the valves.

Subord. II. *MENYANTHÆ*. *Corolla* induplicate in the bud. *L.* alternate, or opposite to the fl.-stems.

5. *LIMNANTHEMUM*. *Cal.* 5-parted. *Cor.* rotate, thin; limb 5-parted, smooth on the disk, hairy or scaly at the base within. *Stam.* 5. *Stigma* with two toothed lobes. *Caps.* 1-celled, not bursting.—*L. simple*, floating.

6. *MENYANTHES*. *Cal.* 5-parted. *Cor.* funnelshaped, fleshy; limb 5-parted hairy within. *Stam.* 5. *Stigma* capitate, notched. *Caps.* 1-celled, 2-valved; valves bearing the seeds along their middle.—*L. ternate*.

Suborder I. *Gentianeæ*. Tribe I. *Chloreæ*.

1. *CHLO'RA* *Linn.* Yellow wort.

1. *C. perfoliata* (L.); lowermost l. elliptic-oblong narrowed below, stem-l. broadly connate, st. perfoliate.—*E. B.* 60. *R.* xvii. 1060.—*St.* 12—18 in. high, simple. *Stem-l.* triangular-ovate, connected by their whole breadth in rather distant pairs, glaucous. *Panicle* forked, many-flowered. *Cal.* divided to its base into linear-subulate segments. *Cor.* bright yellow. *Stigmas* yellow.—Damp chalky places. *A.* VII.—IX. *E.* I.

Tribe II. *Erythrææ*.

2. *ERYTHRÆ'A* *Renealm.* Centaury.

\* *Stamens* from top of *cor.-tube*.

1. *E. pulchella* (Fries); st. much branched acutely quadrangular, l. ovate the uppermost oblong lanceolate, fl. all



stalked axillary and terminal, *cal.* rather shorter than the tube of the opening corolla, cor.-lobes elliptic-oblong blunt.—*E. B.* 458. *Sy. E. B.* 910 b.—St. quite simple, 1 in. high and 1-flowered; or very much branched, 6—8 in. high, with very many flowers. Radical l. very few. Panicles forked, a fl. in the fork; *lateral fl.* distant from the floral leaves. Fl. rose-coloured.—“*β. E. tenuiflora* (Link); branches erect forming a term. fastigiate lengthened corymb.” *Towns. MS.*—In each species the length of the tube must be observed exactly when the flower is opening.—Sandy ground. *β.* Low ground between Cowes and Newport, Isle of Wight. *Townsend.* A. VII.—IX. E. S. I.

2. *E. Centaurium* (Pers.); st. branched above quadrangular, l. elliptic-oblong, the upper ones acute, *fl.* nearly sessile corymbosely paniced, *cal.* not half as long as the tube of the opening corolla, cor.-lobes oval.—*Sy. E. B.* 909.—St. 6—18 in. high, usually simple below. Panicles of fl. lax. *Lateral fl.* apparently stalked, but sessile between the two small floral leaves. Radical l. many.—Dry pastures. A. VII. VIII. E. S. I.

3. *E. latifolia* (Sm.); st. short simple 3-cleft at the top, l. broadly oval blunt 5—7-veined, *fl.* in compact round dense forked term. tufts subsessile, *cal.* about equalling cor.-tube, cor.-lobes lanceolate.—*Sy. E. B.* 907.—St. about 3 in. high. Fl.  $\frac{1}{2}$  as large as in Sp. 2. Lowest l. sometimes almost orbicular.—Sands by sea near Liverpool, very rare. A. VII. E.

4. *E. littoralis* (Fries); st. simple solitary or several from the crown of the root, l. oblong-linear blunt narrowed below, radical l. crowded spatulate, *fl.* sessile between the floral l. densely corymbose, *calyx* as long as the tube of the opening corolla, cor.-lobes oval blunt.—*Sy. E. B.* 908. *E. chloodes* Gren., not *E. linariifolia*.—St. 2—6 in. high. Corymb usually trichotomous, dense; branches sometimes long. Fl. rose-coloured. Radical l. narrow, many.—Sandy sea-shores. A. VII. VIII. E. S.

\*\* *Stamens* from base of cor.-tube.

5. *E. capitata* (W.); st. short simple, st.-l. ovate or subspathulate blunt 3-veined, fl. sessile in compact term. tufts, *cal.* about equalling cylindrical cor.-tube, cor.-lobes oval blunt.—*Linn. Journ.* xviii. t. 15.—St. less than 3 in. high. Fl.-tufts sessile; there are often from outer bracts a few long-stalked tufts overtopping primary tuft. Root-l. in rosette, with 3 long and often 2 short veins. *Cor.-tube* not narrowed at top. Stam. from quite base of cor.-tube, otherwise free.—Downs at Freshwater, Isle of Wight. A. or B. VII. VIII. E.

3. CÍCEN'DIA *Adans.*

1. *C. filifor'mis* (Reich.); cal. bellshaped with 4 ovate acute lobes, st. threadshaped forked.—*Exacum* Sm. *E. B.* 235. *Microcala* Griseb.—St. 1—4 in. high. Radical l. linear-lanceolate, stem-l. subulate, all sessile. Fl. yellow, solitary, on long stalks.—Damp sandy places. A. VII. VIII. E. I.

[*C. pusil'la* (Griseb.); cal. 4-parted with linear segments, st. slender branching from its base.—*E. B. S.* 2994.—St. much branched throughout, 1—4 in. high. L. all narrowly linear-lanceolate. Fl. pink.—On spots sometimes flooded. Paradis, Guernsey. *Capt. Gosse*lin. A. VI. VII.]

Tribe III. *Swertiae*.4. GENTIA'NA *Linn.* Gentian.

\* *Cor. funnel- or somewhat salver-shaped.*

1. *G. Amarell'a* (L.); cor. 5-cleft hairy in the throat, callobes 5 nearly equal lanceolate, l. sessile ovate-lanceolate, radical l. oval-spathulate.—*E. B.* 236. *Sy. E. B.* 917.—Very variable in size and in the number of the flowers, 3—12 in. high, erect. St. square, much branched. Fl. rarely 4-cleft. Caps. stalked or sessile. Cor.-tube obconical or subcylindrical.—*a. G. Amarella* (Sm.); branches of st. erect, cal.-segm. nearly equal, cor.-tube cylindrical a little exceeding calyx, fl. lurid-purple. An early flowering form has 4-cleft cor. and unequal sep.—*β. G. germanica* (Willd.); st. much branched ascending, cal.-segm. unequal, cor.-tube obconical much exceeding calyx, fl. bluish-lilac. *J. of B.* ii. t. 15.—Dry calcareous fields. A. VIII. IX. *Felwort.* E. S. I.

2. *G. campes'tris* (L.); cor. 4-cleft hairy in the throat, callobes 4, 2 outer ones very large ovate, l. ovate-lanceolate.—*E. B.* 237.—St. 3—10 in. high. Fl. pale lilac; cor.-tube slightly thicker upwards. Caps. nearly sessile. Upper l. and sepals pointed.—Dry limestone hills. A. VIII. IX. E. S. I.

3. *G. nivális* (L.); cor. 5-cleft with minute intermediate bifid lobes, throat naked, cal. cylindrical with 5 equal lobes and keeled angles, l. ovate lowermost broadly elliptic.—*E. B.* 896.—St. erect, slightly branched, 1—6 in. high. Fl. bright blue.—Top of Highland mountains, very rare. A. VIII. S.

4. *G. ver'na* (L.); cor. 5-cleft with small intermediate bifid lobes, throat naked, cal. with prominent angles and sharp teeth,

l. ovate lower ones crowded, *st. cæspitose single-flowered* with 1 or 2 pairs of leaves.—*E. B.* 493. *St.* 40. 12.—*St.* prostrate, rooting, each ending in a roselike tuft of l. and a single short flowering shoot. Fl. rather large, vivid blue.—Barren limestone districts. Teesdale, Durham. Burrin, Co. Clare; Gort, Galway, Tuam, &c. *P.* IV.—VI. E. I.

\*\* *Cor.* bellshaped, its throat naked.

5. *G. Pneumonanthæ* (L.); *cor.* 5-cleft, cal. entire with linear blunt lobes, *fl. mostly solitary* slightly stalked, *l. linear* blunt.—*E. B.* 20.—*St.* 4—10 in. high, leafy, simple, erect or ascending. Fl. very large, deep blue within and with a broad greenish band down the middle of each segment.—Moist turfy heaths. *A.* VIII. IX. E.

## Suborder II. *Menyanthææ*.

### 5. LIMNANTHEMUM *Gm.*

1. *L. nymphæoides* (Link); l. roundly heartshaped floating wavy at the edges, ped. clustered 1-fl., *cor.* ciliate.—*Villarsia* Vent. *E. B.* 217. *R.* xvii. 1042.—Floating. *St.* long, round, branched. L. resembling those of *Nymphæa alba*, but much smaller. Fl. yellow. Caps. sometimes 3-valved.—Still places in rivers, rare. *P.* VII. VIII. E.

### 6. MENYANTHES *Linn.* Buckbean.

1. *M. trifoliata* (L.).—*E. B.* 495.—*St.* ascending, round, leafy. L. ternate. Leaflets equal, obovate, wavy. Raceme long-stalked, opposite to a leaf, many-flowered. *Cor.* flesh-coloured, densely fringed within.—Boggy places. *P.* V.—VII. *E.* S. I.

## Order LII. POLEMONIACEÆ.

*Cal.* inferior, 5-parted, persistent. *Cor.* hypogynous, regular, 5-lobed. *Stam.* 5, unequal, on the tube of the corolla. *Ovary* 3-celled. *Stigmas* 3-fid. *Caps.* 3-celled, 3-valved; valves separating at the axis.—No stipules.

1. POLEMONIUM. *Cal.* 5-fid. *Cor.* rotate, with a short tube and 5-lobed limb; throat nearly closed by the dilated bases of the filaments.

### 1. POLEMONIUM *Linn.* Jacob's Ladder.

1. *P. cæruleum* (L.); *st.* angular, l. glabrous pinnate, leaflets

ovate-lanceolate pointed, panicle downy glandular.—*E. B.* 14.—St. 1—2 feet high, simple, hollow. L. alternate, leaflets many. Fl. many, somewhat drooping, bright blue or white.—Bushy hilly places, rare. P. VII. E. S.

### Order LIII. CONVULVULACEÆ.

Cal. inferior, of 5 persistent imbricate often unequal sepals. Cor. hypogynous, regular, plaited, deciduous. Stam. 4—5, from near the base of the corolla. Ovary of 2—4 cells, few-seeded, surrounded by an angular hypogynous disk. Style 1, rarely 2. Caps. with the valves separating from the edges of the dissepiments or bursting transversely.—No stipules.

\* *With leaves and leaflike cotyledons.*

1. CONVULVULUS. Cor. bellshaped, with 5 prominent plaits and 5 shallow lobes. Style simple; stigmas 2. Caps. 2—4-celled; cells 2-seeded.

\*\* *Without leaves or cotyledons.*

2. CUSCUTA. Cal. 4—5-cleft. Cor. roundish-urceolate or bellshaped, 4—5-parted, with as many scales alternating with the segments at the base within. Stam. 4—5. Styles 2, rarely 1. Caps. bursting transversely, 2-celled, 4-seeded.

#### 1. CONVULVULUS Linn. Bindweed.

\* *Bracts minute, distant from the flower.*

1. *C. arven'sis* (L.); l. ovate- or strapshaped-hastate, peduncles mostly 1-flowered.—*E. B.* 312.—St. many, angular, twining or prostrate, leafy, branched. Peduncles sometimes 2-flowered. Cor. beautifully variegated with pink and white. Caps. 2-celled. Roots descending remarkably deep. Plant glabrous, or st. and l. downy.—Fields and hedges. P. VI.—VIII. E. S. I.

\*\* *Bracts large, close to the flower.* CALYSTEGIA R. Br.

2. *C. sépium* (L.); l. ovate- or triangular-hastate, peduncles 1-flowered square.—*E. B.* 313.—St. twining, many feet long, with large rather distant leaves. Fl. solitary, axillary, large, white, rarely pink. Bracts quite enclosing the calyx. Fr. imperfectly 2-celled through the shortness of the dissepiment.—Hedges and thickets. P. VII. VIII. E. S. I.

3. *C. Soldanel'la* (L.); l. reniform slightly angular fleshy, peduncles 1-flowered with 4 membranous angles.—*E. B.* 314.—*St. short, procumbent.* Fl. large, solitary, axillary, very handsome, pink with yellow bands. Bracts rather shorter than the calyx.—Sandy sea-shores. P. VI.—VIII. *Sea-side Bindweed.*  
E. S. I.

## 2. CUSCU'TA Linn. Dodder.

1. *C. europæ'a* (L.); clusters of fl. sessile, cor.-tube cylindrical afterwards ventricose, *scales adpressed* to inside of tube bifid distant below with rounded spaces between them, *cal. much shorter than corolla.*—*E. B.* 378.—*St. threadshaped, branching, reddish.* Fl. in rather large clusters, yellowish.—Parasitical upon herbaceous plants. A. VIII. IX. *Greater Dodder.* E. S.

[*C. Epilinum* (Weihe); clusters of fl. sessile, cor.-tube ventricose, *scales adpressed* fringed with teeth distant below with rounded spaces, *cal. with fleshy segments deltoid below nearly as long as the cor.-tube.*—*E. B. S.* 2850. *C. densiflora* Soy.-Willm.—*St. slender, nearly simple, pale green.* Fl. in rather small distant clusters, whitish. Scales bifid, with 4—8 teeth on each lobe. Ventricose cor.-tube with 5 longitudinal prominences; segm. ventricose. Styles at first erect, soon bowing outwards; stigmas converging.—Parasitical upon Flax and very injurious to the crop. A. VIII. *Flax Dodder.*] E. S. I.

2. *C. Epithymum* (Murr.); *st. twining irregularly,* clusters of fl. sessile, cor.-tube cylindrical, *scales converging usually equalling the tube of the cor.* fringed with teeth and rounded at the end close together below with narrow acute spaces, *cal. bellshaped shorter than the cor.-tube.*—*E. B.* 55.—*St. slender, red.* Fl. small, with a reddish thin cal., and white cor. with spreading ovate-acute segments. Sep. broad, ovate-apiculate, longer than their tube, with patent tips. Anth. blunt or notched at the end. Scales broad; the connecting membrane adpressed throughout.—[See *C. Ulicis* (Godr.); scales less than in Sp. 2 or 3, less deeply fringed converging. *Bull. Bot. Fr.* xxiii.]—Parasitical upon small shrubby plants. A. VII.—IX. *Lesser Dodder.* E. S.

†3. *C. Trifolii* (Bab.); *st. clasping like a ring,* clusters of fl. sessile, tube of the cor. cylindrical, *scales converging usually equalling half the tube of the cor.* fringed with teeth and rounded at the end distant below with rounded spaces, *cal. narrowed below about as long as the cor.-tube.*—*E. B. S.* 2898. Not *C. Epithymum* β. *Trifolii* R. xviii. 1342.—*St. slender, branching, reddish yellow, forming dense broad circular patches.* Fl. small, white.

Cal. fleshy, usually tipped with red; sep. lanceolate, about as long as their tube, adpressed. Anth. apiculate. Scales narrow; connecting membrane not adpressed, forming cuplike spaces between itself and the corolla.—Parasitical upon Clover chiefly. A. VII.—IX. *Clover-Dodder*. E.

[*C. approximata* (Bab.); like Sp. 2 but scales divided by acute spaces and truncate, was found on Bockhara Clover. *A. N. H.* xvi. t. i.; and *C. hassiaca* (Pfeiff.); *fl. stalked*, cor.-tube bellshaped closed with convergiug scales, stig. capitate, Heliotrope-scented, on Lucerne. Both introduced. A. VIII. IX.]

#### Order LIV. BORAGINACEÆ.

Fl. mostly in scorpioidal cymes, symmetrical. Cal. inferior, 5-parted, persistent. Cor. hypogynous, usually regular. Stam. 5, inserted on the corolla. Ovary of two 2-parted carp., with each lobe 1-seeded; ovules pendulous. Style simple, from near the base between the lobes of the ovary. Fr. separating in 4 nutlets or 2 bilocular portions. Seeds without albumen.—L. alternate.—Stip. 0.

Tribe I. *CYNOGLOSSÆ*. Nutlets 4, on the persistent base of the style.—Stam. included.

1. *ASPERUGO*. Cal. 5-cleft with alternate smaller teeth, enlarged and compressed in fruit. Cor. funnelshaped with rounded scales in the throat. Filaments of stam. short. *Nutlets* tubercled, *compressed*, attached by their narrow side, covered by the compressed calyx.
2. *CYNOGLOSSUM*. Cal. 5-cleft. Cor. funnelshaped, the mouth closed with prominent blunt scales. Filaments very short. *Nutlets roundish-ovate*, depressed, muricate, attached by the upper part of their inner edge.

Tr. II. *ANCHUSEÆ*. Nutlets 4, on an hypogynous disk, with an excavated space surrounded by a tumid ring at their base.

3. *BORAGO*. Cal. in 5 deep segments. *Cor. rotate*; tube very short; throat with short erect emarginate scales. *Stam. exerted*; filaments bifid, the inner fork bearing the anther; anthers linear-lanceolate, connivent in the form of a cone.
4. *ANCHUSA*. Cal. 5-fid. *Cor. funnelshaped with a straight tube*; throat closed by prominent blunt scales. *Stam. included, sessile*. Nutlets depressed.

5. **LYCOPSIS.** Cal. in 5 deep segments. Tube of the cor. curved; limb oblique. Otherwise like *Anchusa*.
6. **SYMPHYTUM.** Cal. 5-cleft or 5-parted. *Cor. cylindrical-bellshaped, throat closed by a prominent cone of connivent lanceolate-subulate scales.* Stam. exerted from the tube but covered by the scales; filaments short. Nutlets ovate.
- Tr. III. **LITHOSPERMEÆ.** Nutlets 4, affixed to an hypogynous disk, their base not excavated but attached by a flat or rather convex surface.
7. **ECHIUM.** Cal. in 5 deep segments. *Cor. sub-bellshaped; throat dilated, naked;* limb irregular. *Stam. exerted; filaments very long, unequal.* *Style bifid.* Nutlets wrinkled, attached by a flat triangular base.
8. **PULMONARIA.** Cal. tubular, 5-fid. *Cor. funnelshaped, its throat naked.* *Stam. included in the tube; filaments very short.* *Style simple.* *Nutlets smooth, attached by their truncate base, which has a central tubercle.*
9. **MERTENSIA.** Cal. in 5 deep segments. *Cor. bellshaped, with a short thick cylindrical tube with 5 minute protuberances in its throat.* *Stam. protruded beyond the throat; filaments rather long.* *Style simple.* *Nutlets smooth, inflated, rather drupaceous, attached laterally near their base by a flat surface; seeds free.*
10. **LITHOSPERMUM.** *Cal. in 5 deep segments.* *Cor. funnelshaped; throat naked or with 5 minute scales.* *Stam. included in tube; filaments very short.* *Style simple.* *Nutlets smooth or tubercular, stony, attached by a truncate flat base.*
11. **MYOSOTIS.** Cal. 5-parted. *Cor. convolute in the bud, salvershaped; throat closed with scales; limb 5-fid, blunt.* *Stam. included; filaments very short.* *Style simple.* *Nutlets smooth, convex externally, keeled within, attached by a minute lateral spot near their base.—Distinguished from all the other genera by the convolute corolla.*

### Tribe I. *Cynoglosseæ.*

#### 1. ASPERUGO Linn. Madwort.

†1. *A. procumbens* (L.).—*E. B.* 661.—St. procumbent, angular, rough, with short deflexed bristles. L. oblong, rough, hispid, lower ones stalked, upper sessile. Fl. small, axillary,

solitary, blue, upon short peduncles. Cal. of the fr. much enlarged.—Rich waste ground, rare. A. VI. VII. E. S.

## 2. CYNOGLOSSUM Linn. Hound's-tongue.

1. *C. officinale* (L.); l. downy acute, lower l. elliptic contracting into a petiole, upper l. lanceolate narrowed below subcordate half-clasping, nutlets with thickened prominent margin.—*E. B.* 921.—About 2 feet high. Covered with soft adpressed hairs. Cor. dull crimson, veiny; veins disappearing in drying. Nutlets flat in front. Fetid, rarely subglabrous and nearly scentless.—Waste ground. B. VI. VII. E. S. I.

2. *C. montanum* (Lam.); l. slightly hairy acute nearly glabrous and shining above rough beneath, inferior oblong narrowed into a long petiole, upper l. lanceolate slightly narrowed below clasping, nutlets without thickened edge.—*C. sylvaticum* Sm. *E. B.* 1642. Clothed with straight spreading hairs.—Cor. reddish, changing to blue. L. sometimes very rough.—Shady situations. B. VI. VII. E. I.

## Tribe II. Anchuseæ.

### 3. BORA'GO Linn. Borage.

\*1. *B. officinalis* (L.); lower l. obovate blunt attenuated below, segments of the cor. ovate acute flat spreading.—*E. B.* 36.—Fl. blue. Anth. very prominent. Stem-l. much narrowed below so as to appear stalked, eared at the base. Whole plant hispid with bulbous hairs. St. spreading, 12—18 in. high.—On rubbish and waste ground. B. VI. VII. E. S.

### 4. ANCHU'SA Linn. Alkanet.

†1. *A. officinalis* (L.); l. lanceolate hispid, spikes crowded unilateral, bracts ovate-lanceolate, cal.-segm. blunty hairy on both sides, scales of cor. hairy.—*E. B.* 662.—Fl. deep purple. Cal.-segm. narrow, longer than tube. St. 1—2 feet high, rough with deflexed hairs.—Waste ground, rare. P. VI. VII. E. S.

†2. *A. semper-virens* (L.); l. ovate, lower l. on long stalks, peduncles axillary each bearing 2 dense spikes with an intermediate flower, cal.-segments hairy on the outside only, bracts minute lanceolate, scales of the cor. downy.—*E. B.* 45.—Fl. blue, rather salver- than funnel-shaped. Cal.-segments narrow. St. 1½—2 feet high, rough with spreading somewhat deflexed hairs.—Waste ground near ruins, rare. P. V.—VIII. E. I.



## 5. LYCOP'SIS Linn. Bugloss.

1. *L. arvensis* (L.); l. lanceolate repand-dentate very hispid, cal. of fr. bellshaped erect.—*E. B.* 938.—Fl. small, blue. Whole plant very hispid with strong hairs each rising from a scaly tubercle.—Fields and hedges. A. VI. VII. E. S. I.

## 6. SYM'PHYTUM Linn. Comfrey.

1. *S. officinale* (L.); l. ovate-lanceolate attenuate below, stem-l. very decurrent lanceolate, st. winged in the upper part.—*E. B.* 817.—Height 1—2 feet, branching. Racemes in pairs, drooping. Fl. yellowish white or purple. Cal.-segments somewhat spreading and pubescence rougher in the usually purple-flowered variety, *S. patens* Sibth. *Sy. E. B.* 1116.—Common in damp places. P. V. VI. E. S. I.

2. *S. tuberosum* (L.); l. ovate-oblong attenuate below, stem-l. lanceolate, uppermost slightly decurrent, st. scarcely winged nearly simple.—*E. B.* 1502.—St. 12—18 in. high. Fl. yellowish white, whole plant smaller and slenderer than the preceding. Anth. twice as long as their filaments.—Damp woods and river-banks, rare. P. VI. VII. E. S.

[*S. peregrinum* (*asperinum* Bab.) Bot. Mag. 6466, *S. tauricum*, and *S. orientale* have been noticed in England, but are not natives.]

## Tribe III. Lithospermeæ.

## 7. E'CHIUM Linn. Viper's Bugloss.

1. *E. vulgare* (L.); tubercular hispid, st. erect simple, l. lanceolate 1-ribbed, stem-l. narrowed below sessile, fl. in four lateral scorpioidal cymes, stam. exceeding the corolla.—*E. B.* 181.—St. 1—2 feet high. Lower l. narrowing into a footstalk. Fl. reddish, then bright blue.—Dry places. B. VI. VII. E. S. I.

2. *E. plantagin'eum* (L.); pilose-hispid, st. erect branched diffuse, lower branches prostrate, radical l. oblong-ovate stalked, stem-l. oblong narrowed from a cordate half-clasping base with lateral ribs, spikes paniced long simple, stam. scarcely exceeding the corolla.—*E. B. S.* 2798.—Stam. very unequal, 1 short, 2 intermediate, and 2 longer. Fl. violet-blue. Root reddish.—Jersey. Land's-end, Cornwall. B. VI.—IX. E.

## 8. PULMONA'RIA Linn. Lungwort.

[*P. officinalis* (L.); l. ovate-lanceolate with a long haft, upper l. oblong-sessile.—*Sy. E. B.* 1098.—Fl. pale purple. St. 1 ft.

high. L. spotted.—Woods and thickets, scarcely naturalized. P. IV. V.] E.

1. *P. angustifolia* (L.); l. narrow-lanceolate narrowed to the base, upper l. sessile.—*E. B.* 1628.—Fl. pink, then blue. St. 1 ft. high. L. less frequently spotted.—Thickets. P. IV. V. E.

### 9. MERTEN'SIA *Roth.*

1. *M. maritima* (Gray); st. procumbent branched, l. ovate acute rough with hard dots glabrous fleshy glaucous, nutlets smooth.—*E. B.* 368.—Spreading, very glaucous. Fl. in cymes, purplish blue. Protuberances in throat of cor. yellow. L. tasting like oysters. Nutlets free, forming a pyramid, exceeding the calyx. Pericarp membranous; seeds smaller than the cavity.—Northern sea-shores. P. V.—VIII. E. S. I.

### 10. LITHOSPER'MUM *Linn.* Gromwell.

1. *L. officinale* (L.); st. erect much branched, l. lanceolate acute veined hispid with bulbous adpressed bristles above hairy beneath, throat of the cor. with a crown of scales, nutlets smooth.—*E. B.* 134.—St. 12—18 in. high. Fl. greenish yellow, nutlets grey, highly polished, stony, 2 or 3 ripening in each calyx. Root whitish.—Dry and stony places. P. VI.—VIII. *Gromwell.* E. S. I.

2. *L. purpureo-ceruleum* (L.); barren shoots prostrate creeping, fl.-st. erect forked, l. lanceolate acute hispid, hairs on the upperside adpressed bulbous, cor. much exceeding the calyx, nutlets smooth.—*E. B.* 117.—Fl. large, bright blue. Nutlets "silvery white, highly polished, slightly rugged," rarely produced.—This species and the following have no scales in the throat of the cor., but merely 5 longitudinal downy folds (*Rhyn-tispermum* Link).—Thickets on a limestone soil, rare. P. V.—VII. E.

3. *L. arven'se* (L.); st. erect branched, l. lanceolate rather acute hispid with adpressed bulbous hairs on both sides, cor. rather exceeding the calyx, nutlets tubercled.—*E. B.* 123.—St. 1 ft. high. Fl. small, white. Nutlets pale brown, polished, pitted, 3 or 4 perfected in each calyx. Root bright red.—Corn-fields. A. V.—VII. *Corn Gromwell.* E. S. I.

### 11. MYOSO'RIS *Linn.* Scorpion-grass.

i. Persistentes. *Cal. of fruit shorter than its stalk.*

\* Hairs on cal. all straight and adpressed.

1. *M. palustris* (With.); fr.-cal. open, its teeth short tri-

angular, cor.-limb flat longer than the tube, its lobes slightly emarginate, style about equalling cal., pubescence of st. (usually) spreading.—*E. B.* 1973.—*L.* bluntish, apiculate. Raceme quite leafless. Cor. large, bright blue with a yellow eye. Cal. divided about  $\frac{1}{3}$  down.—*a*; rhizome long leafy stoloniferous, st. angular with decurrent prominent lines, pubescence spreading.— $\beta$ . *M. strigulosa* (R.); rhizome short leafless, st. more slender and erect not angular its pubescence adpressed, l. less blunt, style equalling cor.-tube. *M. coronaria* Dum. Base of st. sometimes patently hairy.—Ditches and watery places. *P. VI.*—*VIII.*  $\beta$ . *VI.* *Forget-me-not.* *E. S. I.*

2. *M. répens* (Don); fr.-cal. open, its teeth narrow lanceolate acute, style falling short of cal., cor.-limb flat longer than the tube, its lobes slightly emarginate, pubescence of the st. spreading.—*E. B. S.* 2703.—*L.* rather acute. Stoloniferous. St. slightly angular. Raceme usually slightly leafy (1—4 leaves) below. Cor. pale blue. Cal. divided fully halfway down.—Boggy places. *P. VI.*—*VIII.* *E. S. I.*

3. *M. cæspitosa* (Schultz !); fr.-cal. open, its teeth narrow lanceolate bluntish, cor.-limb equalling the tube, its lobes entire, style very short, pubescence of the st. adpressed.—*E. B. S.* 2661. *M. lingulata* *Lehm.* (name only).—*L.* usually blunt or even emarginate. St. round, with a decurrent line. Raceme usually slightly leafy below. Cor. smaller than in the preceding, bright blue, segments narrower and rounded at the end. Style about equalling the cal.-tube.—Watery places. *P. VI.*—*VIII.* *E. S. I.*

\*\* Hairs on cal. not all straight, but some or all hooked.

4. *M. alpes'tris* (Schm.); cal. attenuate below deeply 5-cleft open with fruit with straight and a few curved adpressed bristles, pedicels ascending, cor.-limb longer than the tube flat, style equalling  $\frac{1}{2}$  cal., nutlet keeled not rounded at the end, root-l. on long stalks pointed.—*E. B.* 2559. *M. rupicola* Sm.—*L.* oblong-lanceolate, stalks of the lower ones slender. Fl. large, handsome, blue, sweet-scented in the evening.—Summits of the Breadalbane mountains. Micklefell, Teesdale. *P. VII. VIII.* *E. S.*

5. *M. sylvat'ica* (Ehrh.); cal. rounded below  $\frac{3}{4}$  5-cleft closed with fruit, its tube with spreading hooked bristles, pedicels divergent, cor.-limb longer than tube flat, style nearly equalling cal., nutlet keeled on one side upwards bluntish, root-l. on short dilated stalks bluntish.—*E. B. S.* 2630.—*L.* oblong-lanceolate; stalks of the oblong-ovate lower l. dilated. Fl. large, handsome, blue. Cal. divided more than halfway down.—Shady places, rare. *P. V. VI.* *E. S.*

6. *M. arven'sis* (Lehm.); *cal. half-5-cleft* closed with fruit, tube with spreading hooked bristles, pedicels divergent, *cor.-limb equalling tube concave*, cor.-lobes entire, *style very short*, racemes stalked.—*E. B. S.* 2629. *M. intermedia* Link.—L. oblong, acute; lower l. oblong-obovate, blunt. Fl. usually small. A large-flowered (*β. umbrosa*) plant found in shade is often taken for *M. sylvatica*.—Cultivated land and thickets. A. VI.—VIII. E. S. I.

ii. Fugaces. *Cal. of fruit not shorter than its stalk, its tube with spreading hooked bristles.*

7. *M. collina* (Hoffm.); *fr.-cal. open and ventricose* as long as the diverging pedicels, cor.-limb shorter than exerted tube concave, *style about equalling  $\frac{1}{2}$  cal.*, racemes stalked usually with 1 distant flower, fr.-pedicels spreading, hairs on the l. straight.—*E. B.* 2558. *St.* 42. 11. *M. hispida* Koch.—Usually slender, erect; or cæspitose with prostrate branches. L. oblong, blunt; lower obovate. *Fl.* small, unchangeably blue.—[*M. stricta* Link, *M. arvensis* (R.) *St.* 42. 14, has its fr.-cal. closed, very short adpressed fr.-pedicels, sessile racemes leafy below. Lowest fr.-ped. not  $\frac{1}{2}$  as long as calyx. Probably a native of Britain.]—Dry banks. A. IV. V. E. S. I.

8. *M. versic'olor* (Reich.); *fr.-cal. closed oblong bellshaped* longer than the ascending pedicels, cor.-limb much shorter than tube concave, *style equalling cal.*, racemes stalked.—*E. B.* 480.—*St.* erect, simple at the base. L. narrow, oblong, acutish; upper ones frequently opposite. *Fl.* small, at first *pale yellow*, afterwards blue. Lobes of fr.-cal. often erect, therefore cal. not truly closed.—On plants in damp places the fl. are at first white and the cal. is "less deeply divided;" sometimes the whole plant is yellowish and the fl. yellow.—Meadows and banks. A. V. VI. E. S. I.

#### Order LV. SOLANACEÆ.

*Cal. inferior, 5-parted, persistent. Cor. hypogynous, regular or slightly irregular, 5-cleft, deciduous, plicate in bud; the lobes imbricate or imbricate-plicate (in Solanum valvate). Stam. 4—5, inserted on the cor., alternate with the lobes. Ovary 1—2- or imperfectly 4-celled. Fr. a berry or capsule. Seeds many. Embryo usually curved, in fleshy albumen, often not in the axis.—No stipules.*

Tribe I. *SOLANEÆ*. *Cor. rotate; lobes nearly regular and equal, valvate in the bud. Anth. opening by pores.*

1. *SOLANUM*. *Cor.-limb 5-cleft, reflexed. Anth. erect connivent. Berry roundish with 2 or more cells.*

- Tr. II. *ATROPEÆ*. Cor. tubular; tube plicate in bud; lobes slightly unequal, imbricate in the bud. Anth. opening longitudinally at the margin.
2. *ATROPA*. Cor. bellshaped with 5 equal lobes. Cal. 5-parted, patent and dilated with fruit. Stam. included. Fr. a globose 2-celled berry.
  3. *HYOSCYAMUS*. Cor. funnelshaped with a short tube and 5 unequal blunt lobes. Stigma capitate. Fr. a dry 2-celled caps., ventricose below, furrowed, opening transversely by a convex lid.
- [*LYCIUM*. Cal. small and adpressed to base of fruit. Cor. funnelshaped with a short tube and 5 equal patent lobes. Stam. exerted. Fr. a 2-celled berry.]
4. *DATURA*. Cor. funnelshaped, angular, 5-lobed. Cal. deciduous. Stigma 2-lobed. Caps. 4-valved, with 2 partially bipartite cells.

### Tribe I. *Solaneæ*.

#### 1. *SOLA'NUM* Linn. Nightshade.

1. *S. nigrum* (L.); st. herbaceous with tubercled angles, l. ovate bluntly dentate or wavy, fl. drooping.—St. l. 4.—Umbel from the intermediate spaces between the leaves. L. attenuate below. Fl. white. Fr.-stalks thickened upwards. Berries globular, black, or rarely green, or yellowish when ripe. St. a foot or more high.— $\alpha$ ; hairs incurved upwards, l. sinuate, berry black. *Sy. E. B.* 931.— $\beta$ . *S. miniatum* (Bernh.); angles of the st. with prominent tubercles, l. sinuate-dentate more deeply toothed, pubescence patent, berries scarlet. *Sy. E. B.* 932. It may be distinct.—Waste ground.  $\beta$ . Kent. Jersey. A. VII.—X. *Black Nightshade.* E. S. I.

2. *S. Dulcamara* (L.); st. shrubby, zigzag, l. cordate-ovate, upper l. hastate auricled, fl. drooping.—*E. B.* 565. St. 18. 3.—Corymb opposite to leaves. Fl. purple with 2 green spots at the base of each segment. Berries ovate, red. St. climbing to the height of 12—14 feet, nearly round, almost glabrous throughout, or (*S. littorale* Raab) st. and l. downy with patent hairs.— $\beta$ . *marinum* (Bab.); branches of the present year and l. fleshy and usually clothed with hairs incurved upwards, st. angular prostrate diffuse much branched, l. nearly all cordate not hastate. *S. lignosum seu Dulcamara marina* Ray 265.—Woods and hedges, common.  $\beta$ . Pebbly sea-beach. Sh. VI. VII. *Bittersweet.* E. S. I.

Tribe II. *Atropeæ*.2. AT'ROPA *Linn.* Deadly Nightshade. Dwale.

1. *A. Belladonna* (L.); st. herbaceous, l. broadly ovate-acuminate entire, fl. solitary axillary on short stalks.—*E. B.* 592.—St. 3 ft. high. Fl. lurid purple, drooping. Berry violet-black, highly poisonous.—Waste places, rare. P. VI.—VIII. E. S. I.

3. HYOSCY'AMUS *Linn.* Henbane.

1. *H. niger* (L.); l. oblong pinnatifid or sinuate sessile and clasping, lower l. stalked, fl. nearly sessile axillary unilateral. *Sy. E. B.* 936. St. 3. 4.—St. 1—2 feet high. Fl. lurid yellow with dark veins, drooping. Fr. erect. Whole herbage downy glandular, viscid, fetid. Fl. sometimes without dark veins.—Waste places, preferring a calcareous soil. B. V.—VII. E. S. I.

LYC'IUM *Linn.*

\*1. *L. barbarum* (L.); l. narrowly lanceolate narrowed at the base, cal. 2-lipped, cor.-tube as long as the limb, berry oblong.—*Sy. E. B.* 933.—A straggling shrub with long pendulous spinous branches. Fl. bluish. Filaments woolly at the base. Berry red.—An African plant which seems quite naturalized on north Norfolk coast and elsewhere. Sh. VI.—VIII. E.

4. DATU'RA *Linn.* Thorn-apple.

[1. *D. Stramonium* (L.); l. ovate unequally sinuate-dentate glabrous, caps. erect spinose.—*E. B.* 1288.—St. 1—2 ft. high. Fl. white (purplish in *D. Tatula*), large, erect. Fr. densely spinose. Caps. with 4 dissepiments below, only 2 at the top.—Waste ground, rare. A. VI. VII.] E.

## Order LVI. OROBANCHACEÆ.

Cal. variously divided, persistent. Cor. irregular, usually 2-lipped, persistent, imbricate in the bud. Stam. on the cor., 4, didynamous. Anth. 2-celled; cells distinct, parallel. Ovary in a fleshy disk, 1-celled, with 2 or more parietal placentas. Stigma 2-lobed. Fr. capsular, 2-valved, with many minute seeds.—Leafless root-parasites.

1. OROBANCHE. Cal. 4-cleft or of 2 usually bifid sepals. Cor. ringent, 4—5-cleft, deciduous, its base persistent. Bracts 1—3.
2. LATHRÆA. Cor. 2-lipped; the upper lip concave, deciduous, entire; its base persistent. Otherwise like *Orobanche*.

1. OROBAN'CHE Linn. Broom-rape.

\* *Sepals 2, entire or bifid, separate or connected below in front. Bract 1. Valves of caps. cohering at each end.*

1. *O. Rápum* (Thuill.); sep. 2-veined equally bifid nearly equaling the cor.-tube, cor. bellshaped ventricose at the base in front, its back curved, lips wavy obscurely denticulate (not fringed), upper lip concave nearly entire its sides patent, middle lobe of lower lip much longer than lateral lobes, *stam. inserted at the base of the cor.* glabrous below but their upper part and the style glandular-pubescent.—*R. I.* f. 900 & 923. *E. B.* 421. Fries states that *O. major* L. is certainly *O. elatior* Sutt.—*Stig. of 2 distant yellow lobes.* Anth. white when dry.—Parasitical upon Broom, Furze, and other shrubby leguminous plants. P. V.—VII. E. S. I.

2. *O. rúbra* (Sm.); sep. 1-veined lanceolate attenuate exceeding the cor.-tube, undivided, cor. bellshaped its back curved glandular-pubescent externally and the upper lip internally, lips acutely denticulate and crisped, upper lip emarginate its sides patent, lateral lobes of lower lip nearly equal, intermediate lobe rather longer, *stam. inserted near to the base of the corolla* slightly pilose within below but their top and the upper part of the style slightly glandular-pilose.—*Sy. E. B.* 1011. *R. I.* f. 885. Probably *O. Epithymum* DC.—Scarcely a foot high. Purplish red. *Stigma 2-lobed, pale pink.* Anth. fuscous when dry. Sep. with a second faint vein near their anterior margin; and in the dry plant there is an appearance of several more. L. few. Sweet-scented.—Parasitical upon *Thymus Serpyllum*. P. VI.—VIII. E. S. I.

3. *O. caryophyllácea* (Sm.); sep. many-veined lanceolate equally bifid falling short of the cor.-tube touching or connate in front, cor. tubular-bellshaped curved on the back, lips spreading, upper one 2-lobed its lobes porrect, lobes of lower lip nearly equal rounded wavy, *stam. inserted above the base of the cor. hairy within below* but their upper part and the style glandular-hairy.—*E. B. S.* 2639. *O. Galii* Duby.—Scarcely a foot high. *Stigma of 2 nearly separate dark purple lobes.* Anth. fuscous, yellow when dry. Sep. with crisp glandular hairs externally,

each lobe with 1 strong vein and several slender ones. Cor. similarly hairy on both sides.—On *Galium Mollugo* in Kent. P. VI. VII. E.

4. *O. elatior* (Sutt.); sep. many-veined equally bifid equalling the cor.-tube connate in front, cor. curved tubular slightly compressed above, upper lip 2-lobed toothed its lobes inflexed, lobes of lower lip 3 nearly equal acute toothed, stam. inserted above the base of the cor. glandular-hairy in the lower half within.—*E. B.* 568. *O. major* (L.) Fries.—Stem 2—3 ft. high. Stigma bilobed yellow. Upper lip of the cor. usually with an elevated point between the lobes. Cor. glandular externally. Stam. sometimes slightly hairy above. Anth. whitish when dry.—Parasitical upon *Centaurea Scabiosa*, rare. P. ? VI. VII. E.

5. *O. Picridis* (F. W. Schultz); sep. 1—3-veined entire or toothed below in front narrowed into 1 or 2 subulate points, cor. tubular-bellshaped its back nearly straight and compressed slightly curved at each end, lips denticulate wavy upper not notched its sides porrect, stam. inserted below the middle of the cor.-tube hairy in their lower half within, style slightly glandular-hairy below in front and above throughout, stigma bilobed.—*E. B. S.* 2956.—Height 6—18 inches. Stigm.-lobes just touching, purple. Anth. fuscous pale purple or yellowish.—Parasitical upon *Picris*. Cambridgeshire; Kent; Pemb.; Hunts.; Isle of Wight. P. ? VII. E.

6. *O. Hed'erae* (Duby); sep. 1-veined ovate below narrowed into 1 or 2 subulate points about equalling the cor.-tube, cor. tubular arcuate, lips denticulate wavy, upper one bilobed porrect (straight when dry), lobes of lower lip nearly equal the middle one longest, stam. inserted below the middle of the cor.-tube glabrous with a few scattered hairs on their lower part, style glabrous with a few hairs on the upper part, stigma scarcely bilobed.—*O. barbata* *E. B. S.* 2859, not Poir.—St. purplish, about a foot high. Lobes of stigma attached together by at least  $\frac{1}{3}$  of their circumference, yellow. Anth. fuscous, rather paler when dry. St. purplish.—Parasitical upon Ivy. P. VI. VII. E. I.

7. *O. minor* (Sutt.); sep. many-veined ovate below suddenly narrowed into 1 or 2 subulate points equalling or exceeding the cor.-tube, cor. tubular arcuate, lips bluntly denticulate wavy, upper lip porrect (inflexed when dry) notched, lobes of lower lip nearly equal, stam. inserted below the middle of the cor.-tube glabrous with a few scattered hairs below, style glabrous with a line of distant hairs on its anterior side, stigma bilobed.—*E. B.* 422.—Lobes of stig. only just touching at one point, purple. Anth. yellow when dry.—Parasitical chiefly upon *Trifolium pratense*. A. ? VI. VII. E.



8. *O. amethys'tea* (Thuill.); sep. many-veined ovate below narrowed into 1 or 2 subulate points, cor. tubular its *back curved immediately from the base otherwise straight*, lips unequally acutely denticulate wavy, upper lip hooded porrect notched, lobes of lower lip unequal the middle one larger, stam. inserted in the curvature of the cor. glabrous with many hairs at the base within, stigma bilobed.—*Sy. E. B.* 1017. *Atl. Fl. Par.* t. 19. E. *O. Eryngii* Duby.—*Lobes of stig. attached by  $\frac{1}{4}$ , purple.* Anth. dusky brown when dry. Perhaps a form of *O. minor*.—Parasitical upon *Daucus gummifer*. Whitsand Bay, Cornwall. Rock End, Torquay. *Mr. Townsend.* A. ? VI. E.

\*\* *Sepals 4 or 5, connected below. Bracts 3. Valves of capsule separating at the top.* PHELIPÆA Desf.

[*O. arenária* (Bork.); cal. of 5 sep. tubular with triangular-subulate teeth falling short of the cor.-tube, cor. tubular slightly curved in front, middle of the tube compressed on the back. throat slightly inflated externally glandular, *lobes of lips blunt with reflexed edges*, the lower lip hairy within, *suture of anth. hairy.*—*Atl. Fl. Par.* t. 19. L.—Height 12—18 inches. Stigma scarcely 2-lobed; style pale yellow (?), glandular. Filaments glabrous with a few hairs at their base. Lateral bracts linear-subulate, intermediate lanceolate attenuate above. St., scales and cal. glandular-pubescent.—Parasitical upon *Achillea Millefolium*? Alderney! (the Jersey plant is Sp. 9). P. VII. VIII.]

9. *O. cærúlea* (Vill.); cal. of 5 sep. tubular with lanceolate acute teeth falling short of the cor.-tube, cor. tubular curved in front, middle of tube compressed, *lobes of lips acute with reflexed edges*, lower lip hairy within, *anth. glabrous.*—*E. B.* 423. *Atl. Fl. Par.* t. 19. K.—About a foot high. Fl. bluish purple. Stig. scarcely 2-lobed, white. St., scales, bracts, cal. and cor. glandular-pubescent.—Grassy pastures. Herts. Norf. Isle of Wight. Chepstow. P. VI. VII. E.

[*O. ramósa* (L.); cal. of 4 sep. tubular with triangular ovate acuminate teeth, anth. glabrous, st. usually branched.—*E. B.* 184.—Sown with Hemp.]

## 2. LATHRÆA Linn. Toothwort.

1. *L. squamária* (L.); st. simple, fl. pendulous secund, lower lip of the cor. 3-cleft.—*E. B.* 50. *G. E. Smith S. Kent.* t. 3.—Rootstock fleshy, with thick scales. St. 3—8 in. high. Young raceme decurved. Bracts ovate or lanceolate. Style straight or curved. Upper lip nearly entire, or bifid.—Woods and thickets, parasitical upon Hazels &c. P. IV. V. E. S. I.

## Order LVII. SCROPHULARIACEÆ.

Cal. 4—5-cleft, persistent. Cor. irregular or 2-lipped or personate (subrotate in *Verbascum*), deciduous, imbricate in the bud. Stam. on the cor., usually 4, didynamous, or 2 or 5. Ovary free, 2-celled. Style simple; stigma 2-lobed. Fr. capsular, 2-celled; placentas attached to the dissepiment or ultimately central. Embryo straight, in axis of fleshy albumen.—No stipules.

\* *Stamens 5.*

1. VERBASCUM. Cal. of 5 sepals. Cor. rotate; segments unequal, spreading. Stam. unequal, 2 or more, hairy at the base.

\*\* *Stamens 4, didynamous.*

[ERINUS. Cal. in 5 deep segments. Cor. 5-parted with nearly equal emarginate segments and a short tube. Caps. 2-celled.]

2. DIGITALIS. Cal. in 5 deep segments. Cor. bellshaped, oblique, 4—5-fid. Caps. septicial, 2-celled.

3. ANTIRRHINUM. Cal. 5-parted. Cor. personate, gibbous at the base (*no distinct spur*); lower lip 3-fid; a prominent palate closing the mouth. Caps. opening by 2 or 3 pores at the top, 2-celled.

4. LINARIA. Cal. 5-parted. Cor. personate, spurred; lower lip 3-fid; a prominent palate closing the mouth. Caps. with valves or teeth at the top, 2-celled.

5. SCROPHULARIA. Cal. 5-lobed (in *S. vernalis* 5-cleft). Cor. globose; limb minute, of 2 short lips, upper 2-lobed often with a scale (the rudiment of a fifth stam.) within, lower 3-lobed. Caps. 2-valved, the edges inflexed, 2-celled.

6. LIMOSILLA. Cal. 5-cleft. Cor. 5-fid, bellshaped, equal. Caps. globose, 2-valved, 1-celled; placenta central, free or connected with a short dissepiment below.—L. radical.

7. MELAMPYRUM. Cal. tubular, 4-toothed. Cor. ringent; upper lip compressed laterally with reflexed edges; lower furrowed 3-fid. Caps. oblong, obliquely acuminate, compressed. *Seeds 1—2 in each cell, smooth.*

8. MIMULUS. Cal. prismatical, 5-toothed. Cor. ringent; upper lip folded back at the sides. *Seeds many.*

9. **PEDICULARIS.** *Cal. inflated, 5-toothed. Cor. ringent; upper lip compressed laterally; lower plane, 3-lobed. Caps. compressed, acute. Seeds many, angular.*
10. **RHINANTHUS.** *Cal. inflated, 4-toothed. Cor. ringent; upper lip compressed laterally; lower plane, 3-lobed. Caps. compressed, blunt. Seeds many, compressed, with an orbicular margin.—Alecterolophus (Hall).*
11. **BARTSIA.** *Cal. bellshaped, 4-fid. Cor. tubular, ringent; upper lip much arched, not compressed. Caps. pointed; cells many-seeded. Seeds compressed at the hile and with winged ribs at the back (large).*
12. **EUPRAGIA.** *Cal. tubular, 4-cleft. Cor. tubular, 2-lipped. Caps. pointed; cells many-seeded. Seeds slightly angular, very minute, crenate-ribbed; hile basal.*
13. **EUPHRASIA.** *Cal. tubular or bellshaped, 4-fid or 4-toothed. Cor. tubular, 2-lipped; lower lip with 3 notched or emarginate lobes. Anth. unequally pointed. Caps. blunt or emarginate; cells many-seeded. Seeds rather angular, longitudinally ribbed; hile subapical.*
14. **ODONTITES.** *Lower cor.-lip with 3 entire lobes. Anth. with 2 equal points; otherwise like Euphrasia.*
15. **SIBTHORPIA.** *Cal. in 5 deep spreading segments. Cor. rotate, irregularly 5-cleft. Caps. compressed, orbicular, 2-seeded, 2-valved.*

\*\*\* *Stamens 2.*

16. **VERONICA.** *Cal. 4—5-parted. Cor. rotate, unequally 4-lobed, lower lobe the smallest. Caps. compressed, 2-celled.*

1. **VERBAS' CUM** Linn.<sup>1</sup> Mullein.

\* *Leaves strongly decurrent. Raceme dense, nearly simple.*

1. *V. Thapsus* (L.); l. ovate-oblong crenate densely woolly on both sides all decurrent, st. simple, spike dense, pedicels shorter than the calyx, cor. rotate, segments oblong blunt filaments woolly 2 longer nearly glabrous, anth. all nearly equal.—*E. B.* 549. *V. Schraderi* Koch, *R.* xx. 1637.—St. 4—5 feet high. Cor. about twice as long as the calyx. Filaments with white

<sup>1</sup> Hybrids occur in this genus. Four are figured in *Sy. E. B.* t. 843, 844, 845, 846.

wool; the 2 glabrous ones about 4 times as long as their slightly decurrent anthers.—Waste ground. B. VII. VIII. *High-taper*. E. S. I.

\*\* *Leaves not decurrent. Racemes branched, paniced. Anth. all reniform, not decurrent.*

† Flowers yellow or whitish; hairs on the filaments white.

2. *V. Lychnitis* (L.); l. crenate nearly glabrous above woolly and powdery beneath, lower l. elliptic-oblong wedged shaped below scarcely stalked, upper l. sessile ovate-acuminate with a rounded base, st. angular paniced above with ascending branches, stam. equal, filaments all with white hairs.—*E. B.* 58. *R.* xx. 1648.—Height 2—3 feet. Fl. on short stalks, small, many, cream-coloured or rarely white.—Roadsides and waste places. B. VI.—VIII. *White Mullein*. E.

3. *V. pulverulentum* (Vill.); l. obscurely crenate clothed with mealy deciduous wool on both sides, lower l. oblong-elliptic attenuated into a stalk, upper l. sessile acuminate, st. terete paniced above with patent branches, stam. nearly equal scarlet with white hairs.—*E. B.* 487. *R.* xx. 1667.—Height about 3 feet. Fl. on very short stalks, which, as well as the calyx, are densely covered with wool, bright yellow. Cal.-teeth often glabrous.—Roadsides in Norfolk and Suffolk. B. VII. *Hoary Mullein*. E.

†† Flowers yellow; hairs on the filaments purple.

4. *V. nigrum* (L.); l. doubly crenate nearly glabrous above subpubescent beneath, lower l. cordate or ovate-oblong with long stalks, upper l. cordate-ovate nearly sessile, st. angular, raceme elongated, pedicels twice as long as the calyx, stam. equal with purple hairs.—*E. B.* 59. *R.* xx. 1649.—Fl. in clusters in a nearly simple long spike, small, bright yellow.—[*β. tomentosum* (Bab.); l. subpubescent above woolly beneath, fl. smaller.]—Banks and waysides. [*β. Alderney.*] P. VII. VIII. *Dark Mullein*. E. S.

\*\*\* *Leaves not, or very slightly decurrent.*

5. *V. Blattaria* (L.); l. crenate glabrous, lower l. ovate-oblong blunt sinuate, upper l. oblong or subcordate semialexicaul, pedicels solitary nearly twice as long as the bract, stam. and anth. unequal.—*E. B.* 393. *R.* xx. 1653.—Height 5—6 feet. Raceme glandular-pilose. Fl. cream-coloured. Filaments with purple hairs, the 2 longer hairy only on the inside.—On gravelly banks, rare. B. VIII. *Moth-Mullein*. E. \*I.

6. *V. virgatum* (With.); l. doubly serrate slightly glandular-hairy, lower l. oblong-lanceolate sublyrate lobate-crenate-serrate, upper l. oblong-acuminate semiamplexicaul, *pedicels* 1—5 together shorter than the bracts, *stam.* nearly equal.—*E. B.* 550. *R.* xx. 1655.—Usually much stouter than Sp. 5. Height 3—4 feet. Raceme glandular-pilose. Fl. yellow. Filaments with purple hairs, 2 rather longer and hairy only within.—On gravelly banks, rare. B. VIII. E. I.

[*Erinus alpinus* (L.); l. spatulate deeply serrate smoothish, ped. terminal subcorymbose.—Subcæspitose. Fl.-shoots 3—8 in. long, ascending.—In abundance on the old bed of the river near Tanfield, Yorkshire. P. VII.]

## 2. DIGITA'LIS Linn. Foxglove.

1. *D. purpurea* (L.); l. ovate-lanceolate crenate downy beneath, lower l. narrowed into footstalks, sep. ovate-oblong acute 3-veined downy, cor. blunt glabrous externally, upper lip scarcely cloven, segments of the lower lip ovate rounded.—*E. B.* 1297.—L. often crenate-dentate or -serrate. Fl. sometimes white or flesh-coloured. St. 3—4 feet high.—Hedgebanks and woods. P. VI.—VIII. E. S. I.

## 3. ANTIRRH'NUM Linn. Snapdragon.

\*1. *A. május* (L.); l. lanceolate opposite or alternate glabrous, fl. racemose, sep. ovate blunt much shorter than the cor., upper cor.-lip bifid.—*E. B.* 129.—Height 1—2 feet. Cor.  $1\frac{1}{2}$  in. long, purplish-red or white.—Old walls and calcareous cliffs. P. VII.—IX. *Great Snapdragon.* E.

2. *A. Oron'tium* (L.); l. linear-lanceolate opposite or alternate, fl. loosely spiked distant, sep. linear longer than the corolla.—*E. B.* 1155. St. 27.—About a foot high. Fl. purple.—Dry sandy and gravelly fields. A. VII. VIII. E. I.

## 4. LINA'RIA Mill. Toadflax.

\* *Stems trailing. Fl. axillary.*

\*1. *L. Cymbalaria* (Mill.); l. roundish heartshaped 5-lobed glabrous, st. procumbent.—*E. B.* 502.—Stems slender, rooting. Fl. solitary, axillary, upon long stalks, pale blue.—Old walls. P. V.—X. *Ivy-leaved Toadflax.* E. S. I.

2. *L. Elat'ine* (Mill.); l. ovate-hastate, lower l. ovate, cor.-spur straight, peduncles glabrous.—*E. B.* 692.—Fl. solitary, axillary,

upon long slender stalks, small, yellow, with the upper lip purple.—Corn-fields. A. VII.—IX. *Fluellin.* E. I.

3. *L. spúria* (Mill.); l. roundish-ovate, spur curved upwards, peduncles hairy.—*E. B.* 691.—Fl. similar to the last but larger. L. with here and there a small tooth.—In this species and the two preceding some of the fl. are often regular with 5 spurs or partially so with 2, 3, or 4.—Gravelly and sandy corn-fields. A. VII.—IX. E.

\*\* *Stems erect or rarely diffuse.* † Fl. solitary.

4. *L. minor* (Desf.); l. linear-lanceolate blunt glandular-pubescent mostly alternate, fl. axillary, peduncles 3 times as long as the calyx, segments of upper cor.-lip diverging, seeds oblong sulcate.—*E. B.* 2014.—Fl. small; the tube, upper lip, and spur of the cor. purplish; lower lip yellowish. St. erect, 4—10 in. high, branched, glandular-pubescent. Lower l. nearly spatulate.—Sandy and gravelly fields. A. VI.—VIII. E. S. I.

†† Fl. racemose.

[*L. Pelisseriána* (Mill.); glabrous, l. linear, the lower ternate or quaternate, upper alternate, sterile branches radical prostrate with ternate lanceolate or ovate l., fl. racemose, peduncles as long as the bracts, sepals linear acute twice as long as the capsule, seeds nearly flat with a fringed wing one side smooth the other tubercular.—*E. B. S.* 2832.—Fl. purple with darker veins. St. one or more from each root, erect, about a foot high. Caps. bilobed.—Jersey. A. V.]

\*5. *L. supina* (Desf.); glabrous, rachis, ped. and sep. glandular-hairy, l. linear blunt mostly whorled, sep. linear-spathulate shorter than the caps. or spurs, seeds smooth nearly flat with a striate wing.—*Sy. E. B.* 858. *L. maritima* DC. *Icon. Gall.* 12.—St. diffuse or ascending. Fl. capitata-racemose, yellow; throat and spur with slender purple lines. Style entire.—Plymouth and Poole; a ballast plant. Perhaps a native at Hayle, and St. Blazey's Bay, Cornwall. A. VII. VIII. E.

\*6. *L. purpúrea* (Mill.); glabrous, l. linear-lanceolate scattered, lower l. irregularly in fours, fl. narrowly racemose, sep. linear shorter than the caps. and long incurved spur, seeds angular with a network of elevated lines.—*Sy. E. B.* 960.—Fl. purple or yellow with the lips purple; spur two or three times as long as the ped. which is usually shorter than the bract. St. erect, leafy.—Old walls. Naturalized. P. VII. VIII. E.

7. *L. répens* (Ait.); glabrous, l. linear scattered or partly whorled, fl. racemose, *sep. lanceolate* as long as the spur but shorter than the caps., *seeds angular with transverse elevated lines*.—*E. B.* 1253. *L. striata* DC.—Fl. white with blue veins. St. erect, branched, leafy, 1—1½ foot high, slender. Seeds much smaller than those of *L. vulgaris*.—*L. italica* and *L. sepium* may be hybrids between this and *L. vulgaris*.—Calcareous soils, particularly near the sea, rare. P. VII.—IX. E. I.

8. *L. vulgáris* (Mill.); glabrous, rachis and peduncles glandular-hairy, l. linear-lanceolate scattered crowded, fl. racemose imbricate, *sep. ovate acute glabrous* shorter than the caps. or spur, seeds tubercular-asperous with a smooth orbicular margin.—*E. B.* 658.—Fl. large, yellow, rarely milk-white with an orange palate. St. erect, 2 feet high, as well as the l. glabrous. Common and partial flower-stalks occasionally glabrous.—The state called *Peloria* with 5 spurs and an equal and regular cor. is rarely found. *E. B.* 260.— $\beta$ . *latifolia* (Bab.); l. narrowly lanceolate, fl.-l. often lanceolate very glaucous, fl. twice as large in a few-fl. lax raceme, ped. glabrous, spur directed perpendicularly downwards. *Sy. E. B.* 964. *L. speciosa* Ten. ?—Hedges on a gravelly soil. P. VI. VII. *Yellow Toadflax*. E. S. I.

### 5. SCROPHULA'RIA Linn. Figwort.

\* *Cal. of 5 rounded lobes. Cor. purplish; upper lip with a scale (staminode) on its inner side.*

1. *S. nodósa* (L.); l. ovate acute subcordate glabrous *doubly and acutely serrate, lower teeth largest*, st. acutely 4-angular, cymes lax, *sep. roundish-ovate with a narrow membranous margin, staminode wedgeshaped slightly emarginate*.—*E. B.* 1544.—Root tuberous, thick, knotty. St. 2—3 feet high. Bracts small, lanceolate, acute. Fl. greenish purple, lurid, sometimes green or pale. Caps. ovate.—Moist hedges and thickets. P. VI. VII. *Knotted Figwort*. E. S. I.

2. *S. umbrósa* (Dum.)<sup>1</sup>; l. ovate-lanceolate acute subcordate glabrous sharply *serrate, lower teeth smaller*, st. and petioles winged, *cymes lax few- (4—8-) flowered*, *sep. roundish with a broad membranous margin, staminode bilobed with diverging lobes*.—S. Ehrharti, C. A. Stev. *E. B. S.* 2875. Not *S. alata* Gill.—St. tall. *Bracts leaflike*, lanceolate, acute. Fl. dark purple. Caps. subglobose, blunt.—Wet places. P. VIII. IX. E. S. I.

3. *S. aquat'ica* (L. !); l. cordate-oblong roundly blunt glabrous *crenate-serrate*, st. and petioles winged, *cymes dense corym-*

<sup>1</sup> See Du Mortier in *Bull. Belg.* vii. 36.

*bose many- (1—15-) flowered*, sep. roundish blunt with a broad membranous margin, *staminode roundish-reniform*.—*E. B.* 854. *S. Balbisii Horn., Koch.*—St. 2—5 feet high. *Bracts linear, blunt*. Fl. dark purple, occasionally milk-white. Caps. ovoid, pointed.—Specimens from Cambr. and Heref. seem to be the *S. cinerea*, Dum.; their staminode is rounded at the end, not emarginate.—In wet places. P. VII. VIII. E. S. I.

4. *S. Scorodonia* (L.); *l. cordate-triangular with large double teeth downy on both sides*, st. bluntly quadrangular downy, cymes lax few-flowered, sep. roundish downy with a membranous margin, staminode roundish entire.—*E. B.* 2209.—St. 2—3 feet high. L. wrinkled. *Bracts leaflike*, lower exactly like the leaves. Fl. purple. Caps. ovoid, acute.—In moist places. West of Cornwall. Tralee, Kerry. Jersey. P. VII. E. I.

\*\* *Cal. of 5 deep acute segments. Cor. yellow; no staminode.*

†5. *S. vernalis* (L.); *l. downy cordate-acute doubly serrate*, st. winged hairy, *cymes axillary corymbose with leaflike bracts*, sep. oblong with a recurved apex.—*E. B.* 567.—St. about 2 feet high. *Fl. yellow, inflated*; the mouth much contracted. Caps. ovoid, acute. Differing greatly from the other species and allied in appearance to some of the *Calceolarie*.—Waste places, rare. P. IV. V. E. S. ?

#### 6. LIMOSSEL'LA Linn. Mudwort.

1. *L. aquatica* (L.); *l. lanceolate-spathulate on long stalks*, ped. axillary crowded shorter than the petioles.—*E. B.* 357. *St.* 30. 15.—Very small; st. 0, except the naked stoles. Fl. small, white or rose-coloured. Caps. minute, ovoid.—Muddy places where water has stagnated. A. VII.—IX. E. S.

#### 7. MELAMPY'RUM Linn. Cow-wheat.

1. *M. cristatum* (L.); *spikes densely imbricate 4-sided, bracts heartshaped acuminate pectinate-dentate lower ones with a long leaflike recurved point*.—*E. B.* 41.—Bracts rose-coloured at the base. Fl. yellow, tinged with purple. L. linear-lanceolate, acute, entire, with netted veins beneath. St. 8—12 in. high.—Woods and thickets in the Eastern Counties. A. VII. *Crested Cow-wheat*. E.

2. *M. arven'se* (L.); *spikes lax conical, bracts ovate-lanceolate-attenuate pinnatifid with subulate segments and with a few large glandular points beneath*, cal. hispid equalling the cor-tube with long lanceolate-attenuate teeth from an ovate base,



cor. closed.—*E. B.* 53.—Bracts purple-rose-colour. Fl. variegated with yellow, rose-colour, and purple. L. linear-lanceolate, acute, rough-edged, slightly downy on both sides, entire. St. 8—18 in. high.—Corn-fields and dry banks in the Eastern Counties and the Isle of Wight. A. VII. *Purple Cow-wheat.*  
E.

3. *M. pratensis* (L.); *fl. axillary secund in distant pairs*, upper bracts lanceolate with 1 or 2 teeth at the base, *cor.* 4 times as long as the glabrous calyx *closed, lower lip projecting.*—*E. B.* 113.—Teeth and tube of the cal. about equal in length. L. lanceolate, or ovate-lanceolate, or with a cordate base, entire, varying greatly in size. Fl. large, pale yellow, horizontally patent. St. 6—12 in. high.— $\beta$ . *ericetorum* (D. Oliv.); hispid, l. lanceolate or linear-lanceolate, bracts toothed, fl. near together, *cor.*-tube whitish.— $\gamma$ . *M. montanum* (Johnst.); smaller in all its parts, bracts quite entire, l. linear-lanceolate hispid, the 2 lower obovate-lanceolate blunt.—Woods and thickets.  $\beta$ . West of Ireland.  $\gamma$ . Mountains. A. VI.—VIII. E. S. I.

4. *M. sylvaticum* (L.); *fl. axillary secund in distant pairs*, bracts mostly all entire linear-lanceolate, *cor.* about twice as long as the glabrous calyx, *open lips equal* in length.—*E. B.* 804.—Teeth of the cal. longer than the tube. L. linear-lanceolate, entire. Fl. very small, deep yellow, erect. St. 12 in. high.—Alpine woods. A. VII. E. S. I.

### 8. MIM'ULUS Linn.

\*1. *M. hiteus* (L.); l. roundish ovate veined, lower ones stalked, uppermost l. clasping, st. creeping, fl. yellow.—*Sy. E. B.* 967.—An American plant, naturalized in many boggy places. P. VI.—IX. E. S. I.

### 9. PEDICULA'RIS Linn.

1. *P. palustris* (L.); *st. solitary erect branched throughout*, l. pinnatifid, segments oblong blunt lobed, *cal. ovoid pubescent, 2-lobed*, lobes incise-dentate crisped.—*E. B.* 399.—Upper lip of the cor. with a short truncate beak with a triangular tooth on each side. Fl. large, crimson. St. 12—18 in. high, angular, with alternate branches. Whorl of ovate-acute scales at root-crown.—Marshy and boggy places. A.<sup>p</sup> V.—VII. *Lousewort.*  
E. S. I.

2. *P. sylvatica* (L.); *st. branched at the base erect, branches long spreading prostrate*, l. pinnatifid, segments ovate lobed, *cal.*

*oblong glabrous* irregularly 5-lobed, upper lobe lanceolate, other lobes with 3 leaflike divisions.—*E. B.* 400.—Upper lip of the cor. as in the last. Fl. large, rose-colour. Summit of the ped. with a loose membranous cuticle enclosing the base of the calyx. Primary st. erect, often very short; branches prostrate. Whorl of ovate-lanceolate crenate undivided reflexed leaves at root-crown.—Wet heathy and rather hilly pastures. P. ? V.—VIII. *Red Rattle.* E. S. I.

### 10. RHINAN'THUS Linn. Yellow Rattle.

1. *R. Crista-galli* (L.); l. oblong-lanceolate serrate, fl. in lax spikes, cal. glabrous, lobes of the upper cor.-lip short roundish, bracts ovate incise-serrate, seeds with a membranous border.—*E. B.* 657. *R. minor* (Ehrh.).—Lateral lobes of the upper cor.-lip very blunt, shorter than broad, bluish. Bracts green throughout. Style downy near the top. Caps. as long as broad. Cor.-tube straight. St. 1—2 feet high, nearly simple.—Meadows and pastures. A. VI. E. S. I.

2. *R. major* (Ehrh.); l. linear-lanceolate serrate, fl. in crowded spikes, cal. glabrous, lobes of the upper cor.-lip oblong, bracts with an attenuate point incise serrate.—Cal. often slightly downy on its edges. Lateral lobes of the upper cor.-lip longer than broad, purple; the central part truncate. Cor.-tube slightly curved. Style glabrous. Caps. often longer than broad. Anth. very villose. Bracts yellowish with green points.—*a. platypterus* (Fries); seed not twice as broad as its wing. *R. major* Koch, *R. I.* f. 975.—*β. stenopterus* (Fries); seed quite twice as broad as its wing. *R. major* *E. B. S.* 2737.—*γ. apterus* (Fries); seed not winged but rounded and longitudinally ribbed or furrowed on the back. *R. Reichenbachii* Drej.—Cultivated land. *a.* Hastings. *β.* North of England and Scotland. *γ.* Arbroath and Monifieth, Forfarshire. A. VII. VIII. E. S.

### 11. BART'SIA Linn. Red Eye-bright.

1. *B. alpina* (L.); l. opposite ovate slightly clasping bluntly serrate.—*E. B.* 361.—Creeping. St. square, 4—8 in. high, simple. Fl. forming a short dense leafy spike, purplish blue, downy. Cal. purplish, viscid. Anth. hairy.—Alpine pastures, rare. P. VI. VII. E. S.

### 12. EUFRAG'IA Griseb. Marsh Eye-bright.

1. *E. viscosa* (Benth.); l. opposite, upper l. alternate ovate-

lanceolate sessile acutely serrate.—*Bartsia* L. *E. B.* 1045.—L. sometimes linear-lanceolate. St. round, 3—12 in. high, simple. Root fibrous. Fl. distant, axillary, upper ones crowded, yellow. Anth. hairy. St., l., and cal. viscid.—Damp places in the West of E., South-west of S., and South of I. A. VII.—IX.

E. S. I.

### 13. EUPHRA'SIA Linn. Eye-bright.

1. *E. officinális* (L.); l. ovate or oblong-lanceolate nearly sessile serrate (3—5 teeth on each side), lobes of the lower cor.-lip emarginate, of the upper lip patent sinuate-dentate, anth. hairy, seeds with ribs.—*E. B.* 1416.—St. 1—8 in. high. Fl. axillary, solitary, sessile, crowded towards the ends of the branches.—*a*; glandular-pubescent above and on the calyx, caps. oblong-obovate, seeds ovoid greyish. L. usually large and broad, sometimes densely imbricate (*E. ericetorum* Jord. ?).—*β*. *E. nemorosa* (Pers.) pubescent not glandular, caps. linear-oblong, seeds fusiform yellowish. L. usually narrow, sometimes (*E. Salisburgensis* Funk. ?) with very long teeth.—Some authors divide this into many species; but even the above are scarcely distinguishable at all times.—Pastures, woods, heaths. A. VI.—VIII. Common Eye-bright. E. S. I.

### 14. ODONTITES, Duby.

1. *O. rúbra* (Pers.); l. narrowed from near the base opposite linear-lanceolate-attenuate remotely serrate, floral l. usually longer than the fl. with 2—4 teeth and an entire end, cal.-segments as long as their tube lanceolate acute, cor. pubescent open, lobes of the lower lip oblong, style protruded even before the fl. opens, caps. oblong.—*Bartsia Odontites* Huds. *E. B.* 1415.—St. about a foot high, much branched. Fl. many, pink, in leafy unilateral spikes. Filaments, anth., and stam. hairy.—*a*. *O. verna* (R.); branches ascending straightish, l. rounded below, cal.-teeth narrowed below.—*β*. *O. serotina* (R.); branches spreading and curving up, l. narrowed below, cal.-teeth narrowed below.—*Euphrasia rotundata* (Ball), *A. N. H.* ser. 2. iv. 30, has not been again found. I have seen no specimen.—Corn-fields and waste places. A. VII. VIII. E. S. I.

### 15. SIBTHORP'IA Linn.

1. *S. europæ'a* (L.).—*E. B.* 649.—A slender trailing plant with filiform creeping stems, and alternate long-stalked roundish

reniform leaves with a few large crenatures. Fl. very small, axillary, solitary, on short stalks, pinkish, inconspicuous.—Damp shady places in the South and South-west. P. VI.—IX.  
E. I.

16. VERON'ICA Linn. Speedwell.

*Racemes axillary. (Root perennial.)*

1. *V. scutellata* (L.); l. linear-lanceolate acute sessile minutely denticulate, racemes alternate, fruitstalks slender reflexed, caps. of 2 flattish orbicular lobes, st. erect.—*E. B.* 782.—Stoloniferous. St. weak, 1 ft. high, glabrous or hairy. Fl. pale flesh-coloured, with darker lines. Sep. small, lanceolate, acute, shorter than the capsule.—Boggy places. P. VI.—VIII. E. S. I.

2. *V. Anagal'lis* (L.); l. lanceolate serrate acute sessile, racemes opposite, fruitstalks spreading, caps. oval slightly notched, st. erect.—*E. B.* 781.—Stoloniferous. St. glabrous, thick, succulent, hollow, 12—24 in. high. Fl. pale blue. Sep. lanceolate. Racemes sometimes glandular. Whole plant usually glabrous.—In stagnant water. P. VI.—VIII. *Water Speedwell.*  
E. S. I.

3. *V. Beccabun'ga* (L.); l. stalked oval crenate-serrate, racemes opposite, fruitstalks spreading, caps. roundish tumid slightly notched, st. procumbent at the base rooting.—*E. B.* 655.—Glabrous, succulent. Fl. bright blue, rarely pink or flesh-coloured.—Ditches and streams. P. V.—VIII. *Brooklime.* E. S. I.

4. *V. Chamæ'drys* (L.); l. nearly sessile cordate-ovate incise-serrate, racemes opposite, fruitstalks ascending, caps. flat obcordate deeply notched ciliate shorter than the cal., st. bifariously hairy ascending.—*E. B.* 623.—St. about a foot long. Fl. large, many, handsome, blue. Sep. lanceolate, acute. L. on autumnal shoots slightly stalked.—Hedgebanks. P. V. VI. *Germander Speedwell.*  
E. S. I.

5. *V. montana* (L.); l. stalked broadly ovate serrate, fruitstalks ascending, caps. orbicular notched their margins crenulate and ciliate longer than the cal., st. diffuse hairy all round procumbent.—*E. B.* 766.—St. often above a foot long. Racemes lax. Fl. few, pale blue. Caps. very large, quite flat. Sep. ovate-lanceolate, acute.—Woods and thickets in hilly districts. P. V. VI. E. S. I.

6. *V. officinális* (L.); l. shortly stalked oval serrate, racemes spike-like many-flowered, fruitstalks erect, caps. obcordate trun-

cate bluntly notched longer than the cal., st. prostrate creeping hairy.—*E. B.* 765.—St. 6—12 in. long. Racemes erect, much longer than the leaves.—Caps. not always notched. St., l., and calyx sometimes smooth.— $\beta$ . *V. hirsuta* (Hopk.); l. ovate-lanceolate, caps. abrupt undivided. *E. B. S.* 2673. Much smaller than the true *V. officinalis*. I have not seen wild specimens.—Dry banks and heaths.  $\beta$ . Dry heath, Carrick, Ayrshire. P. VI.—VIII. *Common Speedwell.* E. S. I.

\*\* *Racemes terminal; cor.-tube longer than broad.*

7. *V. spicdta* (L.); l. ovate or lanceolate crenate-serrate entire at the end, lower l. blunt stalked, *raceme spikelike long dense*, bracts longer than the pedicels, caps. ovate emarginate with a very long style.—*E. B.* 2.—St. erect, branching at the base about 6 in. high, or in *V. hybrida* taller. Fl. blue. Lower l. oval with a wedgeshaped base, or ovate with a rounded or slightly cordate base (*V. hybrida* L. *E. B.* 673).—Rare. On chalky heaths near Newmarket and Bury; and ( $\beta$ .) on limestone cliffs. P. VII. VIII. *Spiked Speedwell.* E.

\*\*\* *Racemes terminal; cor.-tube very short.* † Seeds flat.

8. *V. saxat'ilis* (L.); l. oval serrate at about the middle, lower l. smaller, *raceme pubescent with crisped hairs not glandular* few-flowered, caps. ovate-attenuate, valves bifid.—*E. B.* 1027.—Raceme persistently subcorymbose. Ped. long, erect. Fl. large, bright blue. St. much branched, decumbent, woody.—*V. suffruticosa* Sm. is probably a mistake.—Exposed alpine rocks. P. VII. S.

9. *V. alpina* (L.); l. elliptic or ovate dentate or entire, lower l. smaller, *raceme hairy with patent hairs not glandular* few-flowered, caps. oval-obovate emarginate crowned with the very short persistent style.—*Sy. E. B.* 980.—St. decumbent, scarcely rooting, erect, simple except quite at the base, 4—6 in. high. Fl.-raceme corymbose; fr.-raceme dense.—Summits of Highland mountains. P. VII. VIII. S.

10. *V. serpyllifolia* (L.); l. ovate or oval slightly crenate, lower l. smaller and rounder, *raceme long many-flowered*, caps. obcordate broader than long crowned with the long persistent style.—*E. B.* 1075.—Fl. whitish with blue veins. St. rooting below, afterwards erect, 2—6 in. high.— $\beta$ . *V. humifusa* (Dicks.); st. quite prostrate, racemes shorter. *Sy. E. B.* 979.—Roadsides and damp places.  $\beta$ . Mountains. P. V.—VII. E. S. I.

[\**V. peregrina* (L.); l. all obtuse and narrowed below, lower l. obovate-oblong, uppermost l. bractlike exceeding the minute fl., raceme slightly spiked many-flowered lax, ped. very short, caps. obcordate, style very short.—*Sy. E. B.* 977.—St. erect, 2—5 in. high. Fl. white faintly tinged with pink. Caps. smooth, broader than long; lobes rounded. Sep. linear-lanceolate.—Fields. A. V.] S. I.

[\**V. repens* (DC.); l. all roundly oval, pan. short few-flowered, ped. exceeding l., cor. 2—3 times as long as the sep., caps. obovate shorter than its style.—*V. tenella* R. xx. 1718.—St. prostrate, rooting; no erect fl.-shoot and raceme. Ped., cal., and caps. glandular-hairy.—A Corsican pl. naturalized at Manchester and York. P. IV. V.]

11. *V. arvensis* (L.); l. cordate-ovate crenate, lower l. stalked, uppermost l. lanceolate entire bractlike exceeding the flowers, raceme slightly spiked many-flowered lax, ped. very short, caps. obcordate broader than long compressed ciliate on the keel.—*E. B.* 734.—St. ascending, 2—6 in. or rarely a foot long. Fl. pale blue. Caps. smooth, with rounded lobes which exceed the style. Seeds 12—14. Sep. lanceolate, unequal.—Gravelly and sandy places. A. IV.—VII. E. S. I.

12. *V. ver'na* (L.); l. pinnatifid, lower l. stalked ovate serrate, upper l. lanceolate entire bractlike, raceme slightly spiked many-flowered lax, ped. very short, caps. obcordate compressed ciliate on the keel.—*E. B.* 25.—St. erect. 1—3 in. high, simple or branched in the lower part. Caps. smooth or downy, with rounded lobes. Style very short. Seeds 12—14. Sep. linear-lanceolate, unequal.—Sandy heaths. Bury, Thetford, and Mildenhall, Suff. A. V. E.

†† Seeds concave.

13. *V. triphyl'los* (L.); l. fingered, lower l. ovate entire or dentate stalked, raceme slightly spiked lax few-flowered, ped. exceeding the calyx, caps. obcordate compressed smooth ciliate on the keel.—*E. B.* 26.—St. erect with spreading branches, 4—5 in. high. Fl. deep blue. Ped. usually longer than the leaves. Sep. oval. Known by its spreading st., deeply fingered l., and dark-blue flowers.—Sandy fields. Bury, Mildenhall, Brandon, &c., Suff. York. A. IV. E.

\*\*\*\* Flowers axillary, solitary. Seeds concave. St. prostrate.

14. *V. agrest'is* (L.); l. all stalked cordate-ovate incise-serrate, sep. oval, stam. inserted at the very bottom of the cor., caps. of 2 turgid keeled lobes, seeds about 6 in a cell.—*E. B. S.* 2603.

*R. I. f. 440.*—Caps. hairy all over, or only ciliate on the keel; hairs all straight and glandular. L. usually exceeding the peduncles. Lower cor.-lip white.—[*V. opaca* (Fries), with spatulate sep., fewer seeds, hairs on caps. incurved short with a few longer and glandular, stam. inserted in the throat of cor., is probably British.]—Fields and waste places. A. IV.—IX. *Green Field Speedwell.* E. S. I.

15. *V. polita* (Fries); l. all stalked cordate-ovate incise-serrate, sep. broadly ovate acute, stam. inserted at the very bottom of the cor., caps. of 2 turgid lobes, seeds 8—12 in a cell.—*E. B. 783. R. I. f. 404, 405.*—Caps. with short dense glandless hairs and other shorter glandular ones, rarely glabrous. L. usually falling short of the peduncles. Fl. wholly blue.—Cor. sometimes as large as that of *V. Buxbaumii*.—Fields and waste places. A. IV.—IX. *Gray Field Speedwell.* E. S. I.

\*16. *V. Buxbaum'ii* (Ten.); l. all stalked cordate-ovate incise-serrate, sep. lanceolate-acute, stam. from the bottom of the cor., caps. of 2 divaricate lobes compressed upwards and sharply keeled, seeds about 8 in a cell.—*E. B. S. 2769.*—St. long, hairy. L. falling short of the peduncles. Fl. twice the size of those of the preceding, as large as those of *V. Chamædryis*, blue.—Fields. A. IV.—IX. E. S. I.

17. *V. hederifolia* (L.); l. cordate with 5—7 large toothlike lobes all stalked, sep. cordate-attenuate ciliate, caps. of 2 turgid lobes, seeds 2 in each cell.—*E. B. 784.*—Fl. pale blue.—Fields and banks. A. IV.—VI. *Ivy-leaved Speedwell.* E. S. I.

[*Acan'thus mollis*, belonging to the Order *Acanthaceæ*, has been introduced by unknown agency to St. Agnes Isle, Scilly, and Traeth Manaccan, Cornw.]

### Order LVIII. LABIATÆ.

Cal. tubular, regular or 2-lipped, persistent. Cor. 2-lipped, upper lip entire or bifid, lower 3-fid, or nearly regular. Stam. 4, didynamous, rarely 2. Ovary free, 4-lobed. Style 1, from the base of and between the lobes; stigma bifid. Fr. in 1—4 small nutlets (not true nuts although so called).—Fl. unsymmetrical, often forming 2 cymes so placed as to resemble a whorl (a verticillaster). L. opposite.—No stipules.

Tribe I. *MENTHOIDEÆ.* Cor. bellshaped, nearly regular. Stam. distant, straight, diverging upwards.

1. *MENTHA.* Cor. 4-fid, tube very short. Cal. regular, 5-toothed. Stam. 4; anth.-cells parallel.

2. LYCOPUS. Cor. 4-fid, scarcely longer than the regular 5-toothed calyx. *Stam.* 2; anth.-cells parallel or ultimately diverging; 2 upper *stam.* imperfect.

Tr. II. *MONARDEÆ*. Cor. 2-lipped. *Stam.* 2, perfect, parallel under the upper lip of the corolla.

3. SALVIA. *Filaments with 2 diverging branches*; only one bearing a perfect anth.-cell. Cor. ringent. Cal. tubular, 2-lipped.

Tr. III. *SATUREINEÆ*. Cor. 2-lipped. *Stam.* 4, distant; cells of anth. separate, diverging; connective dilated.

4. ORIGANUM. *Stam.* diverging; connective subtriangular. Upper lip of cor. straight, nearly flat; lower patent, 3-fid. *Cal. with 5 equal teeth* and 10—13 veins; throat hairy. *Spikes 4-sided*, resembling catkins with imbricate bracts.

5. THYMUS. *Tips of stam. patent*. Anth.-cells at first nearly parallel, afterwards diverging; connective subtriangular. Upper lip. of cor. straight, nearly flat; lower patent, 3-fid. *Cal. 2-lipped* and 10—13-veined; throat hairy. Fl. whorled, axillary or spiked.

6. CALAMINTHA. *Tips of stam. converging* under the upper lip of the corolla. Anth.-cells at length diverging; connective subtriangular. Upper lip of cor. straight, nearly flat; lower patent, 3-fid. Cal. 2-lipped.

[*MELISSINEÆ*. Cor. 2-lipped. *Stam.* distant; anth.-cells connected above.]

[*MELISSA*. *Tips of stam. converging* under the upper lip of the cor. Anth.-cells diverging. *Upper lip of cor. concave*; lower patent, 3-fid. Cal. 2-lipped; upper lip flat, with 3 teeth, the lateral teeth folded at their midrib.]

Tr. IV. *SCUTELLARIEÆ*. *Stam.* approaching, parallel under the upper lip of the cor. Cal. 2-lipped, closed in fruit.

7. SCUTELLARIA. *Tips of stam. incurved*. *Filaments simple*. Anth. of the two longer and *inferior stam.* 1-celled, of the shorter and superior 2-celled. Cor. 2-lipped, upper lip concave. Cal. ultimately closed and compressed; lips entire, upper one with a concave scale on its back.

8. PRUNELLA. Two inferior *stam.* longest. *Filaments bifid*, one branch barren. *Anth. all 2-celled*. Cor. ringent; upper



lip concave, entire. Cal. ultimately closed and compressed; upper lip flat, truncate, 3-toothed; lower bifid.

Tr. V. *NEPETEÆ*. Stam. approaching, parallel under the upper lip of the cor., 2 *inferior shortest*. Cal. tubular.

9. *NEPETA*. Anth.-cells diverging. Cor. ringent; upper lip flat, straight, emarginate or bifid. Cal. 5-toothed.

Tr. VI. *STACHYDEÆ*. Stam. approaching, parallel under the upper lip of the cor., 2 *inferior longest*. Cal. tubular or bellshaped, spreading in fruit.

\* *Stamens exceeding the tube of the corolla.*

10. *MELITTIS*. Anth. approaching in pairs and forming a cross, bursting longitudinally. *Upper lip of the cor. flat*, entire, straight; lower lip with 3 rounded nearly equal lobes. Cal. membranous, bellshaped, ample, variously lobed.

11. *LAMIUM*. Anth. approaching in pairs; cells diverging, bursting longitudinally. *Upper lip of the cor. arched*; lateral lobes of the lower lip minute toothlike or rarely long. Cal. bellshaped, 5-toothed; teeth nearly equal.—*Galeobdolon* (Huds.) has the lobes of the lower lip of the cor. nearly equal and acute.

12. *LEONURUS*. Anth. approaching; cells nearly parallel, bursting longitudinally. *Upper lip of cor. nearly flat*; lower with 3 blunt lobes. Cal. tubular, 5-toothed, 2 lower teeth rather the longest. Nutlets flatly truncate.

13. *GALEOPSIS*. Anth. approaching in pairs; cells opposite bursting by 2 valves transversely. Upper lip of cor. arched; lower lip 3-lobed with 2 teeth on its upper side, lobes unequal. Cal. tubular, 5-toothed; teeth equal or 2 upper ones longest. Nutlets rounded at the end.

14. *STACHYS*. Anth. approaching in pairs; cells diverging, bursting longitudinally. Upper lip of cor. concave; lower of 3 unequal lobes. Cal. tubular-bellshaped with 5 equal teeth. Nutlets blunt and convex at the end.

15. *BALLOTA*. Anth. approaching in pairs; cells diverging, bursting longitudinally. Upper lip of cor. erect, concave, lower 3-lobed, middle lobe cordate. Cal. funnelshaped with 5 equal teeth. Nutlets convex and rounded at the end.

\*\* *Stamens falling short of the tube of the corolla.*

16. MARRUBIUM. Anther-cells diverging, bursting longitudinally. Upper lip. of cor. straight, erect, flattish, cloven; lower 3-lobed, middle lobe the largest. Cal. tubular, teeth nearly equal or 2 longer. *Nutlets flatly truncate.*

Tr. VII. AJUGOIDEÆ. Cor. with the upper lip very short, or deeply bifid and appearing as if wanting.

17. TEUCRIUM. Stam. parallel, protruded between the lobes of the upper lip of the cor., inferior longest; cells bursting longitudinally. *Cor. with the upper lip deeply bifid, lobes long*; lower lip 3-lobed. Cal. tubular, 5-toothed; the teeth equal or the upper one larger (2-lipped).

18. AJUGA. Stam. parallel, protruded far beyond the upper lip of the cor., inferior longest; cells bursting longitudinally. *Cor. with the upper lip very short, 2-lobed*; lower 3-lobed, much longer than the upper. Cal. ovate-bellshaped, nearly equally 5-cleft.

### Tribe I. *Menthoideæ.*

#### 1. MEN'THA Linn. Mint.<sup>1</sup>

\* *Throat of the calyx naked. General inflorescence determinate. Whorls of fl. in terminal spikes or clusters.*—† Stoles aerial, leafy.

†1. *M. viridis* (L.); l. *glabrous, sessile lanceolate acute serrate, spikes lax cylindrical, bracts subulate, cor. glabrous.*—*E. B.* 2424. *Sole Menth* 5. Whorls of the spike rather distant. L. glandular beneath.—β. *crispa* (Benth.); l. deeply cut and crisped.—In marshy places, rare. β. Glen Farg, Perthshire. *Hooker. P. VIII. Spear-Mint.* E. S.

2. *M. rotundifolia* (L.); l. *sessile roundly ovate crenate-serrate wrinkled shaggy beneath, spikes linear cylindrical dense, bracts lanceolate, "fr.-cal. not contracted at the mouth," cor. hairy.*—*E. B.* 446. *Sole* 3.—Viscid. St. 1—2 ft. high. L. usually with a cordate base. Whorls of spike nearly all close together. Scent acrid.—Waste places, rare. *P. VIII. IX. Round-leaved Mint.* E. S. I.

<sup>1</sup> See Mr. Baker's valuable paper in *J. of B.* iii. 233.

## †† Stoles subterranean.

3. *M. alopecuroïdes* (Hull); *l.* subsessile roundly cordate-oval serrate wrinkled hairy beneath, spikes conical-cylindrical, bracts lanceolate, cor. hairy.—*Sy. E. B.* 1021. *M. rotundifolia* Sole 4. *M. velutina* Bab. *M. dulcissima* Dum.—*L.* not felted beneath.—West of S. and east of E. P. VIII. IX. E. S.

4. *M. sylvestris* (L.); *l.* subsessile ovate or lanceolate serrate silky beneath, spikes linear-cylindrical dense, bracts subulate, "fr.-cal. contracted at the mouth," cor. hairy.—*E. B.* 686. Sole 1 & 2.—*L.* lanceolate or oblong, more or less hairy, not shaggy. Whorls of spike nearly all close together. Scent sweet.—Damp waste ground. P. VIII. IX. *Horse-Mint.* E. S.

\*5. *M. piperita* (Sm.); *l.* stalked ovate-lanceolate or oblong serrate, upper *l.* smaller, bracts lanceolate, spikes lax short blunt interrupted below, cal. tubular glabrous below with lanceolate subulate teeth.—*E. B.* 687.—*St.* 1—2 ft. high and *l.* nearly glabrous or hairy on the veins beneath. Cal. glandular.—Wet places, rare. P. VII. VIII. *Pepper-Mint.* E. S. I.

6. *M. aquatica* (L.); *l.* stalked ovate-acute serrate rounded or subcordate below hairy on both sides, fl.-*l.* falling short of fl., fl.-whorls in few axillary and terminal subglobose or ovoid clusters, cal.-teeth triangular  $\frac{1}{2}$  as long as tube.—*M. hirsuta* *E. B.* 447. (*M. citrata* *E. B.* 1025 is a glabrous form.)—*St.* 12—18 in. high. Cal. glandular.—Moist places. P. VII. VIII. *Capitate Mint.* E. S. I.

7. ?*M. hirsuta* (L., Fr.); *l.* stalked ovate or ovate-lanceolate serrate hairy above woolly beneath, fl.-*l.* falling short of the fl., whorls of fl. in subcylindrical thick dense spikes interrupted below, cal.-teeth subulate  $\frac{2}{3}$  length of tube.—*M. pubescens* *Sy. E. B.* 1026, 1027.—*St.* 12—18 in. high. I know little of this. Is it distinct from *M. aquatica*? E.

\*\* *Throat of cal. naked. General inflorescence indeterminate. Fl. in axillary distant whorls; none amongst the uppermost l., or shorter than them.*

8. *M. sativa* (L.); *l.* nearly sessile ovate or ovate-lanceolate sharply serrate, upper *l.* smaller but similar and exceeding the fl., whorls distant dense, cal. tubular or bellshaped, cal.-teeth triangular-lanceolate.—*St.* 1—2 ft. high. Uppermost leaves often above the flowers.—*a. M. sativa* (L.); *l.* hairy on both sides, ped. cal. and cor. hairy. *E. B.* 448.—*β. M. rubra*

(Huds.); l. stalked nearly glabrous, ped. lower part of cal. and cor. glabrous, cal.-teeth hairy, veins of l. purple. *M. gracilis* (Sm.) is a slender green form with subsessile leaves. *M. cardiaca* (Baker) differs by having the upper l. sessile and st.-l. nearly glabrous.— $\gamma$ . *M. pratensis* (Sole); l. rather blunt much veined hairy above, glabrous (except on the veins) beneath, ped. and cal. glabrous, cal.-teeth ciliate.— $\delta$ . *M. gentilis* (L.); l. acute with few veins slightly hairy on both sides, upper l. similar, ped. lower part of cal. and cor. glabrous. *E. B.* 2118.—These forms seem to vary into each other.—Wet places. P. VII. VIII. E. S. I.

9. *M. arvensis* (L.); l. stalked ovate bluntly serrate, upper l. similar and equally large, whorls distant, cal. bellshaped, teeth triangular as broad as long.— $\alpha$ . *vulgaris*; l. narrowed below. *M. arvensis* Sm. *E. B.* 2119.— $\beta$ . *M. agrestis* (Sm.); l. roundish subcordate below, upper ones nearly sessile. *E. B.* 2120.—L. very variable in form, from nearly round and blunt to ovate-acute.—Corn-fields. P. VII.—IX. *Corn Mint.* E. S. I.

\*\*\* *Throat of the calyx closed with hairs.* PULEGIUM Opitz.

10. *M. Pulégium* (L.); l. stalked ovate slightly crenate all similar, whorls all distant globose many-flowered, cal. tubular hispid closed with hairs in the throat.—*E. B.* 1026. *Sole* 23.—St. prostrate. L. often recurved, uppermost axils without flowers. The smallest of our species and remarkably different in habit.—Wet places. P. VIII. IX. *Penny-royal.* E. S. I.

## 2. LYC'OPUS Linn. Gipsywort.

1. *L. europæus* (L.); l. stalked ovate-oblong sinuate-dentate or pinnatifid, sterile stam. wanting, nuts about equalling the tube of the calyx.—*E. B.* 1105.—Subglabrous or pubescent. L. opposite. Fl. small, in dense whorls.—Banks of streams and ditches. P. VII. VIII. E. S. I.

## Tribe II. *Monardeæ.*

### 3. SAL'VIA Linn. Sage.

1. *S. Verbenáca* (L.); l. oblong blunt cordate below sinuate and crenate or dentate stalked, upper l. short broad cordate sessile clasping, bracts cordate acuminate, tube of the cor. equalling the calyx.—*E. B.* 154.—Varies with the l. incise-dentate. Remarkable for its enlarged very broad sessile upper leaves. St.

1—2 ft. high. Cor. purple, small; upper lip concave, laterally compressed, straight except at the tip. Upper cal.-lip broad; teeth small, converging.—Dry gravelly banks. P. V. VI. *English Clary*. E. S. I.

2. ?*S. clandestina* (L.); l. oblong cordate below sinuate-dentate or incise-dentate stalked, *upper l. oblong acute* sessile scarcely cordate or clasping, bracts cordate acuminate, *tube of the cor. exceeding the calyx*.—St. a foot high. Cor. purple, small; upper lip very broad, teeth very small.—Probably distinct from the preceding, but very difficult to characterize. Syme's plant (*Sy. E. B. 1057*) seems different.—Dry gravelly banks, rare. Lizard, Cornwall. Channel Isles. P. VII. E.

3. *S. pratensis* (L.); l. oblong-ovate cordate below, crenate-dentate stalked, upper l. small, sessile lanceolate acute, bracts cordate acuminate, cor. thrice as long as the calyx.—*E. B. 153*.—St. 1—2 ft. high. Known by its large flowers.—Cobham. Kent. Middleton Stoney, Oxfordshire. P. VII. E.

### Tribe III. *Satureineæ*.

#### 4. ORIGANUM *Linn.* Marjoram.

1. *O. vulgare* (L.); l. stalked broadly ovate, bracts ovate exceeding the cal., heads of fl. ovoid or oblong paniced crowded.—*E. B. 1143*.—St. a foot high. Bracts usually purple. L. often slightly toothed. Fl. purple.—Spikes sometimes prismatic and oblong. *O. megastachyum* Link, *Sy. E. B. 1046*.—Dry uncultivated places. P. VIII. E. S. I.

#### 5. THYMUS *Linn.* Thyme.

1. *T. Serpyllium* (L.); *st. prostrate creeping*, l. all oblong or lanceolate narrowed into the flat fringed stalks, *flowering shoots ascending*, fl. capitate, upper cal.-lip with 3 short triangular teeth, lower of 2 subulate teeth, *upper cor.-lip oblong*.—*E. B. 1514*.—Forming a cushion with a fringe of prostrate barren shoots which in the next year produce erect fl.-shoots from their lower joinings and are prolonged at the end. L. narrowed below, their lower half and the stalk often fringed, rather conspicuously veined beneath, often narrow. Nutlets globose, mealy. Cor. purple; upper lip conspicuously notched.—Dry heaths. P. VI.—VIII. E. S. I.

2. *T. Chamæ'drys* (Fr.); stems alike diffuse ascending, l. all broadly oblong with flat fringed stalks, fl. whorled or capitate, upper cal.-lip with 3 triangular teeth, lower of 2 subulate teeth, upper cor.-lip semicircular.—*E. B. S.* 2992.—Forming a tuft of flowering and barren shoots; the st. of preceding year ending in a fl.-shoot. L. less narrowed into the stalks than in *Sp. 1.*, usually only the stalk is fringed, less prominently veined beneath, broad. Nutlets roundish, subcompressed, with a basal apiculus. Cor. purple; upper lip slightly and obscurely notched. In the large forms the st. is stronger; it is not so in *Sp. 1.*—Heaths. P. VI.—VIII. E. S.

## 6. CALAMIN'THA Moench. Calamint.

\* *Fl. in whorls of 2 forked cymes.*

1. *C. Nep'eta* (Clairv.); l. ovate serrate pale beneath shortly stalked, cal. rather bellshaped obscurely 2-lipped its teeth shortly ciliate all nearly of the same shape the upper ones slightly shorter and broader, cyme many-flowered its common stalk about as long as the primary partial stalk.—*Thymus* L. *E. B.* 1414.—St. usually many from the crown of the root, 12—18 in. high; branches short, erect. Fl. purple. Hairs in the throat of the cal. protruded.—Dry banks, rare. P. VII. VIII. Lesser Calamint. E.

2. *C. officinalis* (Moench?) ; l. broadly ovate slightly serrate green on both sides on longish stalks, cal. tubular distinctly 2-lipped its teeth with long cilia those of the upper lip triangular and ascending, of the lower twice as long and subulate, lower lip of the cor. with distant segments the middle one longest, cyme scarcely forked few-flowered, fl. at an angle with the pedicel.—*Thymus Calamintha* L. *E. B.* 1676. *C. menthæfolia* G. & G.—St. usually solitary, or few from the root, 1—2 ft. high; branches long, ascending. Fl. purplish. Hairs in the throat of the cal. included. Larger in all its parts than the preceding.—*a. C. menthifolia* (Host); ped. of cyme  $\frac{1}{2}$  as long as the primary partial stalk.—*β. C. ascendens* (Jord.); ped. of lower cymes as long as or longer than the primary partial stalk. *C. off. β. Briggsii* (Syme). *Sy. E. B.* 1051.—Dry banks, rare. *β. Devon.* P. VII.—IX. Common Calamint. E. I.

3. *C. sylvat'ica* (Bromf.); l. broadly ovate sharply serrate, cal. tubular distinctly 2-lipped, teeth with long cilia those of the upper lip patent or reflexed, lower lip of the cor. with overlapping segments all nearly equally long, cyme many-flowered its common stalk about as long as the primary partial stalk.—*E. B. S.* 2897.

*C. officinalis* G. & G.—Creeping slightly.—St. about 2 ft. high, nearly simple. L. large. Cal.-teeth tinged with purple; hairs in the throat included. Cor. purplish, very large, tube much protruded, middle lobe of lower lip short and broad.—Isle of Wight. Near Torquay. (*J. of B.* xi. 208.) P. VIII.—X. E.

\*\* *Whorls of 6 simple separate peduncles.* ACINOS Moench.

4. *C. A'cinos* (Clairv.); l. ovate subserrate acute with revolute margins, cal. tubular gibbous below distinctly 2-lipped, upper lip with short triangular teeth lower with subulate teeth all converging in fruit.—*Thymus* L. *E. B.* 411.—St. 6—8 in. long. Fl. blue.—Dry gravelly places and limestone rocks. A. VII. VIII. *Basil.* E. S. I.

\*\*\* *Fl. in dense branched axillary clusters. The many setaceous bracts forming a kind of involucre.* CLINOPODIUM Linn.

5. *C. Clinopodium* (Benth.); l. ovate rounded below slightly crenate-serrate, clusters equal many-flowered.—*Clinopodium vulgare* L. *E. B.* 1401.—St. 1—1½ foot high. Fl. purple in 2 or 3 dense whorls, the uppermost terminal.—Dry bushy places. P. VII. VIII. *Wild Basil.* E. S. I.

### [*Melissineæ.*]

#### [*MELIS'SA, Linn. Balm.*]

[*M. officinalis* (L.); l. ovate crenate-serrate acute paler beneath, cal. rather bell-shaped slightly ventricose in front distinctly 2-lipped, upper lip flat truncate with three short broad teeth, lower with 2 lanceolate teeth.—*Sy. E. B.* 1053.—St. 2 feet high. Fl. in axillary second whorls, white.—Escaped in the South. P. VII. VIII.] E.

#### Tribe IV. *Scutellaricæ.*

##### 7. SCUTELLA'RIA Linn. Skull-cap.

1. *S. galericulata* (L.); l. shortly stalked all oblong-lanceolate cordate below crenate-serrate, fl. axillary opposite secund, calyx without glands.—*E. B.* 523.—Cor. large blue. St. 6—12 in. high, stout.—This genus has a long support (carpophore) to its outlets.—Banks of rivers and ditches. P. VII. VIII. E. S. I.

2. *S. minor* (L.); l. shortly stalked, lower broadly ovate, intermediate ovate-lanceolate with the base cordate, upper l. lanceolate with a rounded base, fl. axillary opposite secund, cal. pubescent.—*E. B.* 524.—Cor. small, pale pink. St. 4—8 in. high, slender.—Moist heaths and boggy places. P. VII.—IX. E. S. I.

8. PRUNEL'LA *Linn.* Self-heal.

1. *P. vulgaris* (L.); 1. stalked oblong-ovate blunt, upper lip of the cal. with short truncate mucronate teeth, lower lip with ovate-lanceolate mucronate teeth.—*E. B.* 961.—About 8 in. high. L. nearly entire or slightly toothed; on Continental specimens sometimes pinnatifid. Fl. blue, rarely white, whorled, crowded into a dense spike, with 2 broad kidneyshaped acuminate bracts under each whorl. Cal. reddish purple.—In damp pastures. P. VII. VIII. E. S. I.

Tribe V. *Nepeteæ*.9. NEP'ETA *Linn.*

1. *N. Catária* (L.); 1. stalked cordate acute incise-serrate whitish-pubescent beneath, cymes dense many-flowered spiked, nutlets smooth and glabrous—*E. B.* 137.—Fl. white. St. 2—3 feet high, downy or mealy. Stam. at length curved outwards.—Waste places. P. VII. VIII. *Cat-Mint*. E. S. I.

2. *N. Glechóma* (Benth.); 1. cordate-reniform crenate, whorls axillary stalked secund 3—4-flowered, cal.-teeth ovate-acuminate awned, nutlets oblong with impressed dots.—*Glechoma hederacea* L. *E. B.* 853.—St. procumbent, creeping. Fl. blue-purple; length of cor.-tube and its hairiness are variable. Anth. in pairs forming a cross.—Hedges and thickets. P. IV.—VI. *Ground-Ivy*. E. S. I.

Tribe VI. *Stachydeæ*.10. MELIT'TIS *Linn.* Bastard Balm.

1. *M. Melissophyl'lum* (L.).—*E. B.* 577.—L. oblong-ovate or slightly cordate. Upper lip of the cal. with 2 or 3 teeth. Fl. purple with a white margin or variegated in different ways, large. St. 1—2 ft. high.—*M. grandiflora* (Sm. *E. B.* 636) is only a slight variety.—Woods in the South. P. V. VI. E.

11. LA'MIUM *Linn.* Dead-Nettle.

\* Lower lip of cor. with one large obcordate lobe with 1—2 teeth on each side of its base.

1. *L. amplexicaúle* (L.); 1. roundish-cordate bluntly incise-crenate, lower l. stalked, upper sessile clasping, cal.-teeth longer than the tube (green) at length connivent, cor.-tube straight naked within—*E. B.* 770.—Lower fl.-whorls usually distant.



Fl. purple-red. Nutlets small, smooth, three times as long as broad, with a small triangular oblique terminal space. Cor.-tube much exceeding the calyx, slender. The cor. does not always expand, but the anth. are fertile and the fr. is produced.—Sandy and chalky fields. A. V.—VIII. *Henbit*. E. S. I.

2. *L. intermedium* (Fries); l. incise-crenate, lower l. stalked, upper reniform-cordate sessile, cal.-teeth longer than the tube hispid always spreading, straight cor.-tube with a faint hairy ring within.—*E. B. S.* 2914.—Lower fl.-whorls usually distant. Nutlets large thrice as long as broad, with a large triangular terminal rather oblique space. Tube of the cor. equal, cylindrical. Cal.-teeth usually purple, rigid.—Common in S. rare in E. and I. A. VI.—IX. E. S. I.

3. *L. incisum* (Willd.); l. cordate incise-dentate all stalked, upper broadly ovate crowded, cal.-teeth as long as or longer than the tube always spreading, tube of the cor. straight naked within.—*E. B.* 1933.—Uppermost l. wedgeshaped below. Fl.-whorls usually all contiguous. Fl. reddish. Cor.-tube equal, cylindrical, sometimes with a faint ring of hairs within. Nutlets smooth.—Cultivated and waste ground. A. IV.—VI. E. S. I.

4. *L. purpureum* (L.); l. cordate crenate-serrate all stalked, upper ovate crowded, cal.-teeth as long as the tube always spreading, cor.-tube a little curved below with a ring of hairs within.—*E. B.* 769.—Fl.-whorls contiguous. Cor. pale purple, lip spotted with red; tube narrowed below. Nutlets about twice as long as broad, smooth. A form with more deeply-cut leaves is often taken for *L. incisum*.—Waste and cultivated ground. A. IV.—VIII. *Red Dead-Nettle*. E. S. I.

5. *L. album* (L.); l. cordate-ovate acuminate deeply serrate stalked, cal.-teeth as long as the tube, cor.-tube exceeding the calyx with an oblique ring of hairs within and narrowed below the ring.—*E. B.* 768.—St. 12—18 in. high. Fl. large, white.—Waste ground. P. V. VI. *White Dead-Nettle*. E. S. I.

[*L. maculatum* (L.); l. cordate-ovate acuminate deeply serrate stalked, cal.-teeth longer than the tube, cor.-tube exceeding the calyx with a transverse ring of hairs within and narrowed below the ring.—St. 12—18 in. high. Fl. purple. L. marked with white, cordate-ovate (*L. maculatum*); or green, triangular-cordate (*L. lævigatum* *E. B.* 2550).—An escape. Fifeshire. Clova. P. VI.—VIII.] E. ? S.

\*\* Lower lip of cor. in 3 entire nearly equal acute lobes.

GALEOB'DOLON *Huds.* Archangel.

6. *L. Galeob'dolon* (Crantz); l. ovate acuminate truncate be-

low coarsely serrate stalked, upper l. lanceolate attenuate below, helmet of the cor. long entire, *lower lip in 3 entire nearly equal lobes*.—*G. luteum* Huds. *E. B.* 787. *G. montanum* Reich. !—Fl. yellow. St. 12—18 in. high. Lower l. coarsely and even doubly serrate.—[*G. luteum* Reich. ! has the l. all ovate-acuminate and the lower ones simply crenate with a minute apiculus.]—Woods and thickets. P. V. VI. E. S. I.

### 12. LEONU'RUS Linn. Motherwort.

†1. *L. Cardiaca* (L.); lower l. palmately 5-fid incise-dentate, upper ones 3-lobed entire wedgeshaped below, cor.-tube with an oblique ring of hairs within, helmet nearly flat, lip spreading its middle lobe entire.—*E. B.* 286.—St. 3 feet high. Cor. hairy externally, purple. Fl. in crowded whorls. Cal.-teeth sharp.—Hedges and waste places, rare. P. VIII. E. S.

### 13. GALEOP'SIS Linn. Hemp-Nettle.

1. *G. dubia* (Leers); st. not thickened at joinings with deflexed hairs, l. ovate-lanceolate serrate soft and downy on both sides, upper l. ovate, cal. glandular shaggy, *upper cor.-lip deeply notched*.—*G. villosa* Huds. *E. B.* 2353. *G. ochroleuca* Lam.—Cor. large, pale yellow. St. 10—12 in. high, with gland-tipped hairs on its upper part.—Sandy corn-fields, rare. A. VII. VIII. E.

2. *G. Lad'anum* (L.); st. not thickened at joinings soft with deflexed hairs, l. ovate-lanceolate lanceolate or lanceolate-attenuate at both ends serrate or nearly entire downy on both sides, cal. shaggy with adpressed hairs and a few gland-tipped hairs intermixed, *upper cor.-lip slightly notched*.—*E. B.* 884.—Cor. purple variegated with crimson and white, shaggy externally. St. about a foot high. St. and l. varying much in hairiness, pale green or purplish.—β. *G. intermedia* (Vill.); l. broad not narrowed below, regularly toothed, whorls of fl. all separate.—*Sy. E. B.* 1075.—γ. *G. canescens* (Schultz); l. narrow nearly entire, bracts often reflexed at the end, cal. and upper part of st. clothed with patent hairs.—In gravelly and sandy districts. β. Moray, S. A. VIII. IX. E. S. I.

3. *G. Tet'rahit* (L.); st. thickened at joinings hispid, l. oblong-ovate acuminate serrate, cal. tubular, cal.-teeth and tube nearly equal, *cor.-tube equalling the cal.*, upper lip ovate.—*E. B.* 207.—St. 1—2 ft. high. Cor. purplish variegated with white, large; tube slender slightly inflated; middle lobe of lower lip subquadrate, flat, crenulate, blunt or slightly emarginate. Cal.-teeth as long as their tube, which is shorter but quite as broad

and more strongly ribbed than in the next species, rather inflated below the mouth. Nutlets wholly green, the oblique top longitudinally marked with veins all springing from the wholly acute inner angle of the nut. L. slightly pubescent above.—*β. G. bifida* (Boenn.); middle lobe of lower cor.-lip oblong emarginate purple with pale ultimately revolute edges, base with a yellow 2-lobed spot and several dots. A more slender plant with paler leaves and smaller flowers. *Sy. E. B.* 1079.—Woods and cultivated ground. A. VII.—IX. E. S. I.

4. *G. versicolor* (Curt.); st. thickened at the joinings hispid, l. oblong-ovate acuminate serrate, cal. bellshaped, cal.-teeth shorter than the tube, *cor.-tube much exceeding the cal.*, upper lip roundish-oval.—*E. B.* 667.—St. 2—3 ft. high. Cor.-tube inflated above. Fl. very large, yellow, usually with a broad purple spot upon the lower lip. Cal.-teeth shorter than their tube. Nutlets with the oblique top dark brown, inner angle rounded off almost to the base. Difficult to distinguish upon paper from *G. tetrahit*.—Cultivated ground. A. VII. VIII. E. S. I.

#### 14. STA'CHYS Linn. Woundwort.

1. *S. Betonica* (Benth.); whorls in oblong interrupted term. spike many-flowered, st. erect, lower l. ovate-oblong with a cordate base crenate blunt with long stalks, upper l. oblong-lanceolate serrate acute subsessile, bracts linear-lanceolate equaling the *nearly glabrous cal.*, *stam. falling short of the lip.*—*Betonica officinalis* L. *E. B.* 1142.—St. 1—2 ft. high. Whorls sometimes considerably separated. Cor. purplish red; tube exerted. The English plant has the round crenate not emarginate lower lip of *B. hirta* (R.).—Woods and thickets. P. VII. VIII. *Betony*. E. S. I.

2. *S. germanica* (L.); whorls many-flowered, *st. erect woolly*, l. oblong-ovate or ovate-lanceolate with a cordate base crenate-serrate stalked *densely silky*, *upper l. lanceolate acute sessile, cal. silky*, teeth acute mucronate spinous, bracts equaling the calyx.—*E. B.* 829.—St. 2—3 ft. high. Fl. purple.—Chalky soil, Oxfordshire. B. VII. *Downy Woundwort*. E.

3. *S. sylvatica* (L.); whorls 6—8-flowered, st. erect, l. *cordate-ovate* serrate long-stalked, *floral l. linear entire*, cal.-teeth lanceolate very acute, *bracts minute.*—*E. B.* 416.—Cal.-teeth rather spinous. Petioles and l. nearly equal. Fl. reddish-purple. Nutlets opaque, punctured and irregularly tubercled. L. clothed with scattered adpressed hairs or densely silky on both sides.—Woods and thickets. P. VII. VIII. *Hedge-Woundwort*. E. S. I.

4. *S. palustris* (L.); whorls 6—10-flowered, st. erect, *l. linear-or ovate-lanceolate subcordate below acute crenate-serrate nearly sessile*, cal.-teeth lanceolate very acute, bracts minute.—*E. B.* 1675.—St. 1½—2 ft. high. Cal.-teeth rather spinous. Lower l. with very short stalks, uppermost sessile. Fl. dull purple. Nutlets shining, very minutely dotted.—β. *S. ambigua* (Sm.); l. stalked ovate-lanceolate cordate below serrate. Petioles sometimes half as long as the leaves.—River-sides and damp places. P. VII. VIII. *Marsh-Woundwort.* E. S. I.

5. *S. arvensis* (L.); whorls 4—6-flowered, st. decumbent or ascending, *l. ovate-cordate blunt crenate stalked, floral l. ovate-oblong sessile acute*, cal.-teeth lanceolate awned, cor. scarcely exceeding the cal., bracts minute.—*E. B.* 1154.—Fl. pale purple. Nutlets covered with minute dots and scattered tubercles.—Corn-fields. A. VIII. IX. E. S. I.

[*S. annua* (L.); whorls 4—6-flowered, st. erect, lower l. ovate-oblong blunt crenate-serrate stalked, *floral l. lanceolate acute*, cal.-teeth lanceolate very acute, tube of the cor. exceeding cal., bracts minute.—*E. B. S.* 2669.—Fl. yellowish. Nutlets minutely rough.—Gadshill and Sevenoaks, Kent. A. VIII. IX.] E.

#### 15. BALLO'TA Linn. Horehound.

1. *B. foetida* (Lam.); l. crenate-serrate, bracts linear-subulate, cal. funnelshaped, *cal.-teeth broadly ovate short patent or reflexed*.—*E. B.* 46. *R. I. f.* 1041!—St. 2—3 ft. high. Lower l. cordate; upper ovate. Fl. purple or white. Scent pungent. A hard coarse plant. Whole plant including the cor. covered with hairs.—Waste places. P. VII. VIII. E. S. I.

2. *B. ruderális* (Sw.); l. crenate-serrate, bracts linear-subulate, cal. funnelshaped, *cal.-teeth ovate gradually acuminate erect*.—*R. I. f.* 1039!—St. 2—3 ft. high. Lower l. cordate; upper ovate. Fl. purple or white. Scent agreeable. *Very hairy and soft*.—I still with Grenier consider this distinct from Sp. 1.—Waste places, rare. Abundant at Llanwarne, Herefordshire. P. VII. VIII. E.

#### 16. MARRUBIUM Linn. White Horehound.

1. *M. vulgäre* (L.); st. erect hoary, l. ovate narrowed into a petiole or roundish cordate crenate hoary rough, whorls many-flowered, cal.-teeth 10 subulate patent hooked woolly below, their upper half glabrous.—*E. B.* 410.—St. 1—2 ft. high, with many whorls of small whitish flowers. L. sometimes dentate, rarely roundly cordate.—Waste places, rare. P. VIII. IX. E. S. I.

Tribe VII. *Ajugoideæ*.17. TEUCRIUM *Linn.*

1. *T. Scorodonia* (L.); st. erect, l. oblong-ovate their base cordate crenate-serrate green on both sides, racemes lateral and terminal one-sided, floral l. ovate acute rather longer than the pedicels, *upper cal.-lip ovate*, lower 4-toothed, cor.-tube exserted.—*E. B.* 1543.—St. 1—2 ft. high. L. stalked, with glandular resinous mealiness beneath, wrinkled; sometimes oblong, truncate below or subcordate, coarsely and unequally dentate. Fl. yellowish.—Woods and dry stony places. P. VII. VIII. *Wood-Sage*. E. S. I.

2. *T. Scor'dium* (L.); st. procumbent below, l. *sessile* oblong dentate green on both sides, *floral l. similar*, whorls 2—6-flowered axillary distant, *cal.-teeth short equal*.—*E. B.* 828.—More or less hairy or woolly. St. 1—2 ft. long. L. attenuate or broad or even cordate below. Fl. purple.—Wet places rare. P. VII. VIII. E. I.

\*3. *T. Chamæ'drys* (L.); st. ascending, l. ovate incise-crenate wedgedshaped and entire below green on both sides, *floral l. similar smaller nearly entire*, whorls *racemose* 5-flowered, *cal.-teeth lanceolate nearly equal*.—*E. B.* 680.—St. much branched, 6—8 in. long, lower part woody. Fl. purplish. Lower floral l. exactly like the stem-l., upper l. gradually smaller and broader below.—Ruined walls, rare. P. VII. E. [S. I. ?]

†4. *T. Bótrys* (L.); l. *trifid or pinnatifid* green on both sides, *segm. oblong entire or cut*, floral l. similar, whorls axillary 4—6-flowered, *cal. gibbous at base inflated-tubular*, *cal.-teeth lanceolate equal*.—*E. B. S.* 2946.—St. erect, about 8 in. high; branches ascending. Fl. many, pale purple.—Box Hill and Selsdon, Surrey. A. VIII. IX. E.

18. AJUGA *Linn.* Bugle.

1. *A. reptans* (L.); fl. whorled, st. solitary with long stoles, l. ovate or obovate entire or crenulate stalked, stem-l. sessile.—*E. B.* 489.—St. 6—8 in. high. Lower whorls distant, upper ones spiked. Cor.-tube with a ring of hairs within. Fl. blue, rarely white.—Wet places. P. V. VI. *Common Bugle*. E. S. I.

[*A. alpina*, *E. B.* 477, seems to be *A. genevensis*. It is not stoloniferous, and its upper st.-l. fall short of the flowers.]

2. *A. pyramidális* (L.); fl. whorled most or all of the whorls spiked, st. solitary, l. ovate-oblong entire or crenulate, radical l.

attenuate below, stem-l. sessile *upper l. exceeding the fl.*—*E. B.* 1270.—St. about 6 in. high. L. gradually decreasing upwards. Cor.-tube with a ring of hairs within. Fl. bluish purple. Plant hairy. It has subterranean offsets and short autumnal stoles.—Highland pastures, very rare, S. South Isles of Arran, I. P. V. VI. S. I.

3. *A. Chamæpitys* (Schreb.); *fl. solitary axillary*, st. much branched spreading, *l. deeply trifid with linear entire segments*, floral l. similar exceeding the flowers.—*E. B.* 77.—Hairy. Lowest l. much broader, toothed rather than 3-lobed. Fl. yellow with dark spots. St. reddish purple, branched, 3—6 in. high.—Sandy and chalky fields. A. V.—VII. *Ground-Pine.* E.

### Order LIX. VERBENACEÆ.

Cal. tubular, persistent. Cor. irregular, tubular. Stam. didynamous, or 2. Ovary 2—4-celled; style 1, terminal; stigma bifid. Fr. a caps. or berry, with 2—4 nutlets more or less cohering.—No stipules.

1. VERBENA. Cal. 5-fid. Cor. irregular, 5-lobed, slightly 2-lipped. Stam. included, 4, didynamous, or 2. Capsule dividing into 4 nutlets.

#### 1. VERBENA *Linn.* Vervain.

1. *V. officinalis* (L.); st. erect solitary 4-angular, l. ovate oblong trifid or lacinate-multifid rough, spikes filiform somewhat panicled, *stam.* 4.—*E. B.* 767.—St. rather hispid, 1—2 ft. high. L. lobed and serrate, opposite. Spikes long, slender. Fl. small, distant, pale purple.—Waste ground. P. VII. VIII. E. I.

### Order LX. LENTIBULARIACEÆ.

Cal. permanent, inferior, divided. Cor. irregular, 2-lipped spurred. Stam. 2. Ovary free, 1-celled of 2 carpels. Stigma of 2 plates, one smaller or inconspicuous. Caps. 1-celled; placenta large, free, central. Seeds many.—No stipules.

1. PINGUICULA. Cal. 2-lipped, lower bifid, upper of 3 segments. Cor. ringent, spurred. Stam. at base of corolla.
2. UTRICULARIA. Cal. 2-leaved, lower often notched, upper entire. Cor. personate, spurred. Stam. at base of upper cor.-lip.

## 1. PINGUIC'ULA Linn. Butterwort.

1. *P. vulgaris* (L.); spur subulate shorter than the *cor.-segments* which are very unequal *oblong rounded separated* entire.—*E. B.* 70.—L. all radical, fleshy, covered with minute crystal-line points, pale green; when the plant is gathered they curve backwards so as to hide the root. Fl. violet. Caps. ovoid, acute.—Bogs. P. V. VI. E. S. I.

2. *P. grandiflora* (Lam.); spur subulate cylindro-conical often notched as long as *cor.-segments* which are very unequally *broadly obovate rounded contiguous*.—*E. B.* 2184.—Much larger than *P. vulgaris*. Fl. very large, violet. Caps. ovoid, rounded at the end. Length of spur variable.—Bogs. Kerry and Cork. P. V. VI. I.

3. *P. alpina* (L.); *spur conical shorter than unequal border of cor.* and curved towards lower lip, caps. acute, scape glabrous.—*E. B. S.* 2747.—Fl. small, yellowish; spur remarkably short and conical.—Bogs. Skye. Black Isle, Ross. P. VI. S.

4. *P. lusitanica* (L.); *spur cylindrical blunt decurved shorter than the nearly equal limb of the cor., caps. globose, scape downy*.—*E. B.* 145.—Fl. small, pale yellowish, spur short and cylindrical.—[*P. villosa*, distinguished from this by its acute spur and obconical capsule, may be expected in the North of Scotland.]—Bogs in the Western parts of the country.—P. VI.—IX. E. S. I.

## 2. UTRICULA'RIA Linn. Bladderwort.

\* L. spinose-ciliate.

1. *U. vulgaris* (L.); spur conical, roundish 3-lobed *upper cor.-lip about equalling the bilobed palate, margins of lower lip deflexed at right angles all round*, ped. scarcely thrice as long as ovate bracts thick reflexed with fruit, l. pinnate-multifid, bladders upon the leaves.—*E. B.* 253.—Fl. bright yellow, rather large. Scape 4—6 in. high; fl. lemon-coloured. St. floating in the water. Bladders  $\frac{1}{2}$  in. long. P. VI.—VIII. Greater Bladderwort. E. S. I.

2. *U. neglecta* (Lehm.); spur directed upwards conical projecting, ovate-oblong blunt or emarginate *upper cor.-lip 2 or 3 times as long as the small palate, lower lip with a broad flat spreading margin*, ped. 4—5 times as long as lanceolate bract erect with fr., l. pinnate-multifid, bladders on leaves.—*J. of B.* xiv. 142.—Closely resembling Sp. 1. Ped. slender. Bladders  $\frac{1}{10}$  in. long. Fl. yellow.—Pools. P. VI.—VIII. E.

3. *U. intermedia* (Hayne); spur conical adpressed, *upper lip twice as long as the inflated palate, lower lip with a broad flat spreading margin*, ped. 4—5 times as long as ovate bract erect with fruit, l. 3-parted, segments linear forked, *bladders separate from the leaves*.—*E. B.* 2489.—Fl. paler with a much longer upper lip than in Sp. 1. Bladders on leafless shoots. Increasing by buds at the end of the shoots and seldom flowering.—Ditches and pits, rare. P. VIII. E. S. I.

\*\* L. not ciliate.

4. *U. minor* (L.); *spur very short blunt*, upper lip equalling the palate, lower lip ovate with a nearly flat spreading margin, l. repeatedly forked, bladders upon the leaves.—*E. B.* 254.—Scarcely any spur. Fl. small. Sep. roundish, acuminate. Plant much smaller than either of the others.—Ditches and pits. P. VI.—VIII. *Smaller Bladderwort*. E. S. I.

[5. *U. Bremii* probably grows in Moss of Inshoch, Nairn, and Loch of Spynie. It has larger fl. than Sp. 4, a short conic spur, and orbicular flat lower lip. See *J. of B.* xiv. 142.]

#### Order LXI. PRIMULACEÆ.

Cal. 4—7-cleft, permanent inferior (except <sup>Ov.</sup> in *Samolus*). Cor. regular, 4—7-fid (or none in *Glaux*). Stam. upon the cor. opposite to its segments. Ovary free, 1-celled, with a free central placenta. Style 1. Stigma capitate. Fr. a capsule. Seeds peltate.—No stipules.

Tr. I. *HOTTONIEÆ*. Ovary superior. Caps.-valves 5. Hyle basal.

1. *HOTTONIA*. Cal. 5-parted, *divided almost to its base*. Valves *connected at the top*. Cor. salvershaped, *stam. inserted and included in cor.-tube*.

Tr. II. *PRIMULEÆ*. Ovary superior. Caps.-valves 5. Hyle ventral.

2. *PRIMULA*. Cal. tubular, 5-fid. Cor. *salvershaped, tube cylindrical up to the insertion of the stamens*. Stam. 5, inserted and included in the cor.-tube. Caps. many-seeded, 5-valved with 10 teeth.

3. *CYCLAMEN*. Cal. bellshaped, half 5-cleft. Cor. *with a short bellshaped tube and 5-parted reflexed limb*. Stam. 5, inserted at the bottom of the cor.-tube, included. Caps. many-seeded.

4. *LYSIMACHIA*. Cal. 5-parted. Cor. *rotate, scarcely any tube, limb 5-parted*. Stam. 5, at the base of the cor. Caps.



with 5 valves (in *L. nemorum* sometimes 2-valved or indehiscent, in *L. thyrsoflora* few-seeded).

5. TRIENTALIS. *Cal.* 7-parted. *Cor.* rotate, 7-parted, tube none. *Stam.* 7, inserted at the base of the cor. *Caps.* many-seeded, opening with 5 revolute fugacious valves. *Seeds* with a netlike coat.
6. GLAUX. *Cal.* bellshaped, 5-parted, coloured. *Cor.* none. *Stam.* 5, inserted at the base of the calyx. *Caps.* few-seeded (about 10).

Tr. III. ANAGALLIDÆ. Ovary superior. *Caps.* opening transversely all round. *Hyle* ventral.

7. ANAGALLIS. *Cal.* 5-parted. *Cor.* rotate or funnelshaped, tube none, limb 5-parted. *Stam.* 5, at the base of the cor. *Caps.* many-seeded.
8. CENTUNCULUS. *Cal.* 4—5-parted. *Cor.* with a subglobose inflated tube and spreading 4—5-parted limb. *Stam.* 4 or 5, inserted in the throat of the cor. *Caps.* many-seeded.—*Fl.* usually 4-parted.

Tr. IV. SAMOLEÆ. Ovary inferior. *Caps.* opening by valves. *Hyle* basal.

9. SAMOLUS. *Cal.* 5-parted, its tube adhering to the lower half of the germen, persistent. *Cor.* salvershaped, tube short, limb 5-parted with interposed converging scales. *Stam.* 5, inserted near to the base of the cor.-tube. *Caps.*  $\frac{1}{2}$ -inferior, many-seeded, opening with reflexed teeth.

### Tribe I. *Hottoniæ*.

#### 1. HOTTONIA *Linn.* Water-Violet.

1. *H. palustris* (L.); fl. whorled stalked upon a long solitary cylindrical stalk, cor. exceeding the calyx, sep. subacute equaling cal.-tube, l. pectinate.—*E. B.* 364.—*L.* submerged, crowded. *Fl.* rising above the water, pale pink. *Style* exceeding the cal., *stam.* inserted in tube, *anth.* and filaments about equal in length; or style falling short of cal., *stam.* inserted at top of tube, filaments 3 or 4 times as long as the anthers.—Ponds and ditches.  
P. V. VI. E. I.

Tribe II. *Primuleæ*.2. *PRI'MULA* Linn. Primrose.\* *Cal.-tube angular. L. not mealy.*

1. *P. vulgaris* (Huds.); *l. oblong-ovate* tapering downwards wrinkled crenate, ped. villose radical 1-flowered, *cal. tubular, teeth lanceolate-subulate* very acute, cor.-limb flat with a circle of scalelike folds at the slightly contracted mouth, *caps. ovate*  $\frac{1}{2}$  the length of *cal.*, long straightish teeth of fr.-cal. meeting at top.—*E. B.* 4. *P. acaulis* Jacq.—Scape rudimentary. Fl. erect. *Cal. villose*.— $\beta$ . *P. variabilis* (Goup.); *l. slightly contracted* below, ped. raised on a scape, fl. erect. *Sy. E. B.* 1132, 1133. Often taken for *P. elatior* and called *Oxlip*.—Woods and thickets. P. III.—V. *Primrose.* E. S. I.

2. *P. véris* (L.); *l. ovate abruptly contracted below* then attenuate wrinkled crenate, scape tomentose umbellate many-flowered, *cal. bellshaped, teeth short ovate*, cor.-limb concave with a circle of scalelike folds at the slightly contracted mouth, *caps. oval*  $\frac{1}{2}$  the length of *inflated cal.*, short teeth of fr.-cal. converging.—*E. B.* 5. *P. officinalis* Jacq.—*Cal. tomentose. Cal.-teeth*  $\frac{1}{3}$  of the length of the tube, blunt or slightly acute. Cor.-segm. cordate.—Probably hybrids between this and Sp. 1 are mistaken for *P. elatior*.—Meadows and pastures. P. IV. V. *Cowslip.* Paigle. E. S. I.

3. *P. elatior* (Jacq.); *l. ovate abruptly contracted below* then attenuate wrinkled denticulate, scape umbellate many-flowered, *cal. tubular, teeth lanceolate* acute, cor.-limb concave, segments obcordate-oblong, its *tube not crowned nor contracted at the mouth*, *caps. linear-oblong equalling cal.*, teeth of fr.-cal. patent.—*E. B.* 513.—Cor.-segm. almost square; limb rarely flat. Outer fl. nodding. Fr. erect.—Clayey woods and meadows in the Eastern Counties. P. IV. V. *Oxlip.* E.

\*\* *Cal.-tube not angular. L. mealy beneath.*

4. *P. farinosa* (L.); *l. obovate-lanceolate, cal. oblong-ovate, teeth linear*, cor.-limb flat, *segments obcordate rounded below* distant as long as the tube, *caps. twice as long as calyx*.—*E. B.* 6.—*Umbellate.* Fl. pale lilac with a yellow centre; but the colour and breadth of segments very variable. Germs obovate. Stigma capitate. Rarely the scape is wanting and the fl. are amongst the leaves.—North of England and South of Scotland. P. VI. VII. *Bird's-eye Primrose.* E. S.

5. *P. scot'ica* (Hook.); l. obovate-lanceolate, cal. swollen, teeth short ovate blunt, cor.-limb flat its segments broadly obcordate approximate half the length of the tube, caps. scarcely exceeding calyx.—*E. B. S.* 2608.—Half as large as the preceding. Umbellate. Fl. bluish purple with a yellow centre. Germen globose. Stigma with 5 points. Rarely scape wanting.—Sandy heaths of the extreme North of Scotland. P. V.—VIII. S.

### 3. CYCLAMEN Linn. Snow-bread.

†1. *C. hederifolium* (Willd.); l. cordate-ovate angular denticulate, cor.-throat 5-angled, sep. ovate acuminate denticulate.—*Sy. E. B.* 1136, 1137.—St. a large depressed tuber. L. after the fl., with wavy white blotches above. Fl. pink with a red base (rarely white in *β. ficariifolium* *Sy. E. B.* 548). Fr.-ped. rolled up spirally.—Woods near the borders of East Sussex and Kent! *Mr. W. W. Saunders.* P. VIII. E.

### 4. LYSIMACHIA Linn. Loose-strife.

1. *L. thyrsiflora* (L.); racemes axillary stalked dense, l. opposite lanceolate.—*E. B.* 176.—St. 1—2 ft. high; Fl. small, very many. Cor. divided almost to its base into narrow segments separated by minute teeth, yellow and as well as the cal. spotted with orange. Stam. combined below into a short ring.—Marshes in the North. P. VI. VII. E. S.

2. *L. vulgáris* (L.); st. erect, panicles compound terminal and axillary, l. ovate or ovate-lanceolate nearly sessile opposite or 3 or 4 in a whorl, cor.-segments entire with glabrous edges, *stam.* 5 combined through  $\frac{2}{3}$  of their length.—*E. B.* 761.—St. 2—3 feet high. L. variable in size, shape and pubescence. Panicle much branched or nearly simple. Fl. yellow. Starved forms pass for *L. punctata*.—Sides of rivers and pools. P. VII. E. S. I.

[*L. punctata* (L.); st. erect, ped. axillary opposite or whorled 1-fl., l. ovate-lanceolate slightly stalked opposite or whorled, upper l. narrower, cor.-segm. ovate glandular-ciliate, *stam.* 5 combined below.—*Sy. E. B.* 1142. *L. verticillata* Bieb.—St. 1—1½ ft. high. St. and l. downy. Sep. narrowly lanceolate. Cor. yellow. Ped. rarely branched, downy, falling short of leaves.—Dulverton, Devon! Hingham, Norf.! P. VI. VII.] E.

[*L. ciliata* (L.); st. erect, ped. axillary opposite or whorled l. opposite ovate-lanceolate subcordate with ciliate stalks, cor.-segments roundish crenate, filaments 10 free 5 sterile.—*E. B. S.* 2922.—Fl. yellow. St. 3 feet high?—American. Serbergham, Cumb.! and was there in 1815. P. VI. VII.] E.

3. *L. Nummulária* (L.); *st. prostrate* creeping, *fl. solitary* axillary, sep. cordate-ovate prolonged, *filaments 5 glandular connected at the base*, l. opposite roundish or ovate shortly stalked.—*E. B.* 528.—Fl. occasionally in pairs, yellow.—Damp places. P. VI. VII. *Moneywort.* E. S. I.

4. *L. nem'orum* (L.); *st. prostrate*, fl. axillary solitary, *sep. linear-lanceolate*, *filaments 5 smooth distinct*, l. opposite ovate acute shortly stalked.—*E. B.* 527.—Ped. longer than the l. Caps. 5-valved but usually dividing longitudinally into two parts, sometimes indehiscent. Fl. yellow. Stam. distinct.—Woods and damp shady places. P. VI.—VIII. E. S. I.

### 5. TRIENTA'LIS Linn.

1. *T. europæ'a* (L.); l. oblong-obovate blunt.—*E. B.* 15.—St. 4—6 in. high, with the l. mostly collected at the top. Fl. on slender peduncles, white with a yellow ring. Parts of the fl. and fr. varying from 7 to 9 in each whorl. Valves of the caps. soon falling off.—North of E.; Highlands of S. P. VI. E. S.

### 6. GLAUX Linn. Black Saltwort.

1. *G. marit'ima* (L.).—*E. B.* 13.—St. mostly procumbent. L. opposite, ovate, glabrous. Fl. axillary, sessile, pink, with blunt segments. Remarkable in this Order by its want of petals.—Sea-shores and salt marshes. P. VI.—VIII. E. S. I.

## Tribe III. *Anagallideæ.*

### 7. ANAGAL'LIS Linn. Pimpernel.

1. *A. arven'sis* (L.); *st. procumbent* or erect, fl. axillary solitary, cor. rotate, l. opposite sessile ovate or ovate-oblong.—Pet. slightly exceeding the cal., crenate. Filaments distinct.—*a. A. arvensis* (Sm.); *st. mostly procumbent*, pet. fringed with minute glandular hairs usually scarlet, l. ovate. Fl. sometimes flesh-coloured (*A. carnea* Schrank), wholly white or white with a pink eye. *E. B.* 529.—*β. A. cœrulea* (Sm.); *st. mostly erect*, pet. without glandular hairs (usually blue), l. ovate-oblong. *E. B.* 1823. Probably distinct. Mr. Borrer suspected that each varies with red or blue flowers.—Corn-fields and sand-hills by the sea. A. ? VI. VII. *Scarlet Pimpernel.* E. S. I.

2. *A. tenel'la* (L.); *st. procumbent* rooting, fl. axillary solitary, l. opposite stalked roundish, *cor. funnelshaped*, pet. much exceeding the calyx entire, *filaments connected below*.—*E. B.* 530.

—Ped. long. Fl. rather large, rose-coloured. L. nearly sessile. Spongy bogs. P. VII. VIII. *Bog-Pimpernel*. E. S. I.

8. CENTUN' CULUS *Linn.* Bastard Pimpernel.

1. *C. minimus* (L.); l. alternate ovate acute, fl. nearly sessile, cor. without glands at the base.—*E. B.* 531.—Usually very minute. St. usually prostrate. Cor. very small, pale rose-colour.—Damp sandy and gravelly places. A. VI. VII.

E. S. I.

Tribe IV. *Samoleæ*.

9. SAM'OLUS *Linn.* Brook-weed.

1. *S. Valerian'di* (L.); l. obovate or roundish blunt, upper l. blunt with a point, racemes many-flowered ultimately elongated, caps. subglobose.—*E. B.* 703.—Remarkable in this Order by its cal. adhering to the germen and by having a crown to the small white corolla.—Damp watery places. P. VII. VIII.

E. S. I.

Order LXII. PLUMBAGINACEÆ.

Cal. 5-cleft, persistent, inferior, plicate. Cor. regular, 5-fid or nearly 5-petalous. Stam. 5, hypogynous, or adnate to the base of and opposite to the pet. Ovary free, of 5 carpels, 1-celled, 1-seeded; ovule 1, pendulous by a stalk arising from the bottom of the cell. Styles 5. Fr. a utricle. Seed inverted. Embryo in the axis of farinaceous albumen. Radicle superior.—Stip. 0.

1. STATICE. Fl. spiked. Cal. scarious above. Cor. 5-parted, Styles glabrous. Caps. not bursting.

2. ARMERIA. Fl. in a head with an inverted cylindrical sheath. Styles hairy below. Caps. not bursting.

1. STAT'ICE *Linn.* Sea-Lavender.

\* *L. pinnately veined. Cal.-segm. with intermediate teeth.*

1. *S. Limonium* (L.); l. elliptic-oblong stalked mucronate 1-ribbed strongly veined, st. subterete branched above corymbose, spikelets 1—3-flowered ascending forming dense 2-ranked patent or recurved spikes, cal.-segments entire acute, outer bract pointed rounded on the back.—*E. B.* 102.—St. 6—18 in. high, usually not branched in its lower half, often not until near the

corymbose top. Spikes short; spikelets densely imbricate. Fl. purplish.— $\beta$ . *S. serotina* (G. & G.); pan. lax much spreading. *Sy. E. B.* 1157.—Muddy salt marshes.  $\beta$ . Southern coast. P. VII.—IX. E. S.

2. *S. Bahusien'sis* (Fries); l. oblong-lanceolate stalked mucronate 1-ribbed faintly veined, st. slightly *angular* usually branched from below the middle paniced, spikelets 1—3-flowered unilateral rather distant forming *lax erect or incurved spikes*, cal.-segments acute denticulate, *outer bract pointed keeled on the back*.—*S. variflora*, Drej. *E. B. S.* 2917.—St. 6—18 in. high, *not corymbose*, much branched below. Spikes long; spikelets often 1-flowered, not imbricate. Fl. purplish.—Muddy salt marshes. P. VII. VIII. E. S. I.

\*\* *L. not pinnately veined. Cal.-segm. without interm. teeth.*

3. *S. Dodart'ii* (Gir.); l. obovate-spathulate narrowed into a broadly winged stalk 3-veined below, st. rigid with thick but *not sterile branches*, spikelets 2—4-flowered densely imbricate 2-ranked forming *linear thick suberect spikes*, cal.-segments blunt entire.—*Sy. E. B.* 1160.—St. usually not branched in its lower half; branches often simple, short. Inner bracts obovate, very blunt, with white margins or slightly pink. Fl. purple. Anth. linear.—Rocky shores. Portland. *Henslow!* P. VII. VIII. E.

4. *S. occidentalis* (Lloyd); l. lanceolate-spathulate rather acute narrowed into a long winged stalk obscurely 3-veined below, scapes slender wavy forked branched from near the base, *few lowest branches sterile*, spikelets 2—4-flowered imbricate 2-ranked forming *linear slender suberect spikes*, cal.-segments blunt entire without intermediate teeth.—*S. binervosa* G. E. Sm. *E. B. S.* 2663.—St. usually branching quite from the base; branches repeatedly forked, long, often rough. Inner bracts oval with broad membranous edge, deeply tinged with pink. Anth. oval.— $\beta$ . *intermedia* (Syme); all the branches flowering, spikelets stouter.—Rocky shores. P. VII. VIII. E. S. I.

5. *S. caspia* (Willd.); l. obovate or lanceolate-spathulate narrowed into a petiole, st. branched from near the base granular-rough, *branches many slender repeatedly and acutely forked* uppermost alone bearing dense terminal spikes of 2—3-flowered 2-ranked spikelets, cal.-segments ovate cuspidate denticulate.—*S. reticulata* Sm. *E. B.* 328.—Remarkable by its much divided sterile branches which fork at an acute angle. L. small.—Muddy sea-shores of Norf., and at Frieston, Linc. P. VII. VIII. E.

## 2. ARMERIA Willd. Thrift.

1. *A. maritima* (Willd.); *l. linear 1-veined*, scape downy, inv.-bracts very blunt 1—3 outer ones mucronate, cal.-segments acute, cal.-tube hairy or pilose-striate.—L. all radical, many, narrow. Fl. rose-coloured or white.—*a.* L. flattish above very slender 1-veined. *A. maritima* (Boiss.), *A. pubescens* Link.—*β.* *planifolia* (Syme); *l. nearly flat above broadly linear blunt, early l. 3-veined.* *Sy. E. B.* 1153.—*γ.* *A. duriuscula* (Bab.); *l. subtriquetrous channelled above very slender 1-veined.* *A. pubigera* and *A. duriuscula* Bab.—Muddy and rocky sea-shores, also on mountains. *β.* Scottish Highlands, Ben Lawers! *γ.* Southern sea-shore. P. IV.—IX. E. S. I.

[*A. plantagin'ea* (Willd.); *l. 3—5-veined linear-lanceolate with a narrow membranous margin, scapes glabrous rough, outer inv.-bracts triangular or lanceolate cuspidate, others ovate or obovate with a broad membranous margin blunt, cal.-segm. with long points.*—*E. B. S.* 2928.—Distinguished by its leaves.—Jersey. P. VI. VII.]

## Order LXIII. PLANTAGINACEÆ.

Cal. 4-parted, persistent, imbricate, inferior. Cor. 4-parted, regular, scarious. Stam. 4, hypogynous, or at the base of the tube, alternate with the segments of the cor.; filaments at first doubled inwards. Ovary free, of 1 carpel, 1-celled; or 2—4-celled. Seeds peltate or erect. Style 1. Caps. opening transversely (or indehiscent in *Littorella*). Radicle inferior.

1. PLANTAGO. Fl. perfect. Cal. 4-cleft. Cor. with an ovate tube and 4-parted reflexed limb. Stam. on the corolla. Caps. bursting transversely, 2—4-celled, 2—4-seeded.

2. LITTORELLA. Monœcious. Male fl. stalked; sep. 4; tube of the cor. cylindrical, limb 4-parted; stam. hypogynous; filaments very long. Fem. fl. sessile; sep. 3; cor. oblong, narrowed at both ends; style long; caps. 1-seeded.

## 1. PLANTA'GO Linn. Plantain.

\* *Placenta 3—4-winged, thus forming 3—4 cells. Seeds 1 in each cell.*

1. *P. Coron'opus* (L.); *l. linear pinnatifid or dentate, scape terete, spike slender, bracts subulate from an ovate base erect, midrib of lateral sep. with a ciliate membranous wing.*—*E. B.* 892.—Tube of the cor. glabrous. Extremely variable in size

and amount of pubescence, woolly or nearly glabrous. L. varying in width, nearly entire, or even doubly pinnatifid. Spikes slender,  $1\frac{1}{2}$  in. long, many-flowered; or spherical with 2—6 flowers.—Gravelly barren spots near the sea and inland. A. ? VI. VII. *Buck's horn Plantain*. E. S. I

\*\* *Placenta 2-winged. Seeds 1 in each cell. Cor.-tube pubescent.*

2. *P. maritima* (L.); l. linear channelled fleshy convex on the back, scape terete, spike cylindrical, bracts ovate mucronate, sep. not winged.—*E. B.* 175.—L. usually woolly at their base, nearly flat and broad or linear, toothed or quite entire, glabrous or hairy; 3-veined, veins equidistant. Scape glabrous or hairy. Caps. oblong-conical. Very variable in size. In a Cornish variety the l. are only 1—2 lines in length and semicylindrical, scapes very short, spikes sometimes only 3- or 4-flowered.—The mountain plant may be (1) *P. serpentina* (Vill.) or (2) *P. alpina* (L.); they are both said to have lateral veins of l. nearer the margin than midrib, (1) to have coriaceous, (2) herbaceous leaves. Symes's var. *hirsuta* is a small very hairy form from Shetland. *Sy. E. B.* 1167.—Sea-coast and on mountains. P. VI.—IX. E. S. I.

\*\*\* *Placenta 2-winged. Seeds 1 in each cell. Cor.-tube glabrous.*

3. *P. lanceolata* (L.); l. lanceolate attenuate at both ends 5-veined, scape furrowed, spike ovate or oblong-cylindrical, bracts ovate acute or cuspidate, 2 lateral sep. keeled.—*E. B.* 507.—L. nearly glabrous, length 3—12 times the breadth. Anth. and filaments yellow. Bracts and sep. black at the tip. A very variable plant. Spikes globose, scape and l. silky, neck woolly, in sandy places; spikes very long, l. very long and broad, in rich damp soil.—Meadows, pastures and sandy places. P. V.—VII. *Ribwort*. E. S. I.

[*P. Timbali* (Jord.); rootstock many-headed, bracts with scarious margins; is sometimes found in cultivated fields.]

4. *P. média* (L.); l. elliptic-ovate sessile or with short broad stalks pubescent, scape terete, spike cylindrical, bracts ovate-acuminate, sep. not keeled.—*E. B.* 1559.—L. usually lying flat on the ground, sometimes shortly lanceolate and ascending. Anth. yellow, filaments purple.—Meadows and pastures. P. VI.—IX. *Lamb's-tongue*. E.

\*\*\*\* *Placenta 2-winged. Seeds 2—4 in each cell. Cor.-tube glabrous.*

5. *P. májor* (L.); l. broadly ovate on longish channelled



stalks, scape terete, spike long, bracts ovate keeled about as long as the cal., sep. with a prominent dorsal rib.—*E. B.* 1558. *R.* xvii. 1127.—*L.* ascending. Anth. purple. Seeds about 8.—*P. intermedia* (Lilib.); l. downy, scapes terete downy arcuate-ascending, is probably distinct. It is not very rare in England. *R.* xvii. 86.—Fields and waste places. P. VI.—VIII. *Way-bread.* E. S. I.

[*P. arenária* (W. & K.), a branching leafy plant, was found abundantly on Burnham Sand-hills, Som.] E.

## 2. LITTORELLA Linn. Shore-weed.

1. *L. lacus'tris* (L.).—*E. B.* 468.—With runners. Fl. white. Fertile fl. sessile. Stalks of the male fl. 1—2 in. long. L. all radical, linear, fleshy, somewhat channelled, sometimes hairy above.—Margins of lakes, or under water when it is larger and does not flower. P. VI. VII. E. S. I.

## Division IV. MONOCHLAMYDEÆ.

(Ord. LXIV.—LXXVIII.)

With only a single perianth; that is, the cal. and cor. not distinguishable, or wanting.

### [Order LXIV. AMARANTHACEÆ.]

[Perianth 3—5-parted, scarious, persistent. Stam. hypogynous. Ovary free, 1-celled; ovule 1 or several, suspended from a free central seed-stalk. Style 1 or 0. Stigma simple or compound. Embryo curved round central farinaceous albumen.—*L.* without stipules or sheaths.

[1. AMARANTHUS. Fl. monœcious. Perianth 3—5-parted. Stam. 3—5. Stigmas 3. Caps. 1-celled, 1-seeded.

### 1. AMARANTHUS Linn.

[*A. Blitum* (L.); fl. 3-fid, 3-androus, clusters small lateral, the upper ones in a small naked spike, st. diffuse glabrous.—*E. B.* 2212.—Waste places near towns. A. VIII.] E.

### Order LXV. CHENOPODIACEÆ.

Perianth 5-parted, herbaceous, persistent. Stam. perigynous or hypogynous. Ovary free or adhering to the tube of the

perianth; ovule 1, attached to the base of the cell. Styles divided, or rarely 1. Fr. not bursting, dry, membranous, included in the perianth which often becomes enlarged or fleshy. Embryo curved round farinaceous albumen, or spiral, or doubled together without albumen; radicle next the hile.—L. without stipules or sheaths.

Tribe I. *SALSOLEÆ*. Fl. uniform, perfect. Seeds usually without albumen. Embryo spiral. *St. continuous*, leafy. *L. semicylindrical or terete*.

1. *SUÆDA*. Perianth 5-parted, without appendages. Stam. 5, from the receptacle. Stigmas 2—3, sessile. Pericarp membranous. Seed horizontal or vertical; testa crustaceous.—With bracts.
2. *SALSOIA*. Perianth 5-parted, segments ultimately with a transverse dorsal appendage. Stam. 5, from an hypogynous ring. Styles 2. Pericarp membranous. Seed horizontal; testa membranous.—With bracts.

Tr. II. *BETEÆ*. Fl. uniform, perfect. Seeds with albumen. Embryo curved round the circumference of the seed. *St. continuous*, leafy. *L. flat*.

3. *CHENOPODIUM*. Perianth 3—5-parted, persistent, unaltered. Stam. 5, from the receptacle. Stigmas 2—3. Pericarp thin, free. Testa crustaceous. Seed vertical or horizontal. Without bracts.
4. *BETA*. Perianth 5-parted, persistent. Stam. 5, from a fleshy ring. Styles 2—3. Pericarp imbedded in and adhering to the fleshy tube of the perianth. Seed horizontal, attached laterally. Testa membranous.

Tr. III. *SALICORNIEÆ*. Fl. uniform, perfect. Seeds and embryo as in Tr. II. *St. jointed*, leafless. *Stam. less than 5*.

5. *SALICORNIA*. Perianth fleshy, tumid, undivided, imbedded in an excavation of the rachis. Stam. 1—2. Style very short, stigma bifid. Pericarp membranous. Seed vertical, covered by the persistent perianth.

Tr. IV. *ATRIPLICEÆ*. Fl. monœcious, of two forms, rarely perfect. Seeds, embryo, and stem as in Tr. II.

6. *ATRIPLEX*. Perianth 3—5-parted. Stam. 5. Style 0.—Perianth compressed, of 2 parts not connected above their

middle.—Stam. 0. Stigmas 2. *Pericarp membranous free. Testa crustaceous.* Seed vertical, attached by a lateral hile either near the base or by means of a long seed-stalk in the middle of the side; radicle basal.

7. OBIONE. Perianth 3—5-parted. Stam. 5. Style 0.—*Perianth* of 2 parts *free only at the top*, 3-toothed, wedge-shaped below. Stam. 0. Stigmas 2. *Pericarp very thin, ultimately adhering to the tube of the perianth. Testa membranous.* Seed vertical, pendulous from a long seed-stalk; radicle terminal.

### Tribe I. *Salsolæ*.

#### 1. SUÆ'DA *Forsk.* Sea-Blite.

1. *S. fruticosa* (Forsk.); st. erect shrubby, *l. blunt* semi-cylindrical, styles 3, seeds vertical smooth and shining.—*Salsola* Sm.—*E. B.* 635. *Schoberia* Mey.—St. 2—3 feet high, with many erect leafy branches and axillary flowers.—Sandy and shingly shores of south and east of England, rare. P. VII. VIII. E.

2. *S. maritima* (Dum.); st. herbaceous, *l. acute* semicylindrical, styles 2, seeds horizontal netted shining.—*E. B.* 633.—St. erect or procumbent, with many spreading branches.—Muddy sea-shores. A. VII.—IX. E. S. I.

#### 2. SAL'SOLA *Linn.* Saltwort.

1. *S. Kali* (L.); minutely hairy, st. diffuse, *l. subulate spinous* rough, fl. axillary solitary, segments of the enlarged perianth hard and tough as long as their patent rather coloured roundish wings.—*E. B.* 634.—St. angular, rigid, much branched.—Sandy sea-shores. A. VIII. E. S. I.

### Tribe II. *Beteæ*.

#### 3. CHENOPO'DIUM *Linn.* Goose-foot.

\* *Flowers all pentamerous.* † Leaves undivided.

1. *C. Vulvária* (L.); *l. ovate-rhomboidal mealy, fl. in leafless* dense spikes, perianth covering the fr., seed shining slightly rough (very small), *st. diffuse*.—*E. B.* 1034. *C. olidum* *Curt.*—L. stalked. Covered with a greasy pulverulent *fetid* substance.—Dry waste places near houses. A. VIII. IX. E. S. I.

2 *C. polyspermum* (L.); *l. ovate-elliptic*, fl. in axillary leafless cymose racemes, perianth not covering the fr., seeds shining minutely dotted blunt at the edge.—*E. B.* 1480, 1481.—Racemes more or less cymose or spicate. St. erect or procumbent. L. acute or blunt. *C. acutifolium* and *C. polyspermum* are undistinguishable.—Damp waste places. A. VIII. IX. E.

†† Leaves toothed, angled, or lobed.

3. *C. ur'bicum* (L.); *l. triangular* sinuate-dentate or nearly entire their base contracted into the petiole, *spikes erect* nearly leafless compound, perianth not covering the fr., seeds very minutely rough blunt at the edge.—*E. B.* 717.—L. with short triangular teeth. Spike approaching the stem. Seed almost as large as Rape-seed.— $\beta$ . *C. intermedium* (M. & K.); *l.* with large acute teeth.—Near Oxford. *Sm. Fl. Br.*  $\beta$ . Waste places. A. VIII. E. S. I.

4. *C. album* (L.); *l. rhomboidal-ovate sinuate-dentate* entire below, upper ones lanceolate nearly entire, fl. in compound branched nearly leafless racemes, perianth covering the fr., seeds smooth and shining blunt but keeled at the edge.— $\alpha$ . *C. candicans* (Lam.); fl. in dense spikes shorter than the sinuate-dentate leaves. *E. B.* 1723.— $\beta$ . *C. viride* (L.); fl. in lax cymose racemes exceeding the nearly entire leaves. *Sy. E. B.* 1189.— $\gamma$ . *C. paganum* (R.); fl. in lax compound spikes exceeding the sinuate-dentate leaves. *Sy. E. B.* 1190.—Cultivated and waste places. A. VII. VIII. *Fat Hen*. E. S. I.

5. *C. ficifolium* (Sm.); *l. unequally 3-lobed from a wedged shaped base*, lobes ascending, *middle lobe long oblong-lanceolate dentate blunt*, upper *l. linear-lanceolate* entire, fl. in erect nearly leafless cymose racemes, perianth nearly covering the fr., seeds minutely pitted shining blunt and not keeled at the edge.—*Sy. E. B.* 1191.—Mealy. L. blunt; middle lobe nearly equally broad throughout. Seeds smaller than in *C. album*.—Cultivated and waste ground, rare. A. VIII. IX. E. I.

6. *C. murale* (L.); *l. rhomboidal-ovate* unequally and sharply toothed entire below, fl. in divaricately-branched leafless cymes, perianth nearly covering the fr., seeds minutely granular opaque acute and keeled at the edge.—*E. B.* 1722.—Waste ground near towns and villages. A. VIII. *Sowbane*. E. I.

7. *C. hyb'ridum* (L.); *l. subcordate angulate-dentate* acuminate, teeth large distant, fl. in paniced leafless cymes, perianth not covering the fr., seeds minutely pitted opaque blunt and not keeled at the edge.—*E. B.* 1919.—Seeds very large. L. with 2—4 large teeth on each side.—Waste places, rare. A. VIII. E.

\*\* *Lateral fl.* 3—4-merous with vertical seeds, *term. fl.* 5-merous with horizontal seeds.

† Stigmas short.

8. *C. rubrum* (L.); *l. rhomboidal* irregularly toothed and sinuate entire below, *fl.* in erect compound dense leafy spikes, *seeds* very minute smooth shining *blunt or slightly keeled at the edge*.—*E. B.* 1721.—*St.* erect and often 1—2 ft. high, leafy to the top. Pericarp very loose. *Fl.* generally incomplete. *Cal.* 4- rarely 5-cleft. *Stam.* 1 or 2.—A variety with much more triangular l., shorter spikes and larger seeds, was found near London.—The var. *pseudo-botryodes* (Wats.) has a prostrate spreading st. 4—5 in. long, and a small panicle. *Sy. E. B.* 1197.—Waste places, particularly salt marshes. A. VIII. IX. E. S. I.

9. *C. botryodes* (Sm.); *l. nearly triangular* slightly toothed, thick and fleshy, *fl.* in compound dense spikes nearly leafless at the top, *seeds* very minute smooth shining *keeled at the edge*.—*E. B.* 2247.—Succulently brittle when fresh, limp and flaccid when dry. *St.* erect. Pericarp loose.—Moist sandy places near the sea, on the south-east coast. A. IX. E.

10. *C. glaucum* (L.); *l. oblong sinuate-dentate mealy beneath*, *fl.* in erect nearly simple leafless spikes, *seeds* very minute *netted granular* acutely keeled at the edge.—*E. B.* 1454.—*St.* spreading, often prostrate. *Seeds* reddish.—Rich waste ground. A. IX. E.

†† Stigmas long.

††1. *C. Bonus-Henricus* (L.); *l. triangular-hastate entire*, *fl.* in compound leafless spikes, *seeds* smooth and shining.—*E. B.* 1033.—*St.* a foot high. L. large, dark green, used instead of Spinach. *Stig.* long. *Fr.* exceeding perianth.—Waste places near villages. P. V.—VIII. *Allgood. Mercury.* E. S. I.

#### 4. BETA Linn. Beet.

1. *B. maritima* (L.); *st. many from the crown of the root prostrate*, *l. triangular-ovate* narrowed into a petiole, spikes long simple leafy, bracts lanceolate exceeding the 2—3-flowered clusters, segments of the perianth with entire keels, *stig. lanceolate*.—*E. B.* 285.—*Root* thick, fleshy. *St.* 6—12 in. long prostrate below, spreading in a circle, the ends ascending.—Sea-shores. P. VII.—IX. *Sea-Beet.* E. S. I.

[*B. vulgaris* (L.), *st.* solitary erect, *stig.* ovate, is the cultivated plant.]

Tribe III. *Salicorniæ*.5. SALICOR'NIA Linn.<sup>1</sup> Glasswort.

1. *S. herbácea* (L.); herbaceous, joints of st. compressed rather thickened upwards emarginate, branches all flowering, spikes cylindrical tapering stalked, fl. 3 on each side middle fl. much exceeding the others.—*E. B.* 415.—St. usually erect, branched, 3—12 in. high. Stam. 1 or 2.— $\beta$ . *S. procumbens* (Sm.); st. procumbent, branches divaricate. *E. B.* 2475.—Muddy sea-shores. A. VIII. IX. E. S. I.

2. *S. radicans* (Sm.); st. woody creeping, st.-joints subterete deeply notched scarcely thickened with ascending herbaceous branches some barren, spikes oblong blunt nearly sessile, middle fl. scarcely exceeding the others.—*Sy. E. B.* 1183.—St. procumbent, rooting.—Muddy sea-shores. P. VIII. IX. E.

Tribe IV. *Atripliceæ*.6. A'TRIPLEX Linn.<sup>2</sup> Orache.

A. Fertile fl. of 2 kinds; sep. of fem. fl. distinct, seed vertical; perfect fl. 3—5-parted, seed horizontal.

[*A. nitens* (Rebent.), l. shining above silvery-glaucous beneath; and *A. hortensis* (L.), l. opaque on both sides; are occasionally found.]

B. Monœcious; sep. of fem. fl. united below.

\* Stem with resinous reddish stripes.—† Leaves not lobed.

1. *A. littoralis* (L.); st. erect, l. linear or oblong-lanceolate entire or toothed, fr.-perianth rhomboidal or triangular toothed tubercled on the back.—*a. vera*; l. nearly or quite entire, fr.-perianth with patent points.—*Sy. E. B.* 1200.— $\beta$ . *A. marina* (L.); l. broader usually toothed, fr.-perianth shorter with adpressed points. *E. B.* 708.—Salt marshes. A. VII.—IX.

E. S. I.

†† Lower leaves with lateral spreading or ascending lobes.

2. *A. angustifolia* (Sm. !); st. erect or prostrate, l. lanceolate entire from an acutely wedged shaped base, lower l. with 2 ascending lobes, fr.-perianth rhomboidal acute entire with pro-

<sup>1</sup> See a paper on this genus by Mr. Woods in *Proc. Linn. Soc.* ii. 111.

<sup>2</sup> See Woods in *Phytol.* iii. 585, and in *Tourist's Flora*, 315.

*longed lateral angles* longer than the fr. and *collected into nearly simple interrupted spikes* the larger leaflike and not tubercled, seeds black and polished.—*E. B.* 1774. *A. patula* Wahl., *St.* 79. 5.—Spikes wandlike with distant clusters of fl., valves of the fr.-perianth netted.—Common. A. VII.—X. E. S. I.

3. *A. erecta* (Huds.); primary st. mostly erect, lower l. ovate-oblong with 2 ascending lobes from a bluntly wedgeshaped base irregularly sinuate-dentate, upper l. lanceolate, *fr.-perianth rhomboidal denticulate* acute more or less muricate on the back scarcely exceeding the fr. and *collected in branched dense many-flowered spikes*, seeds black and polished.—*E. B.* 2223! *Koch in St.* 79. 6.—Lower branches procumbent or ascending. Upper l. mostly entire.—Common upon cultivated land. A. VII.—X. E. S. I.

4. *A. deltoidea* (Bab.); st. mostly erect, l. mostly opposite nearly all *hastate-triangular with spreading lobes*, fr.-perianth ovate-triangular muricate on the back scarcely exceeding the fr. united only at the base, *fr. paniced in dense spikes*, seeds thick black and polished or a few dark brown and larger and with large perianths.—*a. A. deltoidea* (Bab.); l. all hastate-triangular toothed, fr.-perianth toothed muricate on back, spikes dense. *E. B. S.* 2860.—*β. salina*; st. and branches prostrate, l. often alternate, uppermost l. lanceolate entire, fr.-perianth slightly toothed or entire.—Cultivated and waste ground. *β.* Sea-coast. A. VI.—X. E. I.

5. *A. hastata* (L.); st. mostly erect, lower l. *hastate-triangular with spreading lobes*, uppermost l. lanceolate entire, *fr.-perianth triangular-rhomboidal* slightly muricate on the back exceeding the fr., *fr. in nearly simple interrupted spikes*, larger seeds dark brown rough compressed, smaller seeds black and shining.—*A. patula* Sm. *E. B.* 936. *A. latifolia* *St.* 79. 7.—Cultivated and waste ground. A. VI.—X. E. S. I.

6. *A. Babingtonii* (Woods); st. spreading procumbent or ascending with spreading branches, l. nearly *ovate-triangular* somewhat 3-lobed unequally sinuate-dentate, upper l. lanceolate dentate and often 3-lobed at the base, *fr.-perianth rhomboidal acute toothed* tubercled on the back, *clusters axillary* and terminal few-flowered, seeds minutely tubercular.—*A. rosea* Bab., *E. B. S.* 2880, not *L.*—A very variable plant. Fr.-perianth large, typically forming a diagonal square a little rounded at the lateral angles.—Sea-shore, common. A. VII.—IX. E. S. I.

\*\* Stem buff-coloured, nearly without stripes. Perianth of fr. hard and thick.

7. *A. farinosa* (Dum.); st. spreading procumbent with spread-

ing branches, l. *triangular-rhomboidal* sinuate mealy beneath, spike of male fl. dense naked, fertile fl. axillary, fr.-perianths *rhomboidal* 3-lobed with the lateral lobes truncate the back 3-ribbed the two lateral ribs often terminating in tubercles, seeds rough opaque.—*A. arenaria* Woods (not Nutt.). *A. laciniata* Sm. (L. in part). E. B. 165.—Plant hoary throughout. Fr.-perianths large, very broad.—Sandy shores. A. VII.—IX.

E. S. I.

## 7. OBI'ONE Gaert. Sea-Purslane.

1. *O. pedunculata* (Moq.); st. herbaceous wavy branched, l. obovate entire attenuate below, upper l. narrower, fr.-perianth long-stalked wedgeshaped 2-lobed with a small intermediate tooth.—E. B. 232. *Atriplex* Sm.—Muddy salt marshes near the east and south coasts; very rare. A. VIII. IX. E.

2. *O. portulacoides* (Moq.); st. woody, l. obovate-lanceolate entire attenuate below, fr.-perianth inversely triangular rounded below *sessile* with 3 equal lobes above and muricate on the back.—E. B. 261. *Atriplex* Sm.—Common on the sea-shore. P. VIII.—X. E. S. I.

## Order LXVI. POLYGONACEÆ.

Perianth 3—6-parted, imbricate. Stam. definite, from the base of the perianth. Ovary 1, free, with 1 erect ovule. Styles and stigmas several. Fr. not bursting, a nut, naked or covered by the enlarged perianth. Embryo inverted, usually on one side of farinaceous albumen; radicle remote from the hile.—Stipules usually cohering in the shape of ocreæ.

1. RUMEX. Perianth 6-parted; 3 inner segments (pet.) large with fr., connivent. Stam. 6, disposed in pairs. Styles 3. Stigmas multifid. Nut triquetrous, covered by the enlarged inner segments of perianth; embryo lateral.

2. OXYRIA. Perianth 4-parted; 2 inner segments (pet.) larger. Stam. 6. Stigmas 2, multifid. Nut compressed, with a membranous wing, larger than the persistent segments of perianth; embryo central.

3. POLYGONUM. Perianth 5-parted. Stam. 5—8. Styles 2—3. Stig. capitate. Nut trigonous or compressed; embryo lateral, incurved; cotyledons not contorted.

[4. FAGOPYRUM. Perianth 5-parted. Stam. 8. Styles 3. Stig. capitate. Nut trigonous; embryo central; cotyledons large, leaflike, plicate-twisted.]



## 1. RUMEX Linn. Dock.

\* *Fl. perfect. Herbage not acid. L. not hastate.*

1. *R. marit'imus* (L.); *enlarged pet. rhomboidal* narrow, each with a lanceolate entire point a prominent narrow oblong tubercle and on each side 2 setaceous *teeth as long as the pet., whorls crowded* many-flowered leafy, l. all linear-lanceolate narrowed at both ends.—*E. B.* 725.—Nut very small; faces elliptic.—Marshes principally near the sea. P. or B. VII. VIII. *Golden Dock.* E. S. I.

2. *R. palus'tris* (Sm.); *enlarged pet. ovate-oblong*, with a lanceolate entire point a prominent narrow oblong tubercle and on each side 2 or 3 setaceous *teeth shorter than the pet., whorls distant* many-flowered leafy, root-l. narrowly lanceolate rounded or slightly decurrent below.—*Curt.* i. 63. *Sy. E. B.* 1213.—Nut 3 times as large as that of *R. maritimus*; faces ovate. Upper-l. linear-lanceolate, narrowed below.—Marshy places, rare.—P. VII.—IX. E. S. I.

3. *R. conglomerátus* (Murr.); *enlarged pet. linear-oblong subacute*, each bearing a large tubercle and entire or *obscurely toothed at the base, whorls distant* leafy, l. oblong pointed, lower l. cordate or rounded at the base.—*R. acutus* Sm. *E. B.* 724.—Nut ovate, acute. Enlarged pet. broadest near the base; the sides nearly parallel. Unopened anth. white. Branches mostly spreading. Uppermost whorls often leafless.—Wet places. P. VI.—VIII. E. S. I.

4. *R. rupes'tris* (Le G.); *enlarged pet. narrow ovate-oblong blunt* all tubercled *entire*, whorls not very distant leafless (except 2 or 3 lowest), l. strapshaped rounded and narrowed at both ends.—*J. of B.* xiv. 2. t. 173.—Nut ovate, acute. Enlarged pet. with nearly parallel sides and very large tubercles.—Shore near Plymouth. P. VI.—VIII. E.

5. *R. sanguin'eus* (L.); *enlarged pet. narrowly oblong blunt entire* only one bearing a tubercle, *whorls distant* leafless, l. ovate-lanceolate, lower l. cordate or rounded at the base.—*E. B.* 1533.—Ped. jointed at their base. Nut ovate-elliptic, acute. Enlarged pet. broadest above their middle. Lowermost whorls often each accompanied by a leaf. Veins of the l. bright red. Branches ascending.— $\beta$ . *R. viridis* (Sibth.); veins of the l. green. Unopened anth. pale yellow. *R. nemorosus* Schrad.—Woody places, rare.  $\beta$ . Woods and roadsides, frequent. P. VI.—VIII. E. S. I.

6. *R. pul'cher* (L.); *enlarged pet. triangular-ovate netted with ribs* toothed below one principally tubercled, branches spreading,

whorls mostly leafy, lower l. fiddle-shaped or cordate-oblong blunt, upper l. lanceolate-acute, st. procumbent.—*E. B.* 1576.—St. straggling. Whorls distant. Nuts ovate, acute.—Dry waste places, local. *B. ? VII.—IX. Fiddle-Dock.* E. I.

7. *R. obtusifolius* (L. ?); enlarged pet. ovate- or oblong-triangular blunt entire or toothed veined one or more tubercled, lower l. cordate-ovate-oblong blunt, upper l. oblong or lanceolate, branches ascending.—*a. R. Friesii* (G. & G.); enlarged pet. sinuate-dentate below with subulate spreading teeth and an oblong entire point one principally tubercled. *E. B.* 1999.—*β. R. sylvestris* (Wallr.); enlarged pet. nearly entire all tubercled, nut but little longer than broad. *J. of B. xi.* 131.—Whorls usually distant and nearly leafless. Height 2—3 ft. Nut nearly twice as long as broad.—Pastures and waste ground. *β.* By Thames near Putney. *P. VII.—IX.* E. S. I.

8. *R. acutus* (L., Fr.); enlarged pet. unequal cordate dilated and toothed at the base with a small entire triangular point one principally tubercled, l. oblong-lanceolate acute, lower l. slightly cordate below.—*R. pratensis* *E. B. S.* 2757.—St. and whorls often tinged with dull red. Whorls near together but not crowded, mostly leafless. Unopened anth. white. Nuts elliptic, abundant. One enlarged pet. larger than the others, sometimes all 3 equally tubercled.—Marshy places, rare. *P. VI.—IX.* E. S. I.

9. *R. crispus* (L.); enlarged pet. cordate entire or crenulate one principally tubercled, l. lanceolate acute wavy crisped.—*E. B.* 1998.—Height 2—3 feet. St. and whorls sometimes tinged with bright red. L. narrowed or truncate below. Nut elliptic. Enlarged pet. equal, sometimes all equally tubercled.—Roadsides, fields, &c. *P. VI.—VIII. Curled Dock.* E. S. I.

[*R. elongatus* (Guss.); enlarged pet. ovate entire one tubercled, l. linear-lanceolate narrowed at both ends not wavy nor crisped.—*J. of B. xi.* 237.—St. 1—2½ ft. high. Rt.-l. 8—12 in. long, 1 in. broad. Of lax habit with distant whorls. A probable form of *R. crispus*.—Surrey side of Thames between Putney and Hammersmith, and Wye at Tintern.]

10. *R. conspersus* (Hartm.); enlarged pet. broadly cordate toothed membranous netted one tubercled, l. oblong blunt cordate or oblique at the base, petioles flat above laterally compressed.—*Sy. E. B.* 1217.—St. 2—3 feet high. Nut ovate, acute. Enlarged pet. equal.—Kinross-shire. *VI.—VIII.* S.

11. *R. aquaticus* (L.); enlarged pet. broadly cordate membranous entire or wavy without tubercles, l. lanceolate, lower l. some-

what cordate, petioles semicylindrical flat and finely margined above.—*E. B. S.* 2698. *R. domesticus* Hartm. *R. longifolius* DC.—Height 3—4 feet. L. very large. Whorls crowded, mostly leafless, forming a large dense lobed panicle. Nut elliptic.—In the North of E. and in S., preferring spots liable to be flooded. P. VII. VIII. E. S.

12. *R. Hydrolap'athum* (Huds.) ; *enlarged pet. ovate-triangular* nearly entire *all tubercled*, l. lanceolate acute *tapering below*, petioles flat but not with raised edges.—*E. B.* 2104.—St. 3—5 feet high. L. often more than a foot long, sometimes cordate at the base. Whorls crowded, mostly leafless. Nut elliptic.—Ditches and river-sides. P. VII. VIII. *Great Water-Dock.* E. S. I.

13. *R. maximus* (Schreb.) ; *enlarged pet. triangular-cordate* denticulate below *all tubercled*, radical l. oblong-acute *obliquely cordate below*, petioles flat or broadly channelled above with raised edges.—*St.* 73. 16. *J. of B.* xii. t. 140—St. 3—5 feet high. L. very long, whorls crowded, mostly leafless.—Near Lewes, Winchester, Kelvedon in Essex, and in Scilly Isles.—Perhaps not distinct from Sp. 12. P. VII. VIII. E.

14. *R. alpinus* (L.) ; *enlarged pet. cordate-ovate* membranous entire or denticulate without tubercles, l. *roundish cordate-blunt with channelled petioles*, upper l. *ovate*.—*E. B. S.* 2694.—Whorls crowded, mostly leafless. Nut elliptic.—Formerly cultivated. P. VII. *Monk's Rhubarb.* E. S.

\*\* *Fl. dioecious or polygamous. Herbage acid. Sorrel.*

15. *R. Acetosa* (L.) ; *enlarged pet. roundish-cordate* entire membranous with a *very minute tubercle* at the base, *sep. reflexed*, l. oblong sagittate, stipules laciniate-dentate.—*E. B.* 127.—Height 1—2 feet. Whorls leafless. Nut elliptic with acute angles.—Woods. P. V. VI. E. S. I.

16. *R. Acetosella* (L.) ; *dioecious, petals scarcely enlarged ovate not tubercled closely adpressed to nut, sep. ascending*, l. lanceolate-hastate or linear with entire lobes, stipules torn.—*E. B.* 1674.—Height 6—10 in. Whorls leafless. L. very variable in breadth.—Dry gravelly places. P. V.—VII. *Sheep's Sorrel.* E. S. I.

[*R. scutatus* (L.), polygamous, l. hastate-ovate slightly fiddle-shaped, is naturalized near Edinburgh &c. *Sy. E. B.* 1222.]

## 2. OXYR'IA Hill. Mountain-Sorrel.

1. *O. reniformis* (Hook.).—*E. B.* 910.—St. 8—10 in. high, usually leafless. L. radical, reniform, slightly notched at the

end; veins radiating from the insertion of the long footstalk. Pedicels thickening upwards, in a spikelike raceme. Permanent pet. not enlarged.—Lofty mountains. P. VII. VIII. E. S. I.

### 3. POLYG'ONUM Linn.

\* *Rhizomatous. Stem simple. Ocreæ truncate. Nut triquetrous. Stam. 8. Styles 3. Bistorta.*

1. *P. Bistor'ta* (L.); *raceme cylindric dense, l. ovate subcordate, root-l. with winged footstalks, faces of nut ovate smooth.—E. B. 509.—St. 1—1½ foot high. Rhizome large. Fl. flesh-coloured.—Moist meadows, rare. P. VI. and X. Snakeweed.*  
E. S. I.

2. *P. vivip'arum* (L.); *raceme slender lax bulbiferous bearing fl. on its upper part, l. linear-lanceolate with revolute margins, lower l. elliptic with wingless footstalks, face of nut ovate-lanceolate smooth and shining.—E. B. 669.—Height 4—8 in. Fl. on the lower part of the raceme replaced by small red bulbs. Fl. white, sometimes very few in number.—Mountain pastures. P. VI. VII.*  
E. S. I.

\*\* *Ocreæ truncate. Root fibrous. Nut compressed or subtriquetrous. Stam. 4—8. Styles 2—3. Racemes spikelike.—Fl. purple or white. L. without or with a central dark spot. Persicaria.*

† Creeping. Perennial. Anthers protruded.

3. *P. amphib'ium* (L.); *raceme dense ovate-cylindrical, l. stalked ovate-oblong (floating) or oblong-lanceolate or narrow-lanceolate rough at the margins, ocreæ membranous close, nut compressed smooth shining, stam. 5.—E. B. 436.—St. long when floating. Racemes generally solitary and terminal. Fl. rose-coloured. Stam. protruded or not. Very variable in the form of its leaves according to its habitation.—Floating or growing upon mud or on boggy ground. P. VII.—IX.*  
E. S. I.

†† Root fibrous. Annual. Anthers included.

4. *P. lapathifólium* (L.); *raceme oblong cylindrical dense, l. oblong-lanceolate or ovate attenuate at both ends glandular beneath, ocreæ close not fringed the upper ones shortly fringed, peduncles and perianth rough with glands, nut compressed its faces roundish acuminate concave smooth shining scarcely covered by the prominently veined perianth, styles distinct at length divergent and reflexed.—E. B. 132.—Height 1—2 feet. St. sometimes spotted, glandular or glabrous; joinings more or less*

thickened. Racemes axillary or terminal, 1—2 together. Fl. pale. Bracts auricled.—Waste and damp places. A. VII.—IX.  
E. S. I.

5. *P. maculatum* (Dyer); raceme long slender, l. lanceolate much attenuate at both ends wavy glandular beneath, *ocreae* loose shortly fringed the floral ones horned, *ped.* and perianths rough with glands, nut compressed its faces roundish acuminate concave shining covered by the strongly veined perianth, styles connected below at length divergent and reflexed.—*P. laxum* E. B. S. 2822. *P. nodosum* R. I. 496 (not Pers.?). *Persicaria maculata* Gray (1821). *J. of B.* ix. 34.—Smaller than the preceding. St. often prostrate; joinings very thick. Racemes slender and interrupted (the typical form); or thicker and continuous (as in *E. B. S.*), and l. white and woolly beneath.—Damp gravelly places. A. VII.—IX.  
E. S. I.

6. *P. Persicaria* (L.); raceme compact ovate-oblong cylindrical, l. lanceolate flat minutely tubercled, *ocreae* loose strongly fringed, *ped.* and perianths smooth, nut compressed and gibbous on one side or trigonous its faces roundish acuminate smooth scarcely covered by the obscurely veined perianth, styles connected halfway up at length patent.—*Sy. E. B.* 1237.—St. 1—2 feet high. L. more or less hairy on both sides; sometimes woolly beneath, when it is *P. incanum* of authors. *Ped.* sometimes slightly hairy.— $\beta$ . *P. biforme* (Wahl.); racemes slender, much resembling *P. maculatum*. *E. B.* 756.—Waste and damp ground. A. VI.—X.  
F. S. I.

7. *P. mite* (Schrank); raceme erect filiform interrupted, l. lanceolate slightly wavy, *ocreae* loose funnelshaped pilose, strongly fringed without glands, perianths without glands, nut (large), compressed its faces roughish ovate acute rather shining convex, *stam.* 5, styles connected halfway up.—*E. B. S.* 2867.—St. 1—3 feet high, often much branched. *Ocreae* all fringed. Racemes thickening upwards.—Wet places. A. VI.—IX.  
E.

8. *P. Hydrop'iper* (L.); raceme drooping filiform interrupted, l. lanceolate wavy, *ocreae* ventricose glabrous fringed glandular, perianths glandular, nut (large) compressed its faces ovate acute rugose-punctate opaque convex, styles 2 nearly distinct.—*E. B.* 989.—St. 1—3 feet high. Upper *ocreae* funnelshaped, scarcely fringed. Nut rounder than in *P. mite*. Racemes sometimes erect.—Wet places. A. VIII. IX. *Water-Pepper*. E. S. I.

9. *P. minus* (Huds.); raceme erect filiform slender lax, l. linear-lanceolate flat, *ocreae* close pilose all fringed without glands, perianths without glands, nut (small) compressed its face ovate acute

smooth shining convex, styles connected for at least half their length.—*E. B.* 1043.—St. usually procumbent, diffuse. Racemes ascending. Much smaller than *P. Hydropiper*, fl. and fr. only half the size.—Wet gravelly places, rare. A. VIII. IX. E. S. I.

\*\*\* *Ocreæ* 2-lobed. *Root fibrous. Nut triquetrous. Stam.* 8.  
*Styles* 3. *Fl. axillary, 1—3 together. Avicularia.*

10. *P. aviculäre* (L.); l. lanceolate or elliptic stalked, *ocreæ lanceolate acute with few distant simple veins at length torn, nut striate with raised points opaque about equalling the perianth.*—*E. B.* 1252. *Sy. E. B.* 1229—1231.—L. usually blunt, sometimes acute, broad or narrow. Fl. either very distant and scattered, or so much collected as almost to form a leafy spike. St. erect or procumbent. *Ocreæ* sometimes long and much torn. Nut usually falling short of the perianth, but sometimes slightly exceeding it. Very variable. Some botanists think that it includes several species; but I am quite unable to divide what I have seen similarly.— $\beta$ . *P. littorale* (Link); st. long, diffuse, prostrate, l. thick broad, nut minutely punctured finely striate.—Waste places. Sands by the sea. A. V.—IX. *Knot-grass.*  
E. S. I.

11. *P. Rãii* (Bab.); l. elliptic-lanceolate flat, *ocreæ lanceolate acute with few distant simple veins at length torn, nut smooth shining exceeding the perianth.*—*E. B. S.* 2805.—St. long, straggling, prostrate. L. bending towards the stem. Filaments broader at the base. Resembling *P. aviculäre* in habit, but *P. maritimum* in fruit. It varies with smaller l. and fl.—Sandy sea-shores. A. or P. VIII. IX.  
E. S. I.

12. *P. maritimum* (L.); l. elliptic-lanceolate coriaceous with revolute edges, *ocreæ lanceolate with many branched veins at length torn, nut smooth shining exceeding the perianth.*—*E. B. S.* 2804.—St. procumbent, quite woody below, often much buried. L. convex above, diverging from the stem. Filaments broader at the base.—Sands of the southern sea-shores. Christchurch, Hants. Braunton Burrows, Dev. Falmouth. Channel Islands. P. VIII. IX.  
E.

\*\*\*\* *Ocreæ semicylindrical. Root fibrous. Nut triquetrous.*  
*Stam.* 8. *Stoles* 3. *Fl. racemose.*

13. *P. Convolvulus* (L.); st. twining angular, l. cordate-sagittate, *segments of perianth bluntly keeled, nut opaque striate with minute points.*—*E. B.* 941.—St. climbing or prostrate, much shorter than in the next species.— $\beta$ . *pseudo-dumetorum* (Wats.); perianth-segm. winged; often taken for Sp. 14.—Cultivated and waste land. A. VII.—IX. *Black Bindweed.*  
E. S. I.

14. *P. dumetorum* (L.); st. twining striate, l. cordate sagittate, segments of perianth winged, nut very smooth and shining.—*E. B. S.* 2811.—St. wiry, climbing to the height of 4 or 5 feet.—Thickets in the South, rare. A. VII.—IX. E.

#### 4. FAGOPY'RUM *Gaert.* Buck-wheat.

[*F. esculentum* (Moench); st. erect without prickles, fl. in cymose panicles, stam. 8, l. cordate-sagittate acute, nut triquetrous acute with entire angles.—*Polygonum Fagopyrum* L. *E. B.* 1044.—An escape. Sown as food for game. A. VII. VIII.]

### Order LXVII. ELÆAGNACEÆ.

Mostly dicecious. Perianth tubular; limb 2—4-toothed, in male fl. 4-parted. Stam. 3 or more, inserted in the throat. Anth. 2-celled, nearly sessile, bursting on the inner side longitudinally. Ovary free, 1-celled, with 1 erect ovule. Fr. crustaceous, enclosed within the fleshy persistent perianth. Albumen thin and fleshy. Radicle inferior.—No stipules.

1. HIPPOPHAEË. Dicecious. Male fl. in a sort of catkin; scales ovate, 1-flowered. Perianth of 2 leaves adhering by their points. Stam. 4, with very short filaments.—Female fl. solitary. Perianth tubular, cloven at the summit. Style short. Stigma long. Nut 1-seeded, in the large coloured berrylike perianth.

#### 1. HIPPO'PHAEË *Linn.* Sea Buckthorn.

1. *H. rhamnoides* (L.).—*E. B.* 425.—A thorny shrub with linear-lanceolate silvery leaves. Fl. appearing with the young leaves. Fr. orange. Height 4—6 feet.—Sandy spots and cliffs of the South-east and East coasts chiefly. Sh. V. E.

### Order LXVIII. THYMELACEÆ.

Perianth tubular, inferior, often coloured, 4—5-cleft. Stam. definite in number, in the tube. Anth. 2-celled, bursting longitudinally. Ovary free, 1-celled; ovule 1, pendulous. Fr. a nut or drupe. Albumen 0 or thin and fleshy. Radicle superior.—No stipules.

1. DAPHNE. Perianth 4-fid, deciduous. Berry fleshy, 1-seeded. Stam. 8, falling short of the perianth, inserted in the tube in 2 rows.

## 1. DAPH'NE Linn.

1. *D. Mezereum* (L.); fl. about 3 together lateral sessile, tube hairy, segments ovate acute.—*E. B.* 1381.—Fl. purple or pale red, appearing before the lanceolate l., which are narrowed below. Berries red. A small deciduous shrub.—Woods, rare. Sh. III. *Mezereon.* E.

2. *D. Lauréola* (L.); racemes axillary of about 5 glabrous drooping bracteate fl., l. lanceolate attenuate below glabrous evergreen.—*E. B.* 119.—A small evergreen shrub, 1—3 feet high, slightly branched, naked below. Fl. yellowish green, funnel-shaped. Berries bluish black.—Woods and thickets. Sh. II.—IV. E.

## Order LXIX. SANTALACEÆ.

Perianth adnate to the ovary; limb 3—5-fid; æstivation valvate. Stam. 3—5, opposite to and inserted at the base of the segments. Ovary 1-celled; ovules 1—4, pendulous from near the apex of a central placenta. Style 1. Fr. drupaceous, 1-seeded. Embryo in the axis of fleshy albumen.—No stipules.

1. THESIUM. Perianth 4—5-cleft, top- or funnel-shaped. Stam. 5, with a fascicle of hairs at their base. Style 1. Stigma simple. Nut drupaceous, crowned with the persistent perianth.

## 1. THE'SIUM Linn.

1. *T. humifusum* (DC.); st. procumbent or ascending racemose, racemes branched or simple, fl.-stalks as long as or longer than the fl., fr.-stalks patent, angles of fr.-stalks and edges of bracts and upper l. denticulate-asperous, fr. oval-oblong, l. 1-veined linear.—*R.* xi. 542. *T. linophyllum* Sm. *E. B.* 247.—Parasitical. Root woody. St. nearly always prostrate, spreading in a circle. L. very slender; veins usually obscure. Bracts 3, lower middle ones exceeding flowers. Perianth-tube very short, open, funnelshaped; segm. triangular, spreading, a tooth on each side. Fr. ovoid, longer than the persistent perianth which is usually inflexed only at the end, strongly ribbed, slightly netted.—Chalky and limestone (oolite) hills. P. VI. VII. E.

[*T. humile* (Vahl); racemes spiked, *fl. nearly sessile*, fr. strongly ribbed and netted sessile crowned with the very short inflexed perianth, *l. fleshy linear* 1-veined.—*R.* xi. 542.—I gathered two specimens of this plant near Dawlish, Devonshire, in 1829. P. VII. VIII.]



## Order LXX. ARISTOLOCHIACEÆ.

Perianth adnate to the ovary below, tubular above, with a lobed dilated usually irregular limb. Stam. 6—12, epigynous. Ovary 3—6-celled. Style simple; stigma radiant. Fr. many-seeded. Seed with a minute embryo at the base of fleshy albumen.—No stipules.

- [1. ARISTOLOCHIA. Perianth tubular, swelling at the base; mouth dilated on one side. Anth. 6, adnate to the short columnar style under the 6-lobed stigma. Caps. 6-celled.]
2. ASARUM. Perianth bellshaped, 3-fid. Stam. 12, inserted at the base of the style. Anth. attached to the middle of the filaments. Stigma 6-lobed. Caps. 6-celled.

1. ARISTOLO'CHIA *Linn.* Birthwort.

[1. *A. Clematidis* (L.); creeping, st. erect simple, l. cordate stalked glabrous, fl. clustered.—*E. B.* 398. *St.* 6. 16.—Fl. pale yellow.—Established near old ruins. P. VII. VIII.] E.

2. AS'ARUM *Linn.* Asarabacca.

1. *A. europæum* (L.); l. reniform blunt.—*E. B.* 1083.—St. short, with 2 evergreen leaves, between which there is a solitary drooping dull-green flower. Perianth-segm. ovate and incurved. Filaments extending beyond the anthers.—Woods and banks, very rare. P. V. E.

## Order LXXI. EMPETRACEÆ.

Dicœious. Perianth of hypogynous scales imbricate in several rows. Stam. equal in number to the inner row of scales and alternate with them, free. Ovary free, on a fleshy disk, 2—9-celled. Ovules solitary, ascending. Style 1. Stigma radiant. Fr. fleshy with bony cells. Embryo in the axis of fleshy albumen. Radicle inferior.—No stipules.

1. EMPETRUM. Cal. 3-parted. Pet. 3. Stam. 3 in the male flowers. Style short; stigma dilated, peltate, with 6—9 rays in the fem. flowers. Berry globose, 1-celled. Seeds 6—9.

1. EMPE'TRUM *Linn.* Crowberry.

1. *E. nigrum* (L.); procumbent, l. linear-oblong their margins meeting in a white line beneath.—*E. B.* 526.—A small leafy heathlike shrub. Fl. axillary, small, purple. Stigma with 9 rays. Berries black.—Mountain heaths. Sh. V. E. S. I

## Order LXXII. EUPHORBIACEÆ.

Fl. monœcious or diœcious. Perianth lobed or 0. Male flowers of 1 or more stamens. Anth. 2-celled. Fem. fl. of 1 superior 2—3-lobed and 2—3-celled ovary. Styles 2—3. Stigma compound or simple. Caps. opening with elasticity; cells 2—3, with 1 or 2 suspended seeds in each. Embryo in fleshy albumen. Radicle superior.—Stip. often present, but very deciduous.

1. BUXUS. Fl. monœcious.—Male. Cal. 3-parted. *Pet.* 2. *Stam.* 4.—Fem. Cal. 4-parted. *Pet.* 3. Caps. with 3 horns, 3-celled, 6-seeded.
2. EUPHORBIA. Fl. incomplete, collected into monœcious clusters consisting of 1 female and many male flowers. Involucre bellshaped with 4 or 5 divisions and 4 or 5 alternate glands.—Males consisting of a single stamen upon a pedicel, intermixed with scales and surrounding the female. Fem. of a single pistil. Styles 3. Stigmas bifid. Caps. 3-celled, bursting at the back. Seeds solitary, pendulous.
3. MERCURIALIS. Fl. diœcious or monœcious. Perianth 3-parted.—Male. *Stam.* 9—16.—Fem. Two barren filaments. Style short, forked. Caps. 2-celled. Cells 1-seeded, bursting at the back.

## 1. BUX'US Linn. Box.

1. *B. sempervirens* (L.); l. ovate-oblong coriaceous shining above, petioles ciliate, anth. ovate-sagittate.—*E. B.* 1341.—A small bushy tree 10—12 feet in height.—Dry chalky hills especially in Surrey and Kent, rare. T. IV.—VI. E.

## 2. EUPHOR'BIA Linn. Spurge.

A. Leaves with stipules, opposite. Clusters axillary.

1. *E. Pep'lis* (L.); st. procumbent forked, heads solitary, caps. keeled, seeds smooth, l. opposite stalked half-oblong-heartshaped nearly entire glabrous.—*E. B.* 2002.—St. usually much tinged with purple, glaucous. Inv.-glands rounded.—Loose sand. Southern sea-coast, rare. A. VII.—IX. E. I.

B. Stipules wanting. L. scattered (except in *E. Lathyris*).

\* *Involucral lobes roundish or transversely oval.*

2. *E. Helioscopia* (L.); umbel of 5 trifid and forked rays, bracts and l. membranous ovate-wedgeshaped serrate upwards, caps.

smooth glabrous, seeds netted rugose.—*E. B.* 883.—St. 6—18 in. high.—Waste and cultivated ground. A. VI.—IX. *Sun Spurge.* E. S. I.

3. *E. stricta* (L.); umbel of 3—5 each 3—5-fid and forked rays, general and first partial bracts and l. oblong clasping, other bracts broadly cordate blunt with a minute apiculus, inv.-lobes oval, *stam. rarely more than 2 in each inv., caps. with prominent cylindrical tubercles*, seeds smooth oval brown and shining, *seed-stalk cordate.*—*E. B. S.* 2974.—St. erect, 2—3 ft. high, much branched. Umbel moderate. Inv., caps., and seeds small.—Limestone woods. Gloucestershire and Monmouthshire. A. VII. VIII. E.

4. *E. platyphyl'la* (L.); umbel of 3—5 repeatedly forked rays, l. and general bracts long-obovate with a cordate base, partial bracts all cordate, *stam. 7—8 in each inv., inv.-lobes oval, caps. with shortly conical tubercles*, seed obovate brown and shining, *seed-stalk nearly reniform.*—*E. stricta* Sm., *E. B.* 333 (starved).—Plant more diffuse than, and quite distinct from, the preceding, erect, slightly branched. Inv., caps., and seeds twice as large.—Fields. A. VI.—VIII. E.

[*E. dulcis* (L.); umbel of 5 bifid rays, l. and general bracts long-obovate blunt narrowed to the base, partial bracts triangular-ovate with a truncate base, inv.-lobes rounded, caps. with few prominent tubercles.—*R. v.* 135.—Erect. A foot high. Umbel-rays long.—Glascoed Dingle, Llansilin. Jedburgh. P. VI.]

5. *E. hiber'na* (L.); umbel of about 5 twice-forked rays, bracts and l. ovate or oblong entire blunt, inv.-lobes reniform, caps. warted glabrous, seeds obovate smooth somewhat shining brownish.—*E. B.* 1337.—Height 1—2 feet. L. broad, usually pilose beneath.—S. and W. of Ireland and N. Devon. P. VI. E. I.

6. *E. pil'osa* (L.); umbel irregular of about 5 trifid and forked rays, bracts all elliptic glabrous, l. broadly lanceolate minutely serrate slightly hairy, inv.-lobes transversely oval, caps. warted, seeds obovate minutely punctate smooth.—*E. B. S.* 2787. *E. palustris* Forst. (not Linn.).—St. 2—4 feet high, leafy throughout, annual. Caps. usually hairy.—Shady places near Bath. P. V. VI. E.

[\*7. *E. coralloïdes* (L.); umbel of 5 trifid and forked rays, bracts ovate-oblong the tertiary ones ovate all hairy, l. lanceolate minutely serrate woolly, inv.-lobes transversely oval, caps. nearly smooth woolly, seeds obovate minutely punctate and with faint

netted bands.—*E. B. S.* 2837.—Height 2—3 feet. Usually naked below.—Slinfold, Sussex. Introduced. B. ? V. VI.] E.

\*\* *Involucral lobes triangular-lunate or with 2 horns.*

† Seeds smooth. Bracts connate.

8. *E. amygdaloides* (L.); umbel of 5 or more forked bifid rays, bracts rounded connate, l. ovate-lanceolate hairy beneath entire, inv.-lobes lunate (yellow) with 2 horns, caps with very minute tubercles glabrous, seeds roundish-ovate.—*E. B.* 256.—St. 2—3 feet high, leafy, purple below, biennial.—Woods and thickets. P. III. IV. *Wood-Spurge.* E. I.

†† Seeds smooth. Bracts separate.

[*E. Cyparis'sias* (L.); umbel of many forked rays, partial bracts reniform or cordate blunt entire, l. linear entire or on the barren shoots setaceous many close together, inv.-lobes with 2 horns, cap. tubercular.—*R. v.* 147.—It may be a native at Whitbarrow in Westmoreland. *Rev. W. H. Hawker.* The other stations belong to *E. Esula.*] E.

†9. *E. Esula* (L.); umbel of many forked rays, partial bracts cordate mucronate, l. lanceolate or sublinear narrowed below glabrous denticulate scattered, inv.-lobes with 2 horns, "caps. asperous, seeds obovate."—*E. B.* 1399. *R. v.* 146.—St. 12—18 in. high, leafy, with a few axillary leafy branches without flowers. Creeping.—Woods rare. P. VI. VII. E. S.

10. *E. Paralias* (L.); umbel usually of 5 forked rays, bracts rather reniform, l. coriaceous oblong, inv.-lobes 5 with 3 or 4 short points, caps. wrinkled.—*E. B.* 195.—Root woody, tough. Flowering st. about a foot high; barren stems shorter, many. L. closely imbricate.—Sandy sea-coasts. P. VIII. IX. *Sea-Spurge.* E. S. I.

††† Seeds rough, tubercled or pitted. L. alternate, scattered.

11. *E. portland'ica* (L.); umbels of 5 forked rays, bracts broadly rhomboidal acuminate, l. obovate or obovate-lanceolate blunt subapiculate, inv.-lobes 4 lunate with long horns, caps. rough at the angles, seeds pitted and netted almost white.—*E. B.* 441.—Glaucous, smooth. Scarcely a foot high. L. spreading. Seed-stalk conical, hooded.—Sea-coasts. P. IV.—IX.

E. S. I.

12. *E. Pep'lus* (L.); umbel of 3 forked rays, bracts ovate blunt mucronate, l. broadly ovate stalked, lower l. suborbicular, inv.-lobes 4 lunate with long horns, caps. smooth with thick-

ened rough keels, seeds ovoid pitted.—*E. B.* 959.—Light green, smooth, erect, 6—10 in. high.—A common weed. A. VII. VIII. *Petty Spurge.* E. S. I.

13. *E. exigua* (L.); umbel of 3 forked rays, bracts lanceolate acute unequal below, l. linear blunt with a mucro or acute, inv.-lobes rounded with 2 horns, caps. smooth with slightly thickened and tubercled angles, seeds angular wrinkled.—*E. B.* 1336.—Height 3—6 in., usually branched at the base. Seeds small, white.—Cornfields. A. VI.—VIII. E. S. I.

††† Seeds rough. Leaves opposite.

14. *E. Lath'yris* (L.); umbel of 3 or 4 forked rays, bracts oblong-ovate attenuate acute, l. linear-oblong sessile, upper l. cordate at the base, inv.-lobes lunate with blunt horns, caps. smooth with a dorsal line.—*E. B.* 2255. St. solitary, 2—3 feet high, purplish. L. many, in 4 rows.—Truly wild in a few stony and rocky woods, where it appears for 2 or 3 years after the bushes have been cut. Also naturalized on cultivated ground. B. VI. VII. *Caper-Spurge.* E.

### 3. MERCURIA' LIS Linn. Mercury.

1. *M. peren'nis* (L.); st. simple, l. stalked oblong-lanceolate rough, female fl. on long common stalks, creeping.—*E. B.* 1872.—St. a foot high, usually naked below. Fl. in lax spikes.—β. *M. ovata* (Steud.); l. nearly sessile ovate.—Woods and thickets. β. Hurstpierpoint, Suss. *Mr. Mitten.* P. IV. V. E. S. I.

2. *M. an'nua* (L.); st. branched, l. stalked ovate or ovate-oblong smooth, female fl. nearly sessile, root fibrous.—*E. B.* 559.—Height 6—12 in. Bright green. Much branched.—β. *M. ambigua* (L.); l. lanceolate, fl. whorled male and female intermixed. *E. B. S.* 2816.—Waste and cultivated land. A. VIII. IX. E. I.

## Order LXXIII. CERATOPHYLLACEÆ.

Fl. monœcious. Perianth free, in many divisions. Anth. 12—20, sessile, 2-celled, 2-pointed; cells again partially divided.—Perianth usually wanting. Ovary 1-celled. Seed 1, pendulous. Embryo straight; cotyledons 4, alternately smaller.—No stipules.—Only one genus.

### 1. CERATOPHYLLUM Linn. Hornwort.

1. *C. demer'sum* (L.); segments of l. linear-filiform, fr. wing-

less having a spine on each side near the base and tipped with the curved subulate style.—*E. B.* 947.—*L.* 2—4 times forked, dark green, segments of uppermost l. broader and more rigid.—Ponds and ditches, rare. *P. VI. VII.* *E. S. I.*

2. *C. submer'sum* (L.); segments of l. setaceous, fr. destitute of spines covered when ripe with cylindrical tubercles and tipped with the short curved style.—*E. B.* 679.—*J. pale green.* Young fr. smooth.—Ponds and ditches. *P. VI. VII.* *E.*

#### Order LXXIV. CALLITRICHACEÆ.

*Fl.* mostly monœcious, solitary, minute, usually with 2 white bracts, no perianth. *Stam.* 1. Filament long. *Anth.* reniform, 1-celled, opening transversely at the top.—*Perianth* none. *Ovary* 1, 4-angled, 4-celled. *Styles* 2, subulate. *Fr.* dry, 4-celled, 4-lobed; separating into 4 1-seeded indehiscent carpels. *Embryo* in axis of fleshy albumen. *Radicle* superior, long. *Cotyledons* short, terete.—*No stipules.*—Only one genus of aquatic plants.

##### 1. CALLIT'RICHE *Linn.* Water-Starwort.

*A. L.* narrowed at the base; lower linear, submerged; upper floating, broad, often spathulate and forming a rosette. *Fr.* of 4 lobes cohering in pairs.

1. *C. ver'na* (L.); fr. nearly sessile longer than broad, lobes parallel bluntly keeled on the back convex on the sides, styles erect deciduous, bracts straightish deciduous.—*E. B.* 722. *C. vernalis* Kütz. *R. v.* 129.—Floating l. ovate-spathulate, in a rosette, rarely wanting. *Fr.* small; keels of each pair of lobes converging.—Ponds and slow streams. *A. or P. IV.—IX.*

*E. S. I.*

3. *C. obtusan'gula* (Le G.); fr. nearly sessile, lobes parallel with rounded edges, styles erect or spreading persistent, bracts persistent.—*L.* all obovate, in a rosette, or lower l. linear. *Fr.* large, quite blunt and rounded at the edges.—Ditches. *A. or P. V.—IX.* *E.*

3. *C. stag'nalis* (Scop.); fr. nearly sessile, lobes winged on the back, styles erect in flower persistent and reflexed closely with fruit, bracts falcate persistent.—*E. B. S.* 2864. Floating l. round-obovate-spathulate, in a close rosette, often wanting. *Fr.* twice as large as in *Sp. 1.*, pale when dry; lobes parallel (*C. platycarpa* Kütz.) or slightly spreading (*C. stagnalis* Kütz.).—Common, especially on mud. *A. or P. V.—IX.* *E. S. I.*

4. *C. hamulata* (Kütz.); fr. broader than long nearly sessile or stalked, lobes parallel with a blunt dorsal ridge flat on the sides, styles divaricate and reflexed over the sides of the fruit, bracts falcate very deciduous.—*R. v.* 130. *Sy. E. B.* 1273.—L. linear; upper ones broader in the middle; uppermost ovate-spathulate. Bracts hooked, overlapping, perhaps confined to the male flowers. Fr. small, the size of that of *C. verna*.— $\beta$ . *C. pedunculata* (DC.); l. linear, upper l. rather broader, fr. stalked. Plant usually very small.—Lakes and streams.  $\beta$ . Marshes. A. or P. VI.—IX. E. S. I.

B. L. uniform, enlarged at the base, linear. Fr. of 4 nearly free lobes.

5. *C. truncata* (Guss.); fr. shortly stalked, lobes with a blunt dorsal ridge, styles reflexed-patent, bracts 0.—*C. autumnalis* Hook. (not L.). *E. B. S.* 2606.—L. all submersed, uniform, very abrupt at the end, translucent green.—Amberley, Suss. Glansiskin, Co. Cork. A. or P. IV.—IX. E. I.

6. *C. autumnalis* (L.); fr. nearly sessile, lobes broadly and acutely winged at the back, styles spreading, bracts 0.—*E. B. S.* 2732.—Fr. 4 times as large as in *C. verna*, dark brown, 1 or 2 of the lobes often abortive. L. dark green, all submersed, very abrupt at the end.—Rare. Anglesea. Several Scottish lakes. Cong, Co. Mayo. A. or P. VI.—IX. E. S. I.

#### Order LXXV. URTICACEÆ.

Fl. monoecious, dioecious, or rarely perfect. Perianth inferior, 4—3—6-parted, imbricate; or in the female flowers tubular notched or scalelike and open. Stam. definite, free, inserted at the base of the perianth and opposite to its lobes. Ovary free, 1-celled; ovules solitary. Stigmas 1—2. Fr. not bursting.

1. PARIETARIA. Fl. polygamous, surrounded by an involucre. Perianth bellshaped, 4-parted. Stam. 4. Style filiform.—L. alternate. Hairs simple.
2. URTICA. Fl. monoecious or dioecious. Males in loose racemes; perianth 4-parted; stam. 4. Females in capitate racemes; perianth 2-parted; stigma sessile.—L. opposite. Hairs stinging.

#### 1. PARIETARIA Linn. ' Wall-Pellitory.

1. *P. diffusa* (Koch); l. elliptic or elliptic-lanceolate 3-veined above the base, bracts combined into an involucre of two 3-lobed

segm., perianth bellshaped equalling the stam. or in the perfect fl. (red) lengthening to twice as long as the stamens.—*E. B.* 879. *Curt.* ii. 203.—St. prostrate or ascending, simple or branched below. Cymes axillary, dense; primary fem. fl. between the inv.-segm. and with 1 or 2 free bracts; each segm. bears on its face 1 or 3 fl., of which the lateral have bracts.—Old walls. P. VI.—IX. E. S. I.

## 2. URTICA Linn. Nettle.

‡1. *U. pilulifera* (L.); l. opposite ovate ovate-lanceolate or cordate-acuminate *coarsely toothed*, clusters of fr. globose stalked.—*E. B.* 148.—About 2 ft. high. Very venomous.—β. *U. Dodartii* (L.); l. ovate or ovate-lanceolate nearly entire. *Sy. E. B.* 1281.—About towns and villages in the East of England, rare. A. VI.—VIII. *Roman Nettle.* E.

2. *U. úrens* (L.); l. opposite elliptic-ovate serrate, *spikes* axillary *nearly simple two together falling short of the petiole*, seeds oblong.—*E. B.* 1236.—Scarcely a foot high, glabrous, with stinging bristles.—Common weed. A. VI.—IX. E. S. I.

3. *U. dioica* (L.); l. opposite cordate serrate, *spikes* axillary *panicled exceeding the petiole*, seeds ovate.—*E. B.* 1750.—Creeping. St. 2—3 ft. high. Hairy. Stinging. Dark green.—β. *angustifolia*; l. ovate-lanceolate rounded but not cordate at the base.—Common. P. VI.—IX. E. S. I.

## Order LXXVI. CANNABINACEÆ.

Fl. dioecious; fem. in catkins. Perianth inferior, of 5 sepals in male, open and scale-like in female flower. Stam. 5, on base of perianth. Ovary free. Ovule solitary. Styles 2, long.—With stipules.

1. HUMULUS. Fem. fl. open nearly flat, 2 together.

### 1. HU'MULUS Linn. Hop.

1. *H. Lupulus* (L.).—*E. B.* 427.—Known by its long twining stems, opposite rough 3—5-lobed serrate leaves; ovoid or globular ripe fem. catkins with ovate scales. Male fl. in loose panicles.—A true native in the south of England. P. VII. E.

## Order LXXVII. ULMACEÆ.

Fl. bisexual, not in catkins. Perianth inferior, bellshaped, 3—8-cleft, imbricate. Stam. definite in number, inserted in



tube of perianth and opposite to its lobes. Ovary free, 1—2-celled; ovules solitary. Styles 2. Fr. not bursting. With deciduous stipules.

1. ULMUS. Perianth bellshaped, 4—5-cleft, persistent. Stam. 5. Styles 2. Caps. compressed, wings all round.—Fl. before the leaves, clustered; ped. short.

### 1. ULMUS Linn. Elm.

1. *U. campestris* (Sm.); l. shortly acuminate more or less doubly serrate, ped. short, fl. 4—5-fid, lobes ciliate, fr. obovate or oblong notched, *seed-cavity chiefly above the middle of fr. and near the notch*.—*U. campestris* (L.) includes this species and Sp. 2.—*a. U. suberosa* (Sm.); l. asperous above pubescent beneath. *E. B.* 1886, 2161.—*β. U. glabra* (Sm.); l. rather coriaceous shining nearly or quite smooth above glabrous except in the axils of the veins beneath, young l. stip. and fr. with sessile glands. *E. B.* 2248. *U. stricta* (Lindl.) which has rigid erect close branches is a form of this variety.—*a.* Throughout England. *β.* South of England and Ireland. T. III.—V. *Common Elm.* E. I.

2. *U. montana* (With.); l. much acuminate doubly serrate asperous above rather downy beneath, ped. short, fl. 5—7-fid, lobes ciliate, fr. oblong or roundish notched, *seed-cavity chiefly below the middle of fr. and distant from the notch*.—*E. B.* 1887. *U. major* Sm.?—Branches large spreading. L. broad. Fr. Hoplike.—Woods and hedges. T. III. IV. *Wych Elm.* E. S. I.

## Order LXXVIII. AMENTIFERÆ.

Fl. monœcious or diœcious, rarely perfect. Barren fl. of all our plants in catkins; sometimes with a membranous perianth. Fertile fl. clustered, solitary or in catkins. Ovary with one or several cells. Stigmas 1 or more. Fruit as many as the ovaries, bony or membranous (or drupaceous in *Myrica*). Albumen usually wanting. Embryo straight or curved, plain. Radicle mostly superior.—Young leaves with stipules.

Subord. I. SALICINÆ. Diœcious. Fl. all in catkins. Fr. naked, 2-valved, 1-celled, many-seeded, Seeds erect, hairy.

1. SALIX. Catkins consisting of imbricate entire scales. Stam. 1—5. Fr. a 1-celled pod with 1—2 glands at its base. Perianth 0.
2. POPULUS. Catkins with slashed scales. Stam. 4—30

from a little oblique cupshaped perianth. Fr. almost 2-celled, with a cupshaped perianth.

Subord. II. *MYRICEÆ*. Dioecious. Fl. in catkins. Fr. drupaceous surrounded by scales become fleshy and adherent to the ovary.

3. *MYRICA*. Catkins with concave scales. Stam. 4—8. Stigmas subulate. Drupe 1-celled, 1-seeded. Perianth 0.

Subord. III. *BETULINEÆ*. Monoecious. Fl. all in catkins. Ovary 2-celled, each with 1 ovule. Fr. naked, indehiscent, 1-celled, 1-seeded. Seeds pendulous, not hairy.

4. *BETULA*. Scales of the barren catkins ternate, the middle one bearing the stamens. Perianth 0. Scales of the fertile catkin 3-lobed, 3-flowered, membranous, deciduous. Styles 2, filiform. Fr. with a membranous margin.

5. *ALNUS*. Scales of the barren catkins 3-lobed, 3-flowered. Perianth 4-parted. Scales of the fertile catkin ovate, 2-flowered, coriaceous, persistent. Styles 2. Ovary compressed, 4 minute scales at its base. Fr. not winged.

Subord. IV. *CUPULIFERÆ*. Male fl. in a catkin. Fem. solitary or clustered or spiked. Perianth adnate to the ovary (glans), with a very minute sometimes evanescent limb, surrounded by a coriaceous involucre.

6. *FAGUS*. Barren catkin globose. Perianth 5- or 6-fid. Stam. 8—15. Fertile fl. 2 together within a 4-lobed prickly involucre. Stigmas 3. Ovaries 3-cornered and 3-celled. Nut by abortion 1—2-seeded.

7. *CASTANEA*. Barren catkin long, cylindrical. Perianth 6-parted. Stam. 8—20. Fertile fl. 3 within a 4-lobed muricate involucre. Stigmas 6. Ovary 5—8-celled. Nut 1-celled with 1—3 seeds.

8. *QUERCUS*. Barren catkin long, pendulous, lax. Stam. 5—10. Perianth 5—7-cleft. Fertile fl. solitary, with a cupshaped scaly involucre. Stigmas 3. Ovary 3-celled. Nut 1-celled, 1-seeded, surrounded at the base by the enlarged cupshaped involucre.

9. *CORYLUS*. Barren catkin long, pendulous, cylindrical. Scales 3-lobed, middle lobe covering the 2 lateral lobes. Stam. 8. *Anth.* 1-celled. Perianth 0. Fertile fl. several, surrounded by a scaly involucre. Styles 2. Nut 1-seeded, enclosed in the enlarged coriaceous cut involucre.

10. *CARPINUS*. Barren catkin long, cylindrical. Scales roundish. Stam. 5—14. *Anth.* 1-celled. Fertile fl. in a lax catkin. Scales large, leaflike, 3-lobed, 2-flowered. Styles 2. Nut ovate, 1-seeded.

### Tribe I. *Salicineæ*.

#### 1. *SA'LIX* Linn.<sup>1</sup> Willow.

Sec. I. *VITISALIX* (Dumort.). Catkin and its leafy stalk usually (but not always) deciduous together, lateral, appearing with the leaves. Scales of uniform colour. Nectary of 2 pieces or urceolate; germen or stamens from the middle. Vernation convolute.

Subsec. 1. *Lycus* (Dumort.). Stam. 4—8. "Nectary urceolate, undivided." L. glossy, glabrous.—Trees or large shrubs. Stipules soon falling. *Pentandræ* Borr.

1. *S. pentan'dra* (L.); l. ovate-elliptic or ovate-lanceolate acuminate glandular-serrate, "*stip. ovate-oblong straight equal*," *stam.* 5 or more, caps. ovate-attenuate glabrous, stalk twice as long as the gland, style short, stig. bifid.—*E. B.* 1805.—Height 6—20 feet. Top of petioles glandular. L. fragrant.—Riversides in the North. Sh. or T. V. VI. *Bay-leaved Willow*.

E. S. I.

†2. *S. cuspidata* (Schultz ?); l. oblong lanceolate acuminate glandular serrate, *stip. half-cordate oblique*, "*stam.* 3 or 4," caps. ovate attenuate glabrous, stalk 3 or 4 times as long as the gland, style short, stig. emarginate.—*E. B. S.* 2961. *S. Meyerriana* Willd.—Height 20—30 feet. Top of the petioles glandular.—Near Shrewsbury. Sh. or T. VI. E.

Subsec. 2. *Amerina* (Dumort.). Stam. 2 or 3. Nectary of 2 pieces, one between the cal.-scale and germen, the other opposite to it.—Naturally trees.

<sup>1</sup> I have ventured to place many plants as varieties which were considered distinct by my able friend Mr. Borrer. I do so with hesitation, being very imperfectly acquainted with many of them, but follow the example of most modern botanists. It is probable that many of the lowland species are not truly natives of Britain; but evidence is wanting by which to determine their respective claims. See *Du Mortier* in *Bull. Soc. R. Bot. Belg.* i. 140; *Babington* in *J. of B.* i. 167; *Leech* in *J. of B.* ix. 360.

i. *Diandra*. Stam. 2. Catkin-scales soon falling.—*Fragiles* and *Albæ* Borr.

3. *S. fragilis* (L.); l. lanceolate pointed serrate, stip.  $\frac{1}{2}$ -cordate, caps. obovate-lanceolate stalked glabrous, stigmas rather thick bifid.—*a. S. decipiens* (Hoffm.); caps. tapering, style longer than the cloven stigmas. *E. B.* 1937. Branches smooth, highly polished, reddish brown; young shoots often crimson.—*β. S. fragilis* (L.); caps. oblong-ovate, style short, stig. bifid. *E. B.* 1807. Branches round very smooth, brown, brittle in the spring. *Crack Willow*.—*β. S. Russelliana* (Sm.); caps. stalked lanceolate-acuminate, style as long as the bifid stigmas. *E. B.* 1808. *L. viridis* Fr.? Branches polished, round, smooth. L. gradually attenuate, very glaucous beneath. Wood and bark highly valuable. *Bedford Willow*.—Damp meadows and osier-ground. T. IV. V. E. S. I.

4. *S. alba* (L.); l. elliptic lanceolate glandular-serrate acute silky on both sides when young, stip. minute, caps. nearly sessile ovate acuminate glabrous, style short, stigmas thick recurved bifid.—*E. B.* 2430.—Height 50—80 feet. Scales shorter than stam., as long as caps. in *a* and *β*, exceeding them both in *γ*. Branches silky.—*β. S. cærulea* (Sm.); l. less silky beneath. *E. B.* 2431.—*γ. S. vitellina* (Sm.); branches bright yellow, l. shorter and broader. *E. B.* 1389.—Wet places. T. V. *White Willow*. E. S. I.

ii. *Triandra*. Stam. 3. Catkin-scales persistent. L. lanceolate approaching to ovate, glabrous. Catkins lax.—Osiers, naturally trees.

\*5. *S. undulata* (Ehrh.) ; l. lanceolate much acuminate serrate glabrous except when young, stip.  $\frac{1}{2}$ -cordate acute, caps. stalked ovate acuminate, pedicel twice as long as the gland, style long, stigma bifid, scales very shaggy.—*S. lanceolata* Sm., *E. B.* 1436.—Height 12—15 feet. L. sometimes wavy, often quite silky when young. Germ. glabrous in *S. lanceolata* (or downy in the foreign *S. undulata*).—Lewes, Suss. Sh. or T. IV. V. E. I.

6. *S. triandra* (L.) ; l. oblong-lanceolate acute serrate glabrous, stip.  $\frac{1}{2}$ -cordate blunt, caps. stalked oblong-ovate glabrous, stigma nearly sessile, scales glabrous.—*E. B.* 1435.—Height 20—30 feet. Germen not furrowed. L. narrowing down to the stalk, somewhat paler beneath, or (*S. triandra* Curt.) lanceolate wavy paler and glaucous beneath.—*β. S. Hoffmanniana* (Sm.); l. somewhat rounded below ovate-lanceolate, stip. larger, scales shaggy towards their base.—*E. B. S.* 2620.—*γ. S. amygdalina* (L.); l. oblong-ovate acute rounded below, caps. ovate tumid

furrowed, young shoots furrowed. *E. B.* 1936.—Wet woods and osier-grounds. Sh. or T. IV. V. E. S. I.

Sec. 2. *CAPRISALIX* (Dumort.). Catkins lateral, sessile, without leaves or with two or three small leaves or leaflike bracts at the base; stalk sometimes elongated in fruit so as to resemble a leafy shoot, but deciduous with the catkin. Catkin-scales often discoloured at the end. Nectary simple (of one piece), on the opposite side of the stam. or germen from the catkin-scale.

Subsec. 1. *Helice* (Dumort.). Filament 1 with a 4-celled anther, or forked with 2 anthers each of 2 cells. *Anth.* purple, ultimately black. Nectary cuneate. Catkins bracteate at the base. Vernation equitant.—*Purpurea* Borr.

7. *S. purpurea* (L.); l. lanceolate broader upwards acuminate attenuate below finely serrate glabrous, caps. ovate very downy sessile, style very short, *stigm.* ovate, *anth.* 1, *stip.* 0.—*a. S. purpurea* (Sm.); decumbent, twigs purple, fertile catkins very compact. *E. B.* 1388.—*β. S. Woolgariana* (Borr.); erect, twigs yellowish grey, l. cuneate-lanceolate glaucous beneath, stigmas blunt. *E. B. S.* 2651.—*γ. S. ramulosa* (Borr.); erect, twigs pale yellowish, l. oblong-lanceolate paler beneath, stigmas sessile bifid.—*δ. S. Lambertiana* (Sm.); erect, twigs purplish glaucous, l. oblong-linear-lanceolate slightly narrowed and somewhat rounded below, stigmas ovate emarginate. *E. B.* 1359.—*ε. S. Helix* (L.); caps. oblong-ovate, *stigmas* almost linear emarginate, twigs pale yellowish or tinged with purple polished. *E. B.* 1343. *Rose Willow*.—Marshes and river-banks. T. III. IV. E. S. I.

8. *S. rubra* (Huds.); linear-lanceolate acuminate glabrous green on both sides, caps. oblong-ovate very pubescent, *style* long, *stigmas* ovate undivided, *anth.* 2 (yellow?), *filaments* combined below, *stip.* linear.—*E. B.* 1145.—Height 10—20 feet. Twigs usually tawny. L. like those of *S. viminalis* but without the white pubescence.—*β. S. Forbiana* (Sm.); l. lanceolate-oblong, style nearly as long as the linear “divided stigmas, *anth.* 1.” *E. B.* 1344. Twigs greyish yellow. L. rather paler and somewhat glaucous beneath. *Stip.* broader, variable. Stigma entire in all that I have seen.—Low meadows. T. IV. V. E. S. I.

Subsec. 2. *Vimen* (Dumort.). Stam. 2, mon- or diadelphous. *Anth.* becoming yellow. Nect. linear. Catkin-scales discoloured at the end. Catkins bracteate at the base. Stigmas not sessile. Pubescence of the l. silky.

\* *Stipules narrow.*

9. *S. viminālis* (L.); l. linear or linear-lanceolate obscurely crenate white silky and shining beneath, stip. small sublanceolate, capsule very shortly stalked lanceolate-subulate, style long, stigmas undivided.—*E. B.* 1898.—Height 10—20 feet. Branches wandlike, long, slender. Gland longer than the stalk of the caps.— $\beta$ . *intricata* (Leefe); l. broader, caps. shorter and broader, style very short, stigmas from the first cloven reflexed and entangled.— $\gamma$ . *stipularis* (Leefe); l. lanceolate, stip. linear-lanceolate denticulate or  $\frac{1}{2}$ -cordate acuminate, stigmas long.—Wet places. Sh. IV. V. Common Osier. E. S. I.

\*\* *Stipules broad.*

10. *S. stipularis* (Sm.); l. lanceolate very obscurely crenate white and downy beneath, stip.  $\frac{1}{2}$ -cordate acute, caps. ovate nearly sessile, style very short (*Sm.*) or elongate (*Hook.*), stigmas linear undivided.—*E. B.* 1214.—Height 10—20 feet with upright brittle reddish-brown twigs.—Wet places. Sh. III. E.

11. *S. Smithiana* (Willd.); l. long-lanceolate obscurely crenate white satiny beneath, stip. reniform  $\frac{1}{2}$ -cordate, caps. stalked lanceolate-subulate, pedicel about as long as the gland, style long, stigmas long linear mostly entire.—Twigs erect, somewhat downy brittle.— $\alpha$ . *S. Smithiana* (*E. B.* 1509); l. rounded at base white with satiny down beneath, stip. small narrow  $\frac{1}{2}$ -lunate.— $\beta$ . *S. rugosa* (Leefe); l. greenish white and rather silky beneath, stip.  $\frac{1}{2}$ -cordate acute, style moderate, stigmas linear broad undivided. *S. holosericea* H. & A.— $\gamma$ . *S. ferruginea* (And.); l. greenish white and rather silky beneath, stip.  $\frac{1}{2}$ -ovate or reniform, style elongate, stigmas linear-oblong undivided. *E. B. S.* 2665. Bushy.—Wet places. Sh. IV. V. E. S. I.

Subsec. 3. *Vetrix* (Dumort.). Stam. 2, free. Anth. becoming fuscous-yellow. Nectary cuneate. Catkin-scales discoloured. Vernation equitant.

i. *Caprææ*. Style short. Stipules reniform, without basal glands.

L. rugose, not turning black; pubescence crisped, not silky.—*Cinereæ* Borr.

12. *S. acuminata* (Sm.); l. lanceolate-oblong pointed finely toothed glaucous ashy and downy scarcely silky beneath, stip.  $\frac{1}{2}$ -cordate, caps. ovate tapering, style conspicuous, stigmas ovate undivided, buds downy.—*E. B.* 1434. *Loud.* 1464.—Height 25—30 feet.—Damp woods and hedges. Sh. or T. IV. E. I.

13. *S. cinerea* (L.); l. elliptic- or lanceolate-obovate pointed subserrate downy and ashy beneath, stip.  $\frac{1}{2}$ -cordate, caps. lanceolate-subulate, stigmas simple or bifid, *buds downy*.—Height 20—30 feet.—*a. S. cinerea* (Sm.); l. obovate-lanceolate rather thick reddish beneath, stip.  $\frac{1}{2}$ -cordate acute, style very short thick. *E. B.* 1897.—*β. S. aquatica* (Sm.); l. ovate-elliptic thinner downy and rather glaucous beneath, stip. reniform, style inconspicuous. *E. B.* 1437.—*γ. S. oleifolia*; l. obovate-lanceolate rather rigid downy and rather glaucous beneath, stip. small rounded. *E. B.* 1402.—Varies greatly. These varieties are scarcely distinguishable.—Wet places. T. or Sh. III. IV. *Sallow.* E. S. I.

14. *S. aurita* (L.); l. obovate repand-dentate recurved-apiculate much *wrinkled* more or less downy above pubescent beneath, stip. roundish or reniform *large stalked*, caps. lanceolate-subulate, stig. generally entire, *buds glabrous* or slightly downy.—*E. B.* 1487.—Height 3—4 feet. Edges of l. deflexed, point hooked. Stigmas and scales often reddish.—Damp woods. Sh. IV. V. E. S. I.

15. *S. cáprea* (L.); l. ovate or elliptic *flat* acute crenate-serrate wavy at the margins deep green with a downy midrib whitish above and cottony beneath, stip. subreniform, caps. lanceolate-subulate, style very short, *buds glabrous*.—*E. B.* 1488.—A small tree, 15—30 feet high. Catkins very thick, blunt. L. mostly large and broad, rarely sublanceolate; spring l. nearly entire.—*β. S. sphacelata* (Sm.); stip. often 0, l. nearly entire. *E. B.* 2333. A bush.—Woods and hedges in dryish and also subalpine places. *β.* Highland valleys. T. IV. V. *Great Sallow.* E. S. I.

ii. *Phylicifoliae*. Style long. Capsule stalked.

\* *Nigricantes*. L. punctate beneath, turning black in drying. Stip. with basal glands.

16. *S. nigricans* (Fries); young shoots and l. downy, l. ovate-elliptic or lanceolate rather glaucous more or less glaucous beneath thin, point of stip. straight.—Shoots dull-coloured.—*a.* germen and stalk silky, st. erect.—*a. S. cotinifolia* (Sm.); l. roundish elliptic or subcordate. *E. B.* 1403.—*b. S. nigricans* (Sm.); l. elliptic-lanceolate. *E. B.* 1213.—*c. S. Forsteriana* (Sm.); l. elliptic-obovate acute, stip. vaulted, catkins long. *E. B.* 2344.—*β. S. rupestris* (Donn); awlshaped germ. and stalk silky or glabrous below, st. trailing, l. elliptic-obovate. *E. B.* 2342.—*γ.* germen glabrous its stalk downy or glabrous, st. erect.—*d. S. Andersoniana* (Sm.); l. elliptic-oblong acute, caps. awlshaped not wrinkled. *E. B.* 2343.—*e. S. damascena* (Forbes)

l. broadly elliptic or subrhomboidal acute, caps. not wrinkled. *E. B. S.* 2709.—f. *S. petraea* (And.); l. oblong or sublanceolate, caps. wrinkled near the top. *E. B. S.* 2725.—? *S. hirta* (Sm.); l. elliptic-cordate pointed, "caps. very pubescent" (*Leefe*) or "glabrous" (*Arnott*) not wrinkled. *E. B.* 1404.—*S. floribunda* (Forbes); l. long elliptical glaucous beneath thinly hairy on both sides. Fem. catkins unknown. *E. B.* 2816.—The varieties are scarcely distinguishable.—Sides of streams, osier-grounds, and mountains. Sh. IV.—VI. E. S. I.

\*\* *Virentes*. L. smooth, scarcely any crisped pubescence beneath, not turning black unless gathered when very young.

17. *S. phlyicifolia* (L.); shoots and leaves soon quite glabrous dark green above and more or less shining glaucous beneath.—Shoots bright chestnut-colour. Stip. none or very small.—a. *germen and stalk silky*.—a. *S. radicans* (Sm.); l. oblong or elliptic-lanceolate, more or less decumbent. *E. B.* 1958.—b. *S. Davalliana* (Sm.); l. oblong-lanceolate pointed, style as long as stigma. *E. B. S.* 2701.—c. *S. Weigeliana* (Willd.); l. roundish or elliptic, style longer than cloven stigma. *E. B. S.* 2656.—d. *S. nitens* (And.); l. elliptic-lanceolate, style longer than undivided stigma. *E. B. S.* 2655.—e. *S. tenuior* (Borr.); l. obovate-lanceolate acuminate, style as long as stigma. *E. B. S.* 2650.—f. *S. tetrapla* (Sm.); lower part of germ. glabrous, style hairy, stigmas divided, l. lanceolate acuminate. *E. B. S.* 2702.—β. *germ. and stalk densely woolly or silky*.—g. *S. laurina* (Sm.); l. elliptic-oblong suberect, stigm. undivided. *E. B.* 1806.—γ; *lower part of germen and stalk glabrous*.—h. *S. laxiflora* (Borr.); l. broadly obovate-acuminate, style about as long as cloven stigma. *E. B. S.* 2749.—i. *S. propinqua* (Borr.); l. elliptic, style longer than the cloven stigma. *E. B. S.* 2729.—δ; *germen glabrous, stalk hairy*.—k. *S. Borreriana* (Sm.); l. lanceolate attenuate at both ends. *E. B. S.* 2619.—ε; *germen and stalk glabrous*.—l. *S. phillyreifolia* (Borr.); l. elliptic acute at both ends. *E. B. S.* 2660.—m. *S. tenuifolia* (Borr.); l. elliptic with a recurved point. *E. B. S.* 2795.—These plants are scarcely distinguishable by description alone.—By streams especially in mountain valleys. Sh. IV. V. E. S. I.

iii. *Incubaceæ*. Style short. Stipules linear.—*Fusca* Bab.  
*Rosmarinifolia* and *Repentes* Borr.

18. *S. rosmarinifolia* (L.); l. linear-lanceolate silky beneath quite entire or remotely glandular-toothed, *stip. minute lanceolate, germens silky lanceolate-acuminate scales short hairy*, "style about as long as the linear divided or entire stigmas."—*E. B.*



1365. *S. Arbuscula* Sm., *E. B.* 1366, exclusive of large leaf, not of *Koch* nor *Fr.*—A slender upright shrub, 1—3 ft. high. Catkins very short, at first drooping. Whole plant becomes nearly black in drying. Style short.—“Several parts of the North.” *Sm.* Sh. V. S. ?

19. *S. Doniána* (Sm.); *l. lanceolate* or *obovate-lanceolate* acute slightly serrate livid with scattered silky hairs beneath, *stip. linear*, germen very silky ovate-oblong longer than the bearded oblong-ovate scales, *style very short, stigmas short emarginate.*—*E. B. S.* 2599.—Shrub about 6 feet high. “Stam. monadelphous, anth. ultimately luteo-fuscous not black.” (*Koch.*)—Forfarshire. Sh. V. S.

20. *S. repens* (L.); *l. elliptic* or *elliptic-lanceolate* acute entire or minutely glandular-serrate glaucous and silky beneath, germ. subconical silky, style rather short, stigmas ovate bifid,—*a. S. repens* (Sm.); *l. elliptic-lanceolate* with a straight point, st. depressed with short upright branches. *E. B.* 183.—*β. S. fusca* (Sm.); *l. oblong-oval* straight, st. decumbent below then erect much branched. *E. B.* 1960.—*γ. S. prostrata* (Sm.); *l. elliptic-oblong* with a twisted point, st. prostrate with long straight branches. *E. B.* 1959.—*δ. S. ascendens* (Sm.); *l. elliptic* with a recurved point, st. decumbent with long somewhat ascending branches. *E. B.* 1962. *S. parvifolia* Sm., *E. B.* 1961.—*ε. S. incubacea* (L.); *l. elliptic-oblong* with a twisted point, stip stalked ovate acute. *E. B. S.* 2600.—*ζ. S. argentea* (Sm.); *l. broadly elliptic* with a twisted point, stip. stalked oval. *E. B.* 1364.—Fries thinks that *S. fusca* (L.) is different.—Heaths at various elevations. Sh. III. IV. E. S. I.

21. *S. ambig'ua* (Ehrh.); *l. oval* obovate or lanceolate slightly toothed with a recurved point somewhat rugose above soft and silky beneath, *stip. stalked half-ovate acute*, germen lanceolate-subulate silky, style very short, stigmas short at length cloven.—*E. B. S.* 2733.—*a*; *l. oval* or obovate moderately hairy.—*β. major*; *l. obovate* very silky on both sides.—*γ. spathulata*; *l. obovate* or *ovate-lanceolate* moderately hairy or silky, style somewhat long.—*δ. undulata*; *l. ovate-lanceolate.*—Gravelly heaths. Sh. V. E. S. I.

iv. *Daphnoideæ.* Style long. Capsule sessile. Catkins sessile, bracteate at the base.—*Arbusculæ* Bab.

22. *S. Arbus'cula* (L., Fries); *l. lanceolate-ovate* or *ovate glabrous* smooth glaucous and opaque beneath finely serrate, germen oblong-ovate silky, stigmas bifid.—*a. S. carinata* (Sm.);

l. ovate finely toothed minutely veined folded into a keel, catkins cylindrical with rounded hairy scales. *E. B.* 1363.— $\beta$ . *S. prunifolia* (Sm.); l. broadly ovate toothed smooth on both sides, st. erect much branched. *E. B.* 1361.— $\gamma$ . *S. venulosa* (Sm.); l. ovate toothed naked netted with prominent veins above, st. erect much branched. *E. B.* 1362.— $\delta$ . *vacciniifolia* (Sm.); l. lanceolate-ovate serrate smooth and even above silky beneath, st. decumbent. *E. B.* 2341.—Highlands. Sh. IV.—VI. S.

23. *S. Lappónum* (L.); l. lanceolate or elliptic subacuminate entire cottony or silky beneath with crisped hairs wrinkled above and when young downy, germen ovate-lanceolate silky, stigmas linear.— $\alpha$ . *S. arenaria* (L.); l. ovate-lanceolate netted and somewhat downy above veined and woolly beneath, style as long as the sessile woolly germen, stigmas linear. *E. B.* 1809. Germen with a long slender reddish style.— $\beta$ . *S. Stuartiana* (Sm.); l. ovate-lanceolate shaggy above densely silky almost cottony beneath, style as long as the almost sessile woolly germen, stigmas capillary deeply divided. *E. B.* 2586.—[ $\gamma$ . *S. glauca* (Sm.); l. ovate-lanceolate even and nearly smooth above woolly and snow-white beneath, germen sessile woolly, style at first very short with thick ovate stigmas. *E. B.* 1810. Germen blunt. Style elongating and the stigmas becoming linear and deeply cloven as the fruit ripens. This is probably not a native. Smith had it from Mr. Crowe's garden. It is hardly the same as  $\alpha$  and  $\beta$ .—*S. glauca* L., Wahl., Koch, has sub-terminal catkins with very long leafy stalks and belongs to the next section.]—Breadalbane and Clova Mountains. Sh. VI. VII. S.

[*S. acutifolia* (Willd.); l. linear-lanceolate acuminate crenate-serrate glabrous glaucous beneath, stip. lanceolate acute sub- $\frac{1}{2}$ -cordate below, "caps. ovate-conic glabrous sessile, stig. linear-oblong."—*Sy. E. B.* 1366.—Shoots purple with a glaucous bloom. Male catkins short, thick, very hairy. Fertile plant not yet found in England. *S. daphnoides*  $\beta$ . Anders.—North Yorkshire, not native. T. IV.] E.

v. *Chrysanthæ*. Style long. Stigma entire. Capsule sessile. Anth. yellow, scarcely changing colour. Catkins appearing before the leaves, sessile, terminal and lateral, with very shaggy and silky scales. Leaves broad, roundish.

24. *S. lanata* (L.); l. broadly oval pointed entire shaggy beneath, stip. oval, catkins with yellow silky hairs, germ. conical glabrous.—*E. B. S.* 2624.—A low (2 ft.) and very beautiful shrub.—Clova and Glen Lochay Mountains. Sh. V. VI. S.

Sec. 3. CHAMELYX (Fries). Catkins on long leafy persistent shoots from the terminal or subterminal buds. Stam. 2. Nectary "of 2 pieces, one between the catkin-scale and germen, the other opposite to it."

i. *Myrsinites*. Catkins at the extremity of the terminal shoot, or of those from the last but one or two of the buds, but in such a manner as to appear to be an elongation of the branch. Small bushy plants.

25. *S. Myrsinites* (L.); l. elliptic or lanceolate serrate shining often hairy with prominent veins, germen sessile ovate-subulate downy, style long.—St. much branching.—*a. S. arbutifolia* (Sm.); l. ovate or lanceolate rather acute. *S. Myrsinites*  $\beta$ . Sm.— $\beta$ . *S. Myrsinites* (Sm.); l. elliptic serrate nearly smooth, catkins short, styles short, stigmas cloven. *E. B.* 1360.—Highlands. Sh. VI. S.

26. *S. procumbens* (Forbes); l. oval minutely serrate bright green and shining on both sides, catkins long cylindrical, germen sessile ovate lanceolate downy, style short deeply cloven, stigmas bifid.—*E. B. S.* 2753.—Scales of the catkin nearly black, longer and more hairy than in *Myrsinites*. A low procumbent much-branched shrub.—Highlands. Sh. VI. S.

[Fries states that *S. retusa* (L.) was found by Mr. Winch in Scotland; but there is no such plant in his Herb.]

27. *S. Grahami* (Borr.); l. oval obscurely crenate shining glabrous silky beneath netted with veins, germ stalked ovate prolonged glabrous, style long, stigma bifid, young branches silky.—*J. of B.* v. t. 66. *Sy. E. B.* 1377.—St. long, trailing. Allied to *S. Myrsinites* and *S. retusa* (can it be the plant referred to by Fries?).—Frouvyn, Sutherl. *Graham*. Muckish, Doneg. *D. Moore, J. of B.* ix. 300. Sh. VI. ? S. I.

ii. *Reticulatae*. Catkins opposite to the terminal leaves, with a bud between them.

28. *S. reticulata* (L.); l. nearly roundly oval very blunt entire netted with veins and glaucous beneath, germen sessile oblong-ovate downy, style short, stigmas bifid.—*E. B.* 1908.—A procumbent much branched shrub. St. usually buried. Catkins on long stalks.—Lofty mountains. Sh. VI. S.

29. *S. Sadleri* (Syme); l. roundly ovate or subcordate entire smooth and cottony netted and glabrous beneath, germen glabrous with long woolly stalks, style long, stigmas linear bifid.—*Edin. Bot. Trans.* xii. 208. t. 1. *J. of B.* xiii. t. 158.—A small

almost subterranean prostrate shrub.—I have not seen this plant, and it may be misplaced here.—Glen Callater, very rare. Sh. VIII.?

S.

iii. *Herbaceæ*. Catkins exactly terminal.

30. *S. herbacea* (L.); l. round blunt or retuse serrate shining glabrous netted with veins, *germens* subsessile ovate-conical glabrous, style short, stigmas bifid.—*E. B.* 1907.—A very minute herblike shrub; the st. extend far amongst loose stones on the tops of mountains. Edges and veins of l. hairy.—Alpine situations. Sh. VI. E. S. I.

*Index to the Salices.*

- |                           |                          |                      |
|---------------------------|--------------------------|----------------------|
| <i>S. acuminata</i> , 12. | <i>glauca</i> , 23.      | <i>purpurea</i> , 7. |
| <i>acutifolia</i> , 23*.  | Grahami, 27.             | radicans, 17.        |
| <i>alba</i> , 4.          | Helix, 7.                | ramulosa, 7.         |
| <i>ambigua</i> , 21.      | <i>herbacea</i> , 30.    | repens, 20.          |
| amygdalina, 6.            | hirta, 16.               | reticulata, 28.      |
| Andersoniana, 16.         | Hoffmanniana, 6.         | retusa, 26*, 27.     |
| aquatica, 13.             | <i>holosericea</i> , 11. | rosmarinifolia, 18.  |
| <i>Arbuscula</i> , 18.    | incubacea, 20.           | rubra, 8.            |
| <i>Arbuscula</i> , 22.    | intricata, 9.            | rugosa, 11.          |
| arbutifolia, 25.          | Lambertiana, 7.          | rupestris, 16.       |
| arenaria, 23.             | lanata, 24.              | Russelliana, 3.      |
| argentea, 20.             | <i>lanceolata</i> , 5.   | Sadleri, 29.         |
| ascendens, 20.            | Lapponum, 23.            | Smithiana, 11.       |
| <i>aurita</i> , 14.       | laurina, 17.             | sphacelata, 15.      |
| Borreriana, 17.           | laxiflora, 17.           | stipularis, 9.       |
| cærulea, 4.               | <i>Meyeriana</i> , 2.    | stipularis, 10.      |
| caprea, 15.               | Myrsinites, 25.          | Stuartiana, 23.      |
| carinata, 22.             | nigricans, 16.           | tenuifolia, 17.      |
| cinerea, 13.              | nitens, 17.              | tenuior, 17.         |
| cotinifolia, 16.          | oleifolia, 13.           | tetrapla, 17.        |
| cuspidata, 2.             | <i>parvifolia</i> , 20.  | triandra, 6.         |
| damascena, 16.            | pentandra, 1.            | undulata, 5.         |
| Davalliana, 17.           | petræa, 16.              | vacciniifolia, 22.   |
| decipiens, 3.             | phillyreifolia, 17.      | venulosa, 22.        |
| Doniana, 19.              | phylicifolia, 17.        | viminalis, 9.        |
| ferruginea, 11.           | procumbens, 26.          | <i>viridis</i> , 3.  |
| Forbiana, 8.              | propinqua, 17.           | vitellina, 4.        |
| Forsteriana, 16.          | prostrata, 20.           | Weigeliana, 17.      |
| fragilis, 3.              | prunifolia, 22.          | Woolgariana, 1.      |
| fusca, 20.                |                          |                      |

1. POP'ULUS *Linn.* Poplar.

1. *P. alba* (L.); l.-buds downy, l. roundish-cordate angularly toothed cottony and snowy white beneath, l. of the young shoots

cordate palmately 5-lobed, *stig.* 2 *linear bifid crosslike* (yellow).—*E. B.* 1618.—With suckers. L. generally lobed. Male catkin-scales hairy.—Damp woods. T. III. IV. *White Poplar*.  
*Abele.* E. S.

2. *P. canes'cens* (Sm.); l.-buds downy not viscous, l. roundish obscurely lobed sparingly *cottony and grey beneath*, l. of young shoots cordate-ovate undivided, *stig.* 2 *wedgedshaped 3—4-lobed* (purple).—*E. B.* 1619.—With suckers. L. not lobed, except occasionally the youngest. Male catkin-scales hairy.—Damp woods. T. III. IV. *Gray Poplar.* E.

3. *P. trem'ula* (L.); l.-buds glabrous slightly viscous, l. nearly round acute serrate *glabrous on both sides*, young l. slightly downy, *stig.* 2 *bifid erect*.—*E. B.* 1909.—A small tree, with suckers. Petioles laterally compressed. Male catkin-scales hairy, palmately cut.—Woods. T. IV. *Aspen.* E. S. I.

4. *P. nigra* (L.); l.-buds glabrous viscous, l. triangular-rhomboidal acuminate serrate *glabrous on both sides*, catkins lax cylindrical, *stig.* 2 roundish 2-lobed.—*E. B.* 1910.—A large tree, without suckers. Petioles laterally compressed. Young shoots glabrous. Male catkin-scales glabrous.—Damp places and river-banks. T. III. *Black Poplar.* E.

## Tribe II. *Myriceæ*.

### 3. MYRI'CA *Linn.* Sweet Gale.

1. *M. Gale* (L.); l. lanceolate broader upwards serrate, st. shrubby.—*E. B.* 562.—Height 3—4 feet. Bushy. Catkins sessile erect. Fr. with resinous glands. L. fragrant when bruised.—Bogs. Sh. V. *Bog Myrtle.* E. S. I.

## Tribe III. *Betulineæ*.

### 4. BET'ULA *Linn.* Birch.

1. *B. verruc'osa* (Ehrh.); l. rhomboid-triangular doubly serrate abruptly acuminate, scales of the fem. catkins 3-lobed, *laterallobes falcate-reflexed*, fr. obovate.—*Fl. Dan.* 2549. *Sy. E. B.* 1295. *B. alba* Koch. *B. odorata* R. xii. 626.—L. usually glabrous often covered with resinous spots above, always having a manifest tendency to a rhomboidal form, flat beneath with raised veins above. Young shoots mostly with resinous tubercles, often very long and pendulous. Stip. ovate-lanceolate, acute, thrice as long as broad, circinate; sides not deflexed. Buds conical. The catkin-

scales and the shape of the l. distinguish this from the next.—  
Rather common. T. IV. V. *White Birch*. E. S. I.

2. *B. glutinosa* (Fries!); l. rhomboid-ovate or cordate unequally serrate acute, scales of the fem. catkins 3-lobed, *lateral lobes ascending*, fr. broadly obovate.—*Sy. E. B.* 1296. *B. alba* R. xii. 623. *B. pubescens* Koch.—L. usually glabrous, always more or less ovate, flat above with raised veins beneath. Stip. ovate, blunt, twice as long as broad; sides deflexed. Buds ovoid. Not so elegant a tree as the preceding and often little more than a bush. Twigs sometimes pendulous.— $\beta$ . *B. pubescens* (Ehrh.); l. peduncles and young twigs downy.—Common. T. IV. V. *Common Birch*. E. S. I.

3. *B. nana* (L.); l. *roundish crenate* glabrous blunt, scales of the fem. catkin digitate-trifid, lobes equal, fr. orbicular with a very narrow membranous margin.—*E. B.* 2326.—A small procumbent shrub. L. minute. Catkins sessile, small.—Turfy places in the Highlands. Sh. V. *Dwarf Birch*. S.

#### 5. AL'NUS Tourn. Alder.

1. *A. glutinosa* (Gaert.); l. roundish blunt wavy serrate glutinous rather abrupt with a wedged-shaped base, axils of the veins beneath downy.—*E. B.* 1508. *R.* xii. 641. *St.* 29. 15.—A moderately large tree. Trunk and branches crooked. Male catkins long and pendent; fem. ones short, ovoid or oblong, very persistent.— $\beta$ . *incisa*; leaves deeply cut.—Wet places and river-banks.  $\beta$ . Wigtonshire. *Dr. Balfour*. T. III. E. S. I.

#### Tribe IV. Cupuliferæ.

#### 6. FA'GUS Linn. Beech.

1. *F. sylvatica* (L.); l. ovate glabrous obscurely dentate ciliate on the edges.—*E. B.* 1846. *R.* xii. 639.—A large tree with triquetrous fruit.—Woods, particularly on calcareous soils. T. III. IV. E.

#### 7. CASTA'NEA Tourn. Chestnut.

[*C. vulgaris* (Lam.); l. oblong-lanceolate acuminate mucronate-serrate glabrous on each side.—*Fagus* Sm., *E. B.* 886. *R.* xii. 640.—Height 50—80 feet. A magnificent tree.—A very doubtful native. T. V. *Sweet Chestnut*.] E.

## 8. QUER'CUS Linn. Oak.

1. *Q. Robur* (L.); l. deciduous stalked obovate-oblong sinuate, lobes blunt, inv. much shorter than the ripe acorn its scales adpressed.—*a. Q. pedunculata* (Ehrh.); young branches glabrous, petioles short, fr.-catkins long-stalked, fr. scattered. *E. B.* 1342.—*β. Q. intermedia* (D. Don); young branches glabrous, petioles short, l. stellate-downy beneath, fr.-catkins shortly stalked, fr. near together. *Mart. Rust.* 11.—*γ. Q. sessiflora* (Salisb.); young branches downy, petioles long, l. glabrous beneath, fr.-catkins subsessile, fr. near together. *E. B.* 1845.—It is generally supposed by foresters that there are two if not three species of Oak in Britain. I have failed in learning how to distinguish them.—Woods. T. IV. V. E. S. I.

## 9. COR'YLUS Linn. Hazel.

1. *C. Avellana* (L.); stip. oblong blunt, l. roundish-cordate acuminate, involucre of the ovoid fr. bellshaped spreading torn at the margin.—*E. B.* 723.—A shrubby tree. Young twigs hairy and glandular. L. downy beneath. Male catkins long, pendulous. Fem. fl. in ovoid buds. Stigmas bright crimson.—Hedges and copses. Sh: III. IV. *Hazel Nut.* E. S. I.

## 10. CARPI'NUS Linn. Hornbeam.

1. *C. Bet'ulus* (L.); scales of the fruit 3-parted, segments lanceolate, the middle one longest.—*E. B.* 2032.—A small tree. L. ovate, acute, plaited when young, deeply and sharply doubly serrate.—Damp clayey woods and hedges. T. V. E.

## Division V. GYMNOSPERMÆ.

Ovules and seeds apparently naked. Carpel spread out flat. No calyx, no corolla.

## Order LXXIX. CONIFERÆ.

Fl. monœcious or dicecious.—Barren fl. of one or more monadelphous stamens, in a deciduous catkin, about a common axis. Anth. of 2 or more lobes bursting outwards, often ending in a scalelike crest.—Fertile fl. usually in cones, sometimes solitary. Ovary spread open in the shape of a scale and placed in the axil of a membranous bract; in the solitary fl. apparently wanting. Ovules naked in pairs on the face of the ovary and inverted; or

(in the solitary fl.) erect. [Or, as some think: ovaries in pairs (or several), inverted, on a scale (which becomes the cone-scale) situated in the axil of a bract; each of 2 connate carpels having together the form of a horseshoe or ring and ultimately producing a wing; ovule between the carpels, its tips exposed; or erect, the 2 carpels uniting in the form of a cup, without a carpellary scale.] Fr. a cone, or solitary seed. Testa hard, crustaceous. Embryo in the axis of fleshy albumen. Radicle next the apex.—Woody tissue marked with circular disks; without medullary rays or proper vascular tissue.

Tribe I. *TAXINEÆ*. Male fl. in catkins. Fem. fl. solitary, naked or bracteate, erect; no carpellary scale.

1. *TAXUS*. Catkins of male fl. oval, scaly below, flowering at the top. Stam. many. Anth. peltate, 3—8-celled. Fem. fl. scaly below. Style 0. Ovule surrounded at the base by a ring which becomes a fleshy cupshaped disk surrounding the seed.

Tr. II. *CUPRESSINEÆ*. Male fl. in catkins. Anth. 4—7, inserted on the edge of the subpeltate scales. Fem. fl. few, in a small catkin, erect; no carpellary scale.

2. *JUNIPERUS*. Anth. 4—7, 1-celled, inserted on the lower edge of the scales. Scales of the fem. catkin imbricate, lower ones barren. Ovules 3, surrounded by a 3-fid fleshy involucre formed of the 3 uppermost connate scales of the catkin.

Tr. III. *ABIETINEÆ*. Fl. in catkins. Anth. 2, 1-celled, adnate to the underside of the scales. Fem. fl. a scalelike open ovary in the axil of a membranous scale, bearing two naked ovules pointing towards the axis [or, perhaps, fem. fl. in pairs on an axillary scale.] Fr. winged.

3. *PINUS*. Male catkins crowded, racemose. Scales of the cone thickened and angular at the end. Fr. with a crustaceous coat.

### Tribe I. *Taxineæ*.

#### 1. *TAX'US* Linn. Yew.

1. *T. baccata* (L.); 1. 2-ranked crowded linear acute, fl. axillary sessile.—*E. B.* 746.—A low tree, trunk often attaining a very considerable bulk. Fr. roundish.—*T. fastigiata* (Lindl.) is not even a permanent variety.—Mountainous woods and limestone cliffs. T. III. IV. *Yew*. E. S. I.



Tribe II. *Cupressineæ*.2. JUNIP'ERUS *Linn.* Juniper.

1. *J. communis* (L.); l. 3 in each whorl spreading linear subulate mucronate keeled exceeding the ripe fruit.—*E. B.* 1100.—Fruticose, erect. L. with a broad flat shallow channel above, the keel beneath with a slender furrow. Berries black, tinged with blue, about half the length of the leaves.—Dry hills, especially on a calcareous soil. Sh. V. E. S. I.

2. *J. nana* (Willd.); l. 3 in each whorl incurved linear lanceolate mucronate keeled equalling the ripe fruit.—*E. B. S.* 2743.—A prostrate shrub with longer berries and shorter leaves than the last.—Mountains. Sh. V. E. S. I.

Tribe III. *Abietineæ*.3. PI'NUS *Linn.* Scotch Fir.

1. *P. sylves'tris* (L.); l. in pairs, young cones stalked recurved ovoid-conical, wing thrice as long as the seed.—*E. B.* 2460.—A lofty tree. Cones of this species have been found at considerable depths in the Irish bogs and English fens.—Highlands. T. V. VI. S.

## Class II. MONOCOTYLEDONES.

Stems destitute of medullary rays, consisting of cellular tissue amongst which the vascular tissue is mixed in bundles, increasing by the addition of new matter within. Leaves mostly alternate and sheathing, with parallel simple veins connected by smaller transverse ones, rarely net-veined. Cotyledon one, or if more they are alternate.

## Division I. DICTYOGENÆ.

Leaves net-veined. Floral envelopes whorled.

## Order LXXX. TRILLIACEÆ.

Perianth inferior, 6- or 8-parted; in 2 whorls; outer whorl or

calyx herbaceous; inner or corolla coloured, or in our plant herbaceous. Stam. 6—10. Anth. linear; filaments subulate. Ovary superior, 3—5-celled. Ovules indefinite in number, in two rows in each cell. Styles as many as the cells, distinct; stigmas inconspicuous. Fr. succulent, not bursting. Seeds with a leathery skin. Embryo minute, in fleshy albumen.—Raphidiferous. Often combined with *Liliaceæ*.

1. PARIS. Perianth subherbaceous, 8-parted; 4 inner divisions narrower than the others. Stam. 8—10. Anth. on the middle of the subulate filament.

1. PAR'IS *Linn.* Herb Paris.

1. *P. quadrifolia* (L.); 1. usually 4 in a whorl.—*E. B.* 7.—St. 1 ft. high, from the end of a long rhizome, usually with 4, rarely from 3—6, ovate l. at its summit. Fl. solitary, terminal. Sep. lanceolate. Pet. subulate. Styles 4. Berry 4-celled; cells with 4—8 seeds. No root-leaves.—Damp woods. P. V. E. S.

Order LXXXI. DIOSCOREACEÆ.

Perianth superior, petal-like, 6-parted. Stam. 6, inserted on the base of the segments of the perianth. Anth. bursting inwards. Ovary inferior, 3-celled. Ovules 2 in each cell, erect. Style 1. Stigmas 3, reflexed. Fr. baccate, or dry and flat. Embryo minute, quite enclosed in the albumen.

1. TAMUS. Perianth bellshaped; limb 5-parted. Male with 6 stamens. Fem. with the perianth adhering to the ovary and persistent, and very short abortive stamens.

1. TA'MUS *Linn.* Black Bryony.

1. *T. communis* (L.); 1. undivided cordate acute shining.—*E. B.* 91.—Dioecious. Root large, thick, fleshy. St. very long, twining. Racemes axillary, on long stalks. Fl. yellowish-green, regular, small. Berry red.—Hedges and thickets. P. V. VI. E.

Division II. FLORIDÆ.

(Orders LXXXII.—XCVI.)

Leaves parallel-veined, persistent. Floral envelopes whorled, or none.—There is no proper perianth distinguishable in Ord. XCII.—XCVI. in our plants.

## Order LXXXII. HYDROCHARIDACEÆ.

Sep. 3, herbaceous. Pet. 3, regular, coloured. Stam. epigynous. Ovary solitary, inferior; placentas parietal sometimes projecting into the centre of the ovary. Stig. 3—6. Fr. dry, or succulent, not bursting, 1- or spuriously many-celled. Seeds many. Albumen 0. Embryo straight, cylindrical.—No raphides.

1. HYDROCHARIS. Dioecious. Cal. 3-parted. Pet. 3. Male with 9—12 stamens in 3 rows surrounding 3 abortive styles.—Fem. with 3 abortive filaments and 3 fleshy scales surrounding the 6 deeply bifid styles. Caps. 6-celled, many-seeded.—L. floating, orbicular.
2. STRATIOTES. Dioecious. Cal. 3-parted. Pet. 3. Male with 12 or more stamens surrounded by many abortive ones. Fem. with 6 deeply bifid styles. Berry 6-celled, many-seeded.—L. swordshaped.
3. ANACHARIS. Dioecious. Fl. from tubular bifid spath. Cal. 3-parted. Pet. 3. "Male. Sep. ovate-oblong. Pet. linear or none. Stam. 9; filaments combined into a column below."—Fem. with a long filiform tube, 3 abortive filaments and ligulate stigmas. Caps. 1-celled, few-seeded.—L. oval-oblong, in whorls.

## 1. HYDROCHARIS Linn. Frog-bit.

1. *H. Morsus-ranæ* (L.).—*E. B.* 808. *R.* vii. 62.—Floating. With runners bearing tufts of stalked roundish-reniform entire leaves. Fl. white, delicate, springing from a pellucid membranous sheath. Seeds covered with beautiful prominent spirally twisted cells.—Ponds and ditches. P. VII. VIII. E. I.

## 2. STRATIO'TES Linn. Water-Soldier.

1. *S. aloides* (L.); l. swordshaped-triangular ciliate-spinous.—*E. B.* 379. *R.* vii. 61.—Creeping in the mud. L. many, rigid. Stalk compressed, 5 or 6 in. high, with two l. near its summit. Fl. white, delicate. The plant rises to the surface of the water to flower, and sinks again afterwards.—Fen-ditches in the East of England; naturalized in Scotland and Ireland. P. VII. E. [S. I.]

## 3. ANACH'ARIS Rich. Water-Thyme.

\*1. *A. Alsinas'trum* (Bab.); l. 3 in a whorl oval-oblong blunt serrulate, fem. fl. from a tubular bifid spath many times exceeding the sessile germen, sep. and pet. broad nearly equal, stigmas reflexed.—*A. N. H.* ser. 2. i. t. 8. *E. B. S.* 2993. *Elodea* Rich. ?—st. long, branching; whorls of leaves many,

close together. Fl. very small but with a very long slender tube often 2 or 3 in. long. The spathe in the axil of a leaflike bract placed within the whorl of leaves. Sep. tinged with green and pink externally, incurved, hooded. Pet. flat, transparent, recurved, oblong. Filaments at first curved outwards, their points placed under the hood of the sepals, afterwards erect, linear, blunt. Anth. 0. Stigmas recurved, linear, or deeply bifid. Sep. pet. and stigmas of about equal length. Style adnate on 3 sides to the tube.—In water. Probably introduced from America. Canals and rivers. P. VII.—IX. E. S. I.

### Order LXXXIII. ORCHIDACEÆ.

Sep. 3, usually coloured. Pet. 3, 2 above, 1 below (lip) frequently lobed and spurred and unlike the others. Stam. 3, united in a central column, 2 lateral abortive, or (in *Cypripedium*) the middle one abortive. Pollen powdery or cohering in masses. Ovary 1-celled, inferior, with 3 parietal placentas. Style forming part of the column with the stamens; stigma a viscid space in front of the column. Caps. 3-valved. Seeds very many, minute. Testa loose, netted.—Raphidiferous.

Tribe I. *OPHRYPDINEÆ*. Anther wholly adnate to the column. Pollen-masses in divisible lobes which are indefinite in number and waxy, stalked.—Root with 2 fleshy entire or palmate knobs below the fibres.

\* *Anther-cells with a prolongation of the stigma between their bases.*

1. *ORCHIS*. Perianth ringent, hooded. Lip 3-lobed, spurred. Glands of the stalks of the pollen-masses in a common pouch.
2. *GYMNADENIA*. Glands of the pollen-masses without a pouch. Otherwise like *Orchis*.
3. *ACERAS*. Lip without a spur. Otherwise like *Orchis*.

\*\* *A plate in front of the glands of the pollen-masses.*

4. *NEOTINEA*. Perianth ringent, hooded. Lip 3-lobed, spurred. Glands of the pollen-masses in separate hollows behind the broad recurved rostellum, which is bounded laterally by 2 semicylindrical ascending processes diverging from its base.

\*\*\* *No prolongation of the stigma between the anther-cells, nor plate in front of them.*

5. *HABENARIA*. Perianth ringent, hooded. Lip 3-lobed or entire, spurred. Glands of pollen-masses naked.

6. **OPHRYS.** Perianth patent. Lip variously lobed, without a spur. Glands of pollen-masses each in a distinct pouch.
7. **HERMINIUM.** Perianth bellshaped, segments all erect. Lip 3-lobed, tumid beneath at the base, without a spur. Glands of the stalks of the pollen masses exserted, naked.

Tr. II. **NEOTTIDEÆ.** Anther attached by its base, persistent. Pollen-masses granular; granules only in a slight state of cohesion; no stalks.—Root of many fleshy fibres, rarely of 2—4 much-thickened ones.

\* *Stigma with a prolongation (rostellum) at its top.*

8. **GOODYERA.** Perianth ringent. Lip gibbous at the base, entire, included. Stigma subcordate. Rostellum erect, bipartite, with a large squarish appendage between its slender segments.
9. **SPIRANTHES.** Perianth ringent. Lip channelled, clawed, fringed. Stigma roundish. Rostellum straight, at length bifid, with a long linear appendage between its points.
10. **LISTERA.** Perianth ringent. Lip deflexed, 2-lobed. Stigma transverse. Rostellum foliaceous, acute, arching over stigma. Column very short.
11. **NEOTTIA.** Perianth hooded. Lip deflexed, 2-lobed, gibbous at the base. Stigma transverse. Rostellum flat, broad, prominent, entire, without an appendage. Column long.
12. **EPIPACTIS.** Perianth patent. Lip interrupted; the basal division concave; terminal one (label) larger with 2 projecting plates at its base above. Stigma nearly square. Rostellum short, terminated by a globose appendage. Anth. terminal, erect, sessile, 2-celled. Column short. Germen straight on a twisted stalk.

\*\* *Stigma without a rostellum.*

13. **CEPHALANTHERA.** Perianth converging (in *C. rubra* spreading). Lip interrupted, the basal division gibbous, jointed to the recurved label. Stigma transverse. Anth. terminal, erect, movable, shortly and thickly stalked, 2-celled. Column long. Germen sessile, twisted.

- Tr. III. *ARETHUSEÆ*. Anther ultimately free, like a lid, deciduous. Pollen in many granules, pulpy or powdery, cohering in 2 stalked masses.—Root (in our plant) of fleshy much-branched fibres.
14. *EPIPOGUM*. Perianth patent. Lip posterior, erect, large, entire, with a small patent lobe on each side of its base and an erect inflated spur. Anth. tumid, seated in the lobed top of the column. Stigma transverse. Germen and stalk not twisted.
- Tr. IV. *MALAXIDEÆ*. Anther terminal, free, usually like a lid. Pollen cohering in a definite number of granules and at length waxy and confluent; no stalks.—Root fibrous, with or without a superior bulb.
15. *CORALLORRHIZA*. Perianth converging. Lip with 2 prominent longitudinal ridges at the base, 3-lobed; lateral lobes small; middle lobe large, slightly emarginate. Spur short or wanting. Stigma triangular. Rostellum wanting; but a large globose appendage. Anth. terminal, 2-celled, opening transversely. Column long. Germen slightly stalked, straight.
16. *MALAXIS*. Perianth patent. Lip posterior, erect, entire, similar to the pet., smaller than the sepals. Spur 0. Stigma rhomboidal. Rostellum short, entire, acute. Anth. terminal, continuous with the short column, out of the apex of which it appears as if it were excavated, with 2 imperfect cells. Pollen-masses connected at their apex. Germen upon a twisted stalk.
17. *STURMIA*. Perianth patent. Lip anterior, erect or oblique, entire, dilated, much larger than the sepals. Spur 0. Stigma roundish. Rostellum wanting; but an appendage consisting of 2 tubercles. Anth. terminal, deciduous, movable like a lid, with 2 distinct cells. Column long. Germen on a twisted stalk.
- Tr. V. *CYPRIPEDIEÆ*. Intermediate anther barren petal-like, 2 lateral anthers perfect.
18. *CYPRIPEDIUM*. Perianth patent. Lip inflated. Column trifid above; the lateral lobes bearing stamens; middle lobe sterile, dilated. Two lower (lateral) sepals combined. Germen straight.

Tribe I. *Ophrydineæ*.1. OR'CHIS *Linn.* Orchis.

\* *Glands of the pollen-masses separate, lip erect in the bud.*

† Bracts mostly 1-veined, root-knobs undivided.

‡ *Lip 3-lobed, lobes broad and short.*

1. *O. Mório* (L.); lip crenulate, middle lobe truncate-emarginate, spur ascending subclavate rather shorter than the germen, *sep. and pet. blunt converging* to form a hood, anth. obovate rather acute.—*E. B.* 2059. *R.* xiii. 363.—St. 6—12 in. high. L. lanceolate, lower spreading, upper adpressed. Fl. few, in a lax spike, purple; sep. (rarely patent) and pet. marked with green veins, converging so as to form a sort of hood; lip pale in the middle, spotted with purple. Fl. sometimes white.—Meadows and pastures. P. V. VI. *Green-winged Orchis.*

E. I.

2. *O. mas'cula* (L.); lip crenate, middle lobe emarginate, spur ascending rather longer than the germen, *sep. acute 2 outer reflexed upwards*, pet. converging, anth. obcordate apiculate.—*E. B.* 631. *E. B. S.* 2995. *R.* xiii. 390, 391.—St. a foot high. L. mostly radical, elliptic-lanceolate, usually spotted with purple. Lowest bract often 3-veined. Fl. in a lax spike, purple; centre of the lip whitish at the base, spotted and downy. Sep. and pet. without green veins.—Woods and pastures. P. V. *Early purple Orchis.*

E. S. I.

‡‡ *Lip pinnately 4-lobed (that is, middle lobe bifid and often with an intermediate tooth).*

3. *O. purpúrea* (Huds.); basal lobes of cor.-lip linear-oblong, *interm. lobe widening gradually upwards* and 2-lobed with an interm. tooth, its lobes denticulate at the tip and broad, spur about  $\frac{1}{2}$  as long as the germen, sep. converging into an *ovate hood* including the pet., bracts minute.—*R.* xiii. 378. *O. fusca* Jacq. *O. militaris* Sm., *E. B.* 16.—St. 1—2 ft. high. L. ovate-oblong, blunt. Fl. in a rather dense spike; *helmet dark purple*, variegated; lip paler with raised rough red points, its term. lobe very broad.—Chalky bushy hills in Kent. P. V. E.

4. *O. militaris* (L.); basal lobes of cor.-lip linear, *interm. lobe broader linear suddenly widened* and 2-lobed with an interm. tooth its lobes mostly entire at the tip and broad, spur about  $\frac{1}{2}$  as long as the germen, sep. converging into an *ovate-lanceolate*

hood including the pet., bracts minute.—*E. B. S.* 2675. *R.* xiii. 376.—Smaller than the preceding. *Helmet ash-coloured or pale purple.* Lip purple, white in the middle with raised rough red spots, with a linear space between the pairs of lobes, and its term. lobes broad.—Chalky hills. Berks., Oxf., Bucks, Herts. P. V. E.

5. *O. Sim'ia* (Lam.); lobes of lip all long narrow 1-veined linear with an interm. setaceous tooth, spur  $\frac{1}{2}$  as long as the germen, sep. converging into an ovate-lanceolate hood including the pet., bracts minute.—*O. tephrosanthos* *E. B.* 1873. *R.* xiii. 373.—More slender than the preceding. *Helmet dark purplish.* Lip with remarkably slender segments, dark purple, with or without small rough raised points; a linear space between the pairs of lobes which are about equal in size.—Chalky hills in Berks., Oxf., and Kent. P. V. E.

6. *O. ustulata* (L.); lobes of lip linear-oblong, spur  $\frac{1}{2}$  the length of the germen, sep. converging into a roundish hood including the pet., bracts long.—*E. B.* 18. *R.* xiii. 368.—St. 4—6 in. high. L. lanceolate, acute. Spike oblong, dense, dark at top, nearly white below. Fl. small, many. *Helmet dark purple.* Pet. linear-lanceolate, blunt. Lip white with purple spots.—Calcareous hills. P. VI. E.

†† Bracts with 3 or more veins, root-knobs undivided.

[*O. laxiflora* (Lam.); lip 3-lobed, lateral lobes rounded and crenulate in front longer than the truncate slightly emarginate interm. lobe, spur shorter than the germen cylindrical, 2 lateral sep. reflexed upwards, pet. converging, anth. obcordate apiculate.—*E. B. S.* 2828. *R.* xiii. 393.—St. 1—2 feet high, round, angular and rough upwards. L. lanceolate or linear-lanceolate. Bracts 3—5-veined. Spike long, lax. Fl. bright purple. Some specimens approach *O. palustris*.—Wet meadows and bogs in Jersey and Guernsey. P. V. VI.]

††† Bracts with 3 or more veins, root-knobs palmate.

7. *O. maculata* (L.); lip 3-lobed flat crenate, spur subulate shorter than the germen, three sep. patent, pet. converging, st. solid, l. lanceolate-obovate.—*E. B.* 632. *R.* xiii. 407.—St. about a foot high. L. usually spotted with purple; lower blunt or rarely acute, broadest towards their top; upper linear-lanceolate, resembling the bracts. Spike ovate, afterwards elongate; lower bracts exceeding the germen, upper equalling it. Fl. pale purple, more or less streaked with purple. Lateral veins of bracts in-



conspicuous. Lip usually flat, deeply 3-lobed, lateral lobes rounded, middle lobe longer and narrower.—Damp woods and pastures. P. V. VI. *Spotted Hand-Orchis*. E. S. I.

8. *O. latifolia* (L.); lip obscurely 3-lobed its sides ultimately reflexed, crenate, spur subulate shorter than the germen, two lateral sep. patent, middle sep. and pet. converging, st. hollow, l. lanceolate acute.—*E. B. S.* 2973. *R.* xiii. 402.—About a foot high. L. often only faintly spotted, spreading, lowest oblong and blunt, from a narrow base but broadest about their middle. Spots on l. ringshaped. Lower bracts exceeding the flowers. Lip usually only slightly lobed. Mostly quite past flowering when Sp. 9 is in perfection in the middle of June.—Marshes and damp meadows. P. V. VI. *Marsh-Orchis*. E. S. I.

9. *O. incarnata* (L.); lip obscurely 3-lobed its sides reflexed crenate, spur subulate shorter than the germen, two lateral sep. patent, middle sep. and pet. converging, st. hollow, l. narrowed from a broad base hooded at the end.—*R.* xiii. 397. *O. latifolia* *E. B.* 2308. *Curt.* ii. 184.—L. erect and approaching the stem. Bracts all usually exceeding the flowers.— $\beta$ . *O. angustifolia* (R.); l. erect-patent linear-lanceolate, upper l. erect, lower bracts equalling the flowers upper ones shorter. *R.* xiii. 394. *O. Traunsteineri* Koch.—Marshes. P. VI. E. S. I.

\*\* Glands of the pollen-masses united, root-knobs undivided.

† Spur filiform. ANACAMPTIS Rich.

10. *O. pyramidalis* (L.); lip with 3 equal lobes and 2 tubercles at the base above, lobes oblong truncate, middle lobe sometimes emarginate, spur filiform longer than the germen, lateral sep. ovate-lanceolate acute spreading, bracts 3-veined.—*E. B.* 110.—St. 12—18 in. high. L. linear-lanceolate, acute. Spike pyramidal, afterwards cylindrical. Fl. rose-purple, sometimes white.—Calcareous pastures. P. VII. *Pyramidal Orchis*. E. S. I.

†† Spur conical. HIMANTOGLOSSUM Spr.

11. *O. hircina* (Scop.); lip 3-parted spiral in the bud, lobes linear, middle one very long twisted, latter much shorter wavy, spur very short.—*E. B.* 34. *R.* xiii. 359, 360.—St. 2—3 feet high. "Cal. green, spotted with dull purple internally." Lip purplish white and spotted at the base, an inch or more long.—Bushy chalk hills, very rare. Kent. Surrey. Gt. Glenham, Suff. † P. V. *Lizard Orchis*. E.

## 2. GYMNADE'NIA R. Br.

1. *G. conops'ea* (R. Br.); lip 3-lobed, lobes equal entire blunt, lateral sep. spreading, spur filiform twice as long as the germen, root-knobs palmate.—*E. B.* 10.—St. a foot high. L. linear-lanceolate. Spike cylindrical, elongated. Bracts 3-veined. Fl. rose-purple, fragrant. Pollen-cells open in front and below, stopped below by oblong glutinous valves quite distinct from the stigma, and to the broader ends of which the glands of the pollen-masses are attached.—Hilly pastures. P. VI. VII. E. S. I.

2. *G. al'bida* (Rich.); lip 3-lobed, lobes unequal entire, middle lobe longest and broadest, sep. and lateral pet. converging, spur much shorter than the germen, root-knobs clustered.—*Habenaria* R. Br. *E. B.* 505. *Peristylus* Lindl.—St. 6—12 in. high. L. oblong, blunt; upper lanceolate, acute. Spike elongated, cylindrical, dense. Bracts 3-veined. Fl. small, yellowish white, fragrant.—Mountain-pastures. P. VI. VII. E. S. I.

## 3. A'CERAS R. Br.

1. *A. anthropoph'ora* (R. Br.); lip 3-parted, segments linear-filiform, middle one bifid and often with an intermediate tooth.—*E. B.* 29. *R.* xiii. 357.—Root-knobs ovate. Height 8—12 in. Spike long, lax. Fl. greenish yellow. Sep. ovate, acute, converging, margined with purple, including the linear-lanceolate blunt petals.—Dry chalky places. P. VI. *Green-man Orchis*. E.

## 4. NEOTIN'EA Reichenb.

1. *N. intac'ta* (Reich.); lip 3-lobed, lobes unequal, lateral lobes linear acute falling much short of the broad oblong interm. lobe.—*Sy. E. B.* 1465. *J. of B.* iii. t. 25.—Height 2—12 inches. Spike dense, cylindrical. Fl. very delicate, pink. Semilanceolate-shaped acute sepals converging over the column and narrow petals and exceeding them. Spur very short. The two ascending fleshy lobes of stigma, and the flat broad plate between them are characteristic of the plant.—Open limestone pastures at Castle Taylor, and elsewhere, Co. Galway. P. IV. V. I.

## 5. HABENA'RIA R. Br. Frog Orchis.

\* Spur very short, usually inflated. CÆLOGLOSSUM, Hartm.

1. *H. vir'idis* (R. Br.); spur 2-lobed, lip linear flat 3-pointed middle point the shortest.—*Orchis* Sm., *E. B.* 94. *R.* xiii. 434. *Peristylus* Lindl.—Lip with 3 tubercles at its base, 1 central, 2 lateral. Stigma oblong, slightly emarginate above. Glands of the pollen-masses connected by an elevated transverse line. Sep.

and pet. connivent. Root-knobs palmate. Fl. green, lip browner. St. 6—8 in. high.—Pastures. P. VI. VII. *Frog Orchis*. E. S. I.

\*\* *Spur slender*. PLATANThERA Rich. Butterfly Orchis.

2. *H. bifolia* (R. Br.); spur twice as long as the germen, lip linear entire, pet. connivent blunt, *anth.* oblong truncate *its cells parallel*.—*E. B. S.* 2803. *R.* xiii. 429.—About a foot high. Root-l. usually 2, elliptic. Stem-l. small, lanceolate, resembling the bracts. Spike slender. Fl. white. Central line between the *anth.*-cells a furrow in front and a keel behind, stalks of pollen-masses short, gland oval. Stigma truncate, emarginate with pointed lobes.—Heathy places. P. VI. VII. E. S. I.

3. *H. chloran'tha* (Bab.); spur twice as long as the germen, lip linear entire, pet. connivent blunt, *anth.* truncate *its cells twice as distant at the base as at the top*.—*Orchis bifolia* Sm., *E. B.* 22. *H. montana* *R.* xiii. 430.—Usually taller and stouter than the preceding. Spike usually lax, but sometimes dense. Fl. larger. Central line between *anth.*-cells a prominent ridge in front and a groove behind. Space between the bases of the *anth.*-cells usually, not always, spread open. Stalks of pollen-masses long, gland circular. Stigma very broad, slightly pointed in the middle.—Moist woods and thickets. P. V. VI. E. S. I.

## 6. O'PHRYS Linn.

1. *O. apifera* (Huds.); lip tumid 5-lobed, 2 lower lobes prominent and with a hairy base, 2 intermediate reflexed truncate, *terminal acute long usually reflexed*, *anth.* with a hooked point, pet. oblong bluntish downy.—*E. B.* 383.—About a foot high. Fl. few, large, rather distant. Sep. whitish tinged with purple. Lip velvety, brown variegated with yellow. "All the lobes of the lip sometimes reflexed, interm. overlapping term. one."— $\beta$ . *O. Trollii* (Heg.); term. lobe of lip acute triangular not reflexed.—On calcareous soils.  $\beta$ . Reigate. P. VI. VII. *Bee Orchis*. E. I.

2. *O. arachnites* (Reichard); lip somewhat tumid entire or with 4 shallow marginal lobes and a *terminal inflexed flat rather heart-shaped appendage*, *anth.* with a straight or hooked point, *pet. deltoid downy*.—*E. B. S.* 2596. *R.* xiii. 461.—Sep. pinkish. Lip velvety, dark purple, variegated with yellow; appendage green, never reflexed.—Chalk downs. Folkestone and Sittingbourne, Kent. P. IV.—VI. *Late Spider Orchis*. E.

3. *O. aranifera* (Huds.); lip tumid obscurely 3-lobed, *middle lobe large emarginate without an appendage*, *anth.* acute, *pet. linear glabrous*.—*E. B.* 65. *R.* xiii. 449.—Smaller than the two

preceding and with fewer flowers. Sep. green. Pet. green, quite glabrous. Lip deep brown, hairy with paler or yellowish glabrous lines often resembling the Greek letter  $\Pi$ , entire at the end or notched with or without a central point.— $\beta$ . *O. fucifera* (Sm.); lip usually undivided often with a gland in the notch, pet. rough. *E. B. S.* 2649.—Chalky places, rare.  $\beta$ . Kent, Sussex, and Isle of Wight. P. IV. V. *Spider Orchis*. E.

4. *O. muscifera* (Huds.); lip oblong trifid with a broad pale spot in the centre, middle lobe long bifid, anth. short blunt, pet. filiform.—*E. B.* 64. *R.* xiii. 447. *St.* 40. 15.—Slender, about a foot high. Sep. green. Lip brownish purple; central spot subquadrate, bluish. Pet. very narrow, purple.—Damp calcareous thickets and pastures. P. V. VI. *Fly Orchis*. E. I.

#### 7. HERMIN'IUM *R. Br.* Musk Orchis.

1. *H. Monorchis* (R. Br.); lip 3-lobed, central lobe longest, pet. with a lobe on each side.—*E. B.* 71.—Root-knobs very unequal and distant. L. usually 2, lanceolate. St. about 6 in. high. Sep. ovate, greenish. Spike dense, slender.—Calcareous soil in the South, rare. P. VI. VII. E.

### Tribe II. Neottideæ.

#### 8. GOODYE'RA *R. Br.*

1. *G. répens* (R. Br.); l. ovate stalked netted, sep. pet. and lip ovate-lanceolate.—*E. B.* 289.—St. 6—8 in. high, bearing linear adpressed bracts. Creeping. Whole upper part of the plant covered with minute stalked glands. L. netted with brown. Fl. white, small.—Fir forests of the North. P. VIII. S.

#### 9. SPIRAN'THES *Rich.* Lady's Tresses.

1. *S. autumnális* (Rich.); root-fibres few ovate-oblong thick, root-l. ovate-oblong in a lateral cluster, stem-l. like bracts, spike dense.—*E. B.* 541.—St. 4—6 in. high. Spike spiral. Fl. greenish white. Column and lid acute; a blunt ovate membranous process between them on each side.—Dry calcareous and gravelly places. P. VIII. IX. E. I.

2. *S. æstivális* (Rich.); root-fibres few long cylindrical, root-l. oblong-lanceolate round the base of the st., stem-l. narrowly lanceolate, spike lax.—*E. B. S.* 2817.—St. 3—12 in. high. Spike spiral. Fl. with a larger lip. Column and lid acute; the in-

term. processes lanceolate acute.—Bogs. Between Lyndhurst and Christchurch, Hants. Wire Forest, Worcest. St. Owen's Pond, Jersey. P. VII. VIII. E.

3. *S. gemmip'ara* (Lindl.); root-fibres few long cylindrical, root-l. linear-lanceolate, stem-l. triangular-lanceolate surrounding the base of the st., bracts shorter than the fl., spike dense 3-ranked, sep. and pet. equal blunt adhering together, lip blunt spathulate.—*Sy. E. B.* 1474.—*S. cernua* Bot. Mag. 5277. *S. Romanzoffiana* Hook.—Spike about  $1\frac{1}{2}$  in. long. Fl. fragrant. Lateral sep. united at the base. See *Rep. Bot. Congr. Lond.* 176.—Castleton, Bearhaven, and between Bandon and Dunmanway, Co. Cork. P. VIII. IX. I.

#### 10. LIS'TERA R. Br. Tway-blade.

1. *L. ovata* (R. Br.); l. 2 opposite ovate, lip bifid, column with a crest which includes the anther.—*E. B.* 1548. *St.* 29. 14.—*St.* 1 ft. high. Spike long, very lax. Fl. small, greenish. L. large.—Woods and pastures. P. V. VI. *Tway-blade.* E. S. I.

2. *L. cordata* (R. Br.); l. 2 opposite cordate, lip 4-lobed, column without a crest.—*E. B.* 358.—Height 3—5 in. *St.* slender. Fl. very small, in a lax spike, greenish. Lip with 2 basal and 2 terminal linear lobes.—Turfy mountain moors. P. VI.—VIII. E. S. I.

#### 11. NEOT'TIA Linn. Bird's-nest.

1. *N. Nidus-avis* (Rich.).—*E. B.* 48. *Listera* Sm.—Whole plant pale reddish brown. Root formed of many short thick fleshy fibres from the extremities of which the young plants are produced. (See *Leight. Fl. Shrop.* 434.) *St.* a foot high, with sheathing brown scales. L. none. Spikes dense, cylindrical, many-flowered. Lip linear-oblong with 2 spreading lobes.—Shady woods. P. ? VI. E. S. I.

#### 12. EPIPAC'TIS Rich. Helleborine.

1. *E. latifolia* (All.); st. solitary, l. broadly ovate exceeding the joints, upper l. ovate-oblong, lower bracts exceeding the fl., label roundish-cordate with a small recurved point falling short of the broadly ovate sep. and pet., basal hunches smooth.—*E. B.* 269.—L. ovate, very broad, the very uppermost sometimes lanceolate-attenuate; lowermost leafless sheaths close.

Lower bracts leaflike, lanceolate, attenuate. Fl. green with the lip purple, sometimes all purple. Peduncle shorter than the downy germen. Label of the lip broader than long, crenate. In a slender form of this plant the upper l. are lanceolate, label cordate blunt with a minute apiculus, and sep. ovate-lanceolate.—Mountain woods. P. VII. VIII. E. S. I.

2. *E. média* (Fries!); st. solitary, l. ovate-oblong the upper ones lanceolate acute, lower bracts exceeding the fl. and fr., label entire triangular-cordate acute equalling the lanceolate sep. and pet., basal hunches plicate rugose.—*Fr. Herb. Norm.* viii. 65!—Narrower and longer in all its parts than *E. latifolia*. Only the very lowest l. ovate, intermediate lanceolate, upper l. lanceolate-attenuate and merging gradually into the linear-lanceolate bracts; sheaths funnelshaped. Fl. green tinged with purple. Ped. shorter than the downy germen. Label longer than broad, crenate. Fr. abruptly obovate.—Woods. P. VIII. E. S. I.

3. *E. violácea* (Bor.); st. many clustered, l. ovate-lanceolate the upper ones narrower passing gradually into slender bracts, label triangular-ovate acuminate falling short of the ovate-lanceolate sep. and pet., hunches plicate-crenate.—*E. B. S.* 2775. *R.* xiii. 486.—Fl. "yellow-green tinged with pink." St. and l. much tinged with purple. Ped. shorter than the downy germen. Label longer than broad, entire, with an attenuate point.—Woods. P. VIII. E.

4. *E. ovális* (Bab.); st. solitary, l. ovate-oblong acute the upper ones lanceolate 1 or 2 lowest bracts exceeding the fl. but falling short of the fr., label transversely oval mucronate equalling the ovate acute sep. and pet., basal hunches plicate-rugose.—*E. B. S.* 2884. Helleborine &c. No. 2. *Ray* 383.—L. small; sheaths funnelshaped, rather close. Bracts all much smaller than even the most uppermost leaf. Fl. varying from a dark yellow to blackish red, peduncle shorter than the downy germen. Label transversely oval, crenate, with a small acute point, and elevated folded and tubercularly crenate hunches above. St. 6—18 in. high.—May be *E. atro-virens* Hoffm.; but that is more often combined with *E. media*.—Settle, Yorkshire. Little Doward Hill, Herefordshire. Ormes Head. Durness, Sutherland. Burren, Clare. P. VII. E. S. I.

5. *E. palustris* (Sw.); l. lanceolate, bracts falling short of the somewhat drooping fl., label roundish blunt crenate equalling the perianth.—*E. B.* 270.—St. 12—18 in. high. Cal. purplish green, pet. and lip white tinged with purple.—Moist places, not rare, P. VII. VIII. E. S. I.

13. CEPHALANTHERA *Rich.*

1. *C. grandiflora* (Bab.); l. ovate or ovate-lanceolate, bracts exceeding the glabrous germen, lip blunt included.—*E. B.* 271. *C. pallens* Koch.—St. 12—18 in. high. Fl. white; lip marked with several elevated longitudinal lines. Sep. erect, blunt.—Woods, usually on a calcareous soil. P. VI. E. S.

2. *C. ensifolia* (Rich.); l. lanceolate, bracts much falling short of the glabrous germen, lip blunt included.—*E. B.* 494.—Height 12—18 in. Fl. white; lips with several elevated white lines and a yellow spot in front. Outer sep. acute.—Woods, rare. P. V. VI. E. S. I.

3. *C. rubra* (Rich.); l. lanceolate acute, bracts exceeding the downy germen, lip acute equalling the pet.—*Sy. E. B.* 1483.—Fl. purple; lip white with a purple margin, marked with many wavy longitudinal lines.—Woods, very rare. In two or three places in Gloucestershire. P. VI. VII. E.

Tribe III. *Arethuseæ.*14. EPIPOGUM *Gm.*

1. *E. aphyllum* (Sw.).—*Sy. E. B.* 1486. *Bot. Mag.* 4821. *R.* xiii. 468. St. 18. 16.—St. 3—7 in. high, sheathed. L. none. Sep. and pet. narrowly lanceolate, acute, pale yellowish. Middle lobe of lip ovate, furrowed, white, with 4 rows of purple tubercles. Spur very thick. Column short, dilated above the stigma to receive the base of the anther. Appendage triangular.—Damp wood near Tedstone Delamere, Herefordshire. Only once found. P. VIII. E.

Tribe IV. *Malaxideæ.*15. CORALLORRHIZA *Hall.*

1. *C. innata* (R. Br.); spur very short or wanting.—*E. B.* 1547.—Root of thick fleshy much-branched fibres. Height 6—12 in. Spike of few yellow flowers. Sep. and pet. lanceolate acute. Lip oblong, white, with a few purple spots, sometimes with 3 equal lobes.—Boggy woods, rare. P. VII. S.

16. MALAXIS *Sw.*

1. *M. paludosa* (Sw.); st. with 3—5 oval concave leaves, lip

concave acute.—*E. B.* 72.—St. 1—4 in. high, 5-edged. Sep. ovate, spreading, 2 turning upwards. Lip superior, erect, 3-veined, its base surrounding the column. L. fringed at the end with little bulbs. Forming a small bulb at its base.—This plant and the next rather grow upon the moss as epiphytes than amongst it.—Spongy bogs. P. VIII. IX. E. S. I.

### 17. STURM'IA *Reichen.*

1. *S. Loesel'ii* (R.); l. oblong-lanceolate, st. triangular, lip obovate exceeding the petals.—*Malaxis* Sm., *E. B.* 47. *Liparis* Hook.—St. 6—10 in. high. Fl. 6—12, in a lax spike, yellowish. Sep. lanceolate. Pet. linear. Forming a large ovate bulb at its base, enclosed in the whitish sheaths of the decayed leaves. An epiphyte?—*Liparis* is the name of a genus of insects, and also of fish. I follow Koch in adopting *Sturmia*.—Spongy bogs in Norf., Suff., and Camb., now very rare. P. VI. E.

### Tribe V. *Cypripediæ.*

#### 18. CYPRIPE'DIUM *Linn.* Lady's Slipper.

1. *C. Calcéolus* (L.); st. leafy, middle lobe of the column nearly ovate blunt deflexed, lip slightly depressed falling short of the calyx.—*E. B.* 1.—St. 12—18 in. high, downy, bearing 3 or 4 large ovate pointed leaves. Fl. usually solitary or 2, large; sep. 1—1½ in. long, dark brown; pet. dark brown, rather narrower than sep.; lip 1 in. long, inflated, yellow, netted with darker veins.—Dense Northern woods, very rare. P. V. VI. E.

### Order LXXXIV. IRIDACEÆ.

Perianth tubular, 6-parted, petal-like, in 2 often unequal rows. Stam. 3, epigynous, opposite the outer segments of the perianth. Anth. bursting outwards. Ovary inferior, 3-celled. Style 1. Stigmas 3, dilated, often like petals. Caps. 3-celled, 3-valved; valves bearing the dissepiments in the middle. Seeds many. Embryo cylindrical, enclosed in horny or fleshy albumen. Radicle pointing towards the hile.

1. SISYRINCHIUM. Perianth 6-cleft; segments nearly equal, patent. Style short. Stigmas 3, rolled inwards, filiform. Filaments connate below.



2. IRIS. Perianth 6-cleft; alternate segments reflexed. Stigmas 3, like petals, covering the stamens.
3. GLADIOLUS. Perianth irregular, 6-cleft, 2-lipped; segments unequal. Style filiform. Stigmas 3, widening upwards. Seeds more or less winged.
4. ROMULEA. Perianth regular, 6-cleft; segments spreading. Style filiform. Stigmas 3, bifid; lobes slender.
5. CROCUS. Perianth regular, funnelshaped with a long tube; limb bellshaped. Style filiform. Stigma 3-fid or 3-parted; lobes widening upwards.

### 1. SISYRINCHIUM Linn.

1. *S. Bermudia'num* (L.); scape 2-edged nearly simple nearly leafless, spath 1—6-flowered falling short of the fl., segments of perianth emarginate mucronate.—*Sy. E. B.* 1491. *Redoute Lil.* v. 282.—St. about 1 foot high. L. grasslike. Spath with about equal lanceolate valves, falling short of the fl. in my Irish specimens. Perianth blue; segments narrowed below.—Near Woodford, Galway, abundant. P. VII. I.

### 2. IRIS Linn. Flag.

1. *I. Pseud-ac'orus* (L.); l. swordshaped, st. roundish, perianth beardless its inner segments narrower and falling short of the stigmas.—*E. B.* 578.—St. terete, 2—5 ft. high. Fl. yellow, or in *I. Bastardi* (Bor.) fl. pale lemon-coloured. Caps. oblong, trigonous, apiculate.—*a*; outer perianth-segm. uniform clear yellow with broadly obovate blade and rather short claw, stigm. narrow and long.—*β. I. acoriformis* (Bor.); outer perianth-segm. with deeper blotch at the base, the blade suborbicular and claw long narrowish yellow with prominent purplish veins. Yellow of fl. paler than in *a*.—Wet places. P. VI. VII. *Yellow Flag.* E. S. I.

2. *I. fœtidis'sima* (L.); l. swordshaped, st. compressed, perianth beardless its inner segments about equalling the stigmas.—*E. B.* 596. *R. ix.* 347.—Herb green, not glaucous, yielding an unpleasant smell when bruised. St. 1-angled, 2 ft. high. Fl. lead-coloured or bluish, rarely yellow.—Woods and thickets. P. V.—VII. *Gladdon.* E. I.

[\**I. tuberosa* (L.); l. tetragonal, segments of the perianth acute, root tuberous.—Penzance. Cork.—*E. B. S.* 2818.] E. I.

3. GLA'DIOLUS *Linn.*

1. *G. illyricus* (Koch); corm clothed with nearly parallel fibres netted above with long narrow openings, fl. secund, filaments longer than the anthers, cor.-tube nearly thrice as long as germen, caps. oval emarginate with 3 rounded angles.—*Sy. E. B.* 1493. *J. of B.* i. t. 4. *St.* 83. 3.—Height about 20 in. Corms ovate-acuminate. L. slender, swordshaped. Sheaths 2-edged. Fl. reddish changing to bluish; basal pet. rather acute, exceeding the blunt lower lateral pet.; “edges of upper pet. not covered by the 2 adjoining ones” when the fl. are in their prime. Stigm. narrow below widening from the middle upwards and there fringed.—Amongst *Pteris* in the New Forest. Isle of Wight. *Mr. A. G. More.* P. VII. E.

4. ROMULEA *Mar.*

1. *R. Colum'næ* (R.); scape 1-flowered usually solitary slightly nodding, l. filiform compressed furrowed recurved, spath exceeding the tube of the cor., style falling short of the stam., stigmas bifid.—*E. B.* 2549. *R.* ix. 354. *T. Bulbocodium* Sm.—A small plant not more than 4 in. high, with a corm. Fl. greenish without, pale with purple stripes and yellow at the base within.—Sandy places. Dawlish Warren, Devon. Jersey and Guernsey. P. III. IV. E.

5. CROCUS *Linn.*

\* *Scapes enveloped in a tubular sheath.*

1. *C. ver'nus* (Willd.); l. and fl. at the same time, spath simple, throat of the cor. fringed with hairs, stigma shortly 3-fid, lobes erect wedgeshaped jagged at the end, corm clothed with slender netted fibres.—*E. B.* 344.—Fl. 1—2, violet-purple.—Near Nottingham; and Mendham, Suff. P. III. E.

2. *C. nudiflorus* (Sm.); l. succeeding the fl., spath simple, stigma in 3 deeply lacinate divisions erect, corm with a membranous coat.—*E. B.* 491. *C. speciosus* Hook. *E. B. S.* 2752.—L. linear, appearing in March. Fl. solitary, purple. Stigmas only a little higher than the anthers, or rising considerably above them.—Meadows. P. IX. E.

\*\* *Scapes naked.*

[*C. argenteus* (Sal.); l. and fl. at the same time, spath double,

*stigma* longer than the *stam.* erect deeply trifold, *divisions* truncate and *slightly notched* at the end, *corm* with a membranous coat.—*E. B. S.* 2645.—Fl. pale lilac with yellow and purple stripes.—And *C. aureus* (Sibth.); l. and fl. together, *spath simple, stigma shorter than the stam. shortly 3-fid*, segments truncate or slightly notched at the end, *corm* coated with compact fibres.—*E. B. S.* 2646.—Fl. yellow.—On site of old garden, Barton, Suff. P. III.]

### Order LXXXV. AMARYLLIDACEÆ.

*Stam.* 6. *Anth.* bursting inwards. Raphidiferous. Otherwise like *IRIDACEÆ*.

1. NARCISSUS. Perianth tubular below; limb 6-parted, spreading, with equal segments, and a bellshaped crown within. *Stam.* alternately shorter, within the crown.
2. LEUCOJUM. Perianth bellshaped, 6-parted; segments all equal and thickened at their points. *Stam.* equal.
3. GALANTHUS. Perianth 6-parted, 3 outer segments spreading, 3 inner shorter erect emarginate. *Stam.* equal, subulate.

#### 1. NARCIS'SUS *Linn.*

†1. *N. biflorus* (Curt.); l. linear blunt slightly keeled, scape compressed 2-edged striate 1—2-flowered, crown very short concave crenate at the pale (ultimately white) margin.—*E. B.* 276. *R.* ix. 365.—Pet. of a pale sulphur-colour.—Sandy fields in the South. A doubtful native. P. IV. V. E. I.

[2. *N. poet'icus* (L.); l. linear blunt slightly keeled, scape compressed 2-edged mostly 1-flowered, crown very short concave crenate at the red margin.—*E. B.* 275. *R.* ix. 364.—Pet. white, broadly ovate, crown yellow.—Heathy open fields on a sandy soil. Norf., Kent. P. V.] E.

3. *N. Pseudo-narcis'sus* (L.); l. linear blunt not keeled, scape 2-edged 1-flowered, perianth-segm. scarcely exceeding tube, crown bellshaped crisped at the margin and crenate equalling the perianth, pedicel within the *spath* short.—*E. B.* 17. *R.* ix. 369.—L. 2 or 3, scarcely a foot long. Fl. large, yellow.—Open woods and pastures. P. III. IV. *Daffodil.* E. S.

[*N. lobularis* (Schult.); “perianth-segm. exactly twice as long as tube, crown 6-lobed.” Tenby.—*N. minor* (L.); “crown 6-

lobed equalling the perianth, pedicel long." Penrice Castle, Glam.—*N. incomparabilis* (Curt.) *Sy. E. B.* 1502; crown erect  $\frac{1}{2}$  as long as perianth, otherwise like Sp. 3. In several places in the South.—These are not native plants.]

## 2. LEUCOJUM *Linn.* Snowflake.

†1. *L. æstivum* (L.); spath many-flowered, style thickened upwards.—*E. B.* 621. *R.* ix. 362.—Bulbous. Height 2—2 $\frac{1}{2}$  feet. Fl. æstival, white, drooping; tips greenish. L. broadly linear, keeled, hibernal. Scape 2-edged.—Wet meadows. P. V. E.

†2. *L. ver'num* (L.); spath 1-flowered, style thickened upwards.—*Sy. E. B.* 1506. *J. of B.* iv. t. 49.—Bulbous. Height 8—10 inches. L. and white drooping flowers vernal.—Near Bridport and Bicester. P. II.—IV. E.

## 3. GALAN'THUS *Linn.* Snowdrop.

1. *G. nivâlis* (L.).—*E. B.* 19. *R.* ix. 363.—Bulbous. Fl. solitary, white, drooping; inner segments greenish. L. 2, keeled, broadly linear, glaucous.—Thickets in the West. P. II. III. E.

## Order LXXXVI. ALISMACEÆ.

Perianth free, of 6 leaves; 3 inner or all coloured. Stam. 6. Ovaries 3—6 or many, always distinct or ultimately separable; carp. opening at the suture or not at all; seeds 1 or many. Embryo straight or curved; albumen 0.—No raphides.

### Suborder I. ALISMOIDEÆ.

Three inner perianth-segm. petal-like. *Seeds* 1—2 in each cell, erect or ascending, on the suture. Embryo cylindrical, doubled upon itself; radicle next the hile.

1. ALISMA. Fl. perfect. Stam. 6. *Carp. many, forming a ring or head*, small, 1-seeded, not bursting.—L. ovate or narrow.
2. ACTINOCARPUS. Fl. perfect. Stam. 6. *Carp. 6—8, rather large, 2-seeded, combined at the base and spreading in a radiant manner.* L. cordate-oblong.
3. SAGITTARIA. Fl. monœcious. Male fl. with many sta-

mens. Female fl. with *many* 1-seeded compressed *carpels forming a head*, upon a globose receptacle.—*L. sagittate*.

### Suborder II. BUTOMEÆ.

Three inner perianth-segm. petal-like; three outer herbaceous or slightly coloured. *Seeds many*, minute. Placenta ramified over the inner surface of each carpel. Embryo straight or curved; radicle next the hile.

4. BUTOMUS. Perianth-segm. 6, all coloured, resembling a corolla. Stam. 9. Carpels 6, connected below.

### Suborder III. JUNCAGINEÆ.

*Perianth uniform*, herbaceous, inconspicuous, or 0. Seeds 1—2, erect, close together and at the base of the carpel. Embryo straight; radicle next the hile; plumule coming through a lateral cleft in the embryo.

5. SCHEUCHZERIA. Perianth of 6 reflexed leaves. Stam. 6, with slender filaments. Ovaries 3. Stigma sessile, downy. Carpels compressed, inflated, diverging, 1—2-seeded, free.
6. TRIGLOCHIN. Perianth of 6 erect deciduous leaves. Stam. 6; anth. almost sessile. Ovaries 3—6. Stigmas sessile, feathery. Carp. attached to an angular axis, from which they at length separate at the base.

### Suborder I. *Alismoideæ*.

#### 1. ALIS'MA *Linn.* Water-Plantain.

1. *A. Plantago* (L.); fl.-stalk paniced with whorled compound branches, *carp.* ranged in a circle *compressed blunt* obovate, *style below the top of inner edge of carp.*, l. cordate-ovate or lanceolate.—*E. B.* 837. *R.* vii. t. 57.—*L.* all radical, on long stalks. Submersed l. linear. Fl.-stalks 2—3 ft. high. Fl. pale rose-colour. Sep. ovate-oblong, styles twice as long as ovary.— $\beta$ . *A. lanceolata* (With.); l. lanceolate narrowed below. *Sy. E. B.* 1438. Sep. ovate, styles equalling the ovary. Perhaps distinct.—By water. *P.* VII. VIII. *E. S. I.*

2. *A. ranunculoides* (L.); fl.-stalks umbellate, *carp. angular acute* forming a globose squamose head, *style terminal*, l. linear-lanceolate acute.—*E. B.* 326. *R.* vii. t. 55.—*L.* all radical, on long stalks. Fl.-stalks from 3 to 24 in. long, ending in 1 or 2 umbellate whorls of simple peduncles. Fl. pale purple.— $\beta$ . *A.*

*repens* (Dav.); stoloniferous, producing l. roots and fl. at the nodes, fl. larger. *E. B. S.* 2722.—Turfy bogs.— $\beta$ . By lakes in Wales and Ireland. *P. VI. VII.* E. S. I.

3. *A. natans* (L.); st. floating and rooting leafy, *peduncles simple* from the joinings of the stem, *carp.* striate beaked, floating l. stalked oblong blunt, radical leafless petioles broadly subulate.—*E. B.* 775. *R. vii. t.* 54.—St. slender, often very long. Root-petioles in small tufts. Fl. rather large, white with a yellow spot.—Lakes, rare. *P. VIII.* E. I.

## 2. ACTINOCAR'PUS *R. Br.* Thrumwort.

1. *A. Damasónium* (R. Br.); stalks with 1—3 whorls of fl., *carp.* subulate compressed opening longitudinally, l. cordate-oblong.—*E. B.* 1615. *Alisma* L.—L. all radical, floating, on long stalks, 5-veined. Pet. white. *Carp.* large; with 2 stalked seeds, one from the lower angle erect, one from the upper horizontal.—Ponds and ditches, rare. *P. VI. VII.* E.

## 3. SAGITTA'RIA *Linn.* Arrowhead.

1. *S. sagittifolia* (L.); aërial l. arrowshaped with lanceolate straight lobes, fl.-stalk simple, fl. whorled.—*E. B.* 84. *R. vii. t.* 53.—Stoloniferous; each runner ending in a tuber. The l. that rise above the water are remarkably arrowshaped, with the 3 parts nearly equal. The submersed leafless petioles are linear. Fl. white.—Ditches and rivers. *P. VIII.* E. I.

## Suborder II. *Butomeæ.*

### 4. BU'TOMUS *Linn.* Flowering Rush.

1. *B. umbellatus* (L.).—*E. B.* 651. *R. vii. t.* 58.—Flowerstalk radical, 2—3 feet high, overtopping the leaves, bearing an irregular many-flowered simple umbel with scarious bracts and a membranous 3-leaved involucre. Fl. rose-coloured. L. radical, slender, triangular.—Rivers and ponds. *P. VI. VII.* E. S.

## Suborder III. *Juncagineæ.*

### 5. SCHEUCHZE'RIA *Linn.*

1. *S. palus'tris* (L.).—*E. B.* 1801. *St.* 78.4. *R. x.* 419.—St. 6—8 in. high, erect. L. distichous, few, alternate, semicylindrical, blunt, with a minute pore on the upper side of the apex. Raceme terminal, of about 5 greenish flowers. Caps. about 3, much inflated.—Sphagnous bogs, rare. *P. VII.* E. S.

6. TRIGLO'CHIN *Linn.* Arrow-grass.

1. *T. maritimum* (L.); fr. ovoid of 6 combined carpels.—*E. B.* 255. *R.* vii. t. 52.—L. radical, linear. Fl. in a lax simple spike or raceme, greenish.—Muddy salt marshes. P. VII. VIII. E. S. I.

2. *T. palustris* (L.); fr. linear angular of 3 combined carpels.—*E. B.* 366. *R.* vii. t. 51.—Slenderer than the preceding but closely resembling it, 8—10 in. high. Stoloniferous.—Marshy places. P. VI. VII. E. S. I.

## Order LXXXVII. ASPARAGACEÆ.

Perianth inferior, petal-like, 6-parted or 4—8-parted. Stam. 6 or 4—8, inserted into the receptacle or on the perianth. Anth. bursting inwards. Ovary superior, 3- (rarely 1-) celled. Ovules 1 or many in each cell. Styles 1—3. *Fr. succulent*, not bursting. *Root not bulbous*.—Raphidiferous.

1. ASPARAGUS. Perianth 6-parted, bellshaped, tubular below. Stam. 6. Ovary 4-celled; cells 2-ovuled. *Style* 1. *Stigmas* 3, *reflexed*.—Fl. by abortion diœcious.
2. CONVALLARIA. *Perianth* bellshaped, 6-parted, deciduous. Ovary 3-celled; cells 2-ovuled. *Stigma blunt*, trigonous. Berry with 1-seeded cells. Fl. jointed to pedicel.
3. POLYGONATUM. *Perianth* tubular, 6-toothed, tardily deciduous. Ovary 3-celled; cells 2-ovuled. *Stigma blunt*, trigonous. Berry with 1—2-seeded cells. Fl. not jointed to pedicel.
4. MAIANTHEMUM. *Perianth* 4-parted; segments horizontally patent or reflexed, deciduous. Stam. 4. *Style* 1, bifid. *Stigma blunt*. Berry 2-celled; cells 1-seeded.
5. RUSCUS. Diœcious. *Perianth* 6-parted to the base, persistent. Filaments forming an ovoid tube, on the top of which the 3 anth. are placed.—Fem. the same, but the anthers barren. *Style* 1. *Stigmas* capitate. Berry 1-celled, rarely 2-seeded.—Fl. on disk of persistent leaflike shoots.

1. ASPAR'AGUS *Linn.*

1. *A. officinalis* (L.); st. herbaceous mostly erect without spines branched, l. clustered terete flexible setaceous.—*E. B.* 339.—Creeping. Stems many.—*a. A. prostratus* (Dum.); st.

prostrate at the base, branches short about 1 ft. long. *Bull. S. Bot. Belg.* i. t. 2.—\**β. campestris*; st. erect about 3 feet high, branches long.—Sea-coast, rare. Kynance Cove, Cornwall. South coast of Anglesea. Giltar Point, Pemb. Tramore, Waterf.—*β.* Escape from cultivation. P. VIII. E. I.

## 2. CONVALLA'RIA Linn. Lily of the Valley.

1. *C. majalis* (L.).—*E. B.* 1035. *St.* 14. 10.—About a foot high. L. 2, ovate-lanceolate, radical. Scape semicylindrical. Fl. racemose, nodding, pure white, globose-bellshaped, fragrant.—Woods and thickets. P. V. E. S.

## 3. POLYGONA'TUM Tourn. Solomon's Seal.

1. *P. verticillatum* (All.); *l. linear-lanceolate whorled*, st. erect angular.—*E. B.* 128. *R. x.* 435.—*St.* 2 feet high. L. 3—5 in a whorl. Berries red.—Woods. Perthshire. Smalesmouth, Northumberland. P. VI. E. S.

2. *P. officinale* (All.); *l. ovate-oblong half-clasping glabrous alternate, st. angular*, peduncles 1—2-flowered, cor. not narrowed in the middle, *filaments glabrous*.—*Convallaria Polygonatum* (L.), *E. B.* 280. *R. x.* 434.—Height 1—1½ foot. Berry bluish.—Woods, rare. P. V. E.

3. *P. multiflorum* (All.); *l. ovate-oblong half-clasping glabrous alternate, st. round*, peduncles 1- or many-flowered, cor. narrowed in the middle, *filaments downy*.—*E. B.* 279. *R. x.* 433.—Height 2 feet. Berry bluish.—Woods. P. V. E. S.

## 4. MAIAN'THEMUM Wiggers.

1. *M. bifolium* (DC.); st. with 2 alternate stalked triangular-cordate leaves.—*Ger. Herb.* p. 409. *Smilacina, Sy. E. B.* 1510. *R. x.* 436.—*St.* 6—8 in. high. Root filiform. L. very deeply cordate. Raceme terminal, resembling a spike. Fl. small; segments reflexed.—Woods. Near Scarborough, in plenty. Howick, Northumb. (now eradicated). Caen Wood, Middlesex. Dingley Wood, Preston; Harwood, Blackburn. *Gerard.* P. V. E.

## 5. RUS'CUS Linn. Butcher's Broom.

1. *R. aculeatus* (L.); leaflike flattened shoots ovate-attenuate very acute rigid bearing the fl. upon the middle of their upper surface, fl. 1 rarely 2 with a flat subulate scarious 1-veined bract.—*E. B.* 560. *R. x.* 437.—Evergreen. Fl. small. L. very minute.—Thickets. Sh. III. IV. E. S. ?



## Order LXXXVIII. LILIACEÆ.

Perianth inferior, petal-like, 6-leaved, 6-parted or with 6 teeth. Stam. 6, inserted on the receptacle or on the perianth. Anth. bursting inwards. Ovary superior, 3-celled. Ovules many in each cell. Style 1. Stigmas 3 or 1. *Fr. dry*, capsular, bursting with 3 valves bearing the dissepiment on their middle.

Tribe I. *TULIPEÆ*. Perianth-l. nearly or quite distinct. Cells of caps. many-seeded. Seeds flat (in *Lloydia* angular), placed closely one above another; testa pale or fuscous, not crustaceous.—St. usually more or less leafy. Bulbous.

1. *TULIPA*. Perianth without nectaries, deciduous. *Anth. erect*. Style 0. Stigma 3-lobed. Seeds flat.
2. *FRITILLARIA*. Perianth deciduous; a nectariferous depression at the base of each leaf. *Anth. attached above their base*. Style 3-fid at the apex. Seeds flat.
- [3. *LILIUM*. Perianth deciduous, spreading or reflexed; a longitudinal nectariferous furrow at the base of each leaf. *Anth. attached above their base*. Stigma capitate. Seeds flat.]
4. *LLOYDIA*. Perianth persistent, patent. Stam. inserted at the base of perianth. *Anth. erect*. Style filiform. Stigma trigonous. Seeds angular above, flat beneath.

Tr. II. *ASPHODELEÆ*. Fl. not jointed to their stalks. Leaves of perianth distinct. Cells of caps. few-seeded. Seeds various in form, usually with a black crustaceous testa.—St. usually leafless. Bulbous.

5. *ORNITHOGALUM*. Perianth-l. 6, spreading, persistent. Stam. on the receptacle and adhering only slightly to the perianth. *Anth. attached by their backs*.—Fl. white or yellow, never blue.
6. *GAGEA*. Perianth-l. 6, spreading, persistent. Stam. adhering to the base of the perianth. *Anth. erect*.—Fl. corymbose or umbellate, yellow.
7. *SCILLA*. Perianth-l. 6, spreading, deciduous. Stam. on the base of the perianth. *Anth. attached by their backs*.—Fl. racemose, never white or yellow.
8. *ALLIUM*. Perianth-l. 6, rather spreading. Stam. at the base of the perianth. *Anth. attached by their backs*.—Fl. umbellate. Spath of 1 or 2 leaves.

Tr. III. *ANTHERICEÆ*. Fl. jointed to their stalks. Leaves of perianth slightly connected below. Cells of caps. few-seeded. Seeds various in form.—Not bulbous.

9. SIMETHIS. Perianth-l. 6, spreading, deciduous. Stam. on the base of the perianth. Filaments bearded. *Anth.* attached by their backs. Caps. 3-celled, 6-seeded.

Tr. IV. *HEMEROCALLIDEÆ*. Leaves of the perianth combined below. Cells of the caps. few-seeded. Seeds various in form; testa (in our plants) black.—Bulbous.

10. ENDYMION. Perianth tubular-bellshaped, of 6 connivent leaves with reflexed points combined below. Stam. inserted below the middle of the perianth; filaments decurrent.
11. MUSCARI. Perianth globose or subcylindrical, narrowed at the mouth, 6-toothed. Stam. inserted at about the middle of the tube; filaments not decurrent.

### Tribe I. *Tulipeæ*.

#### 1. TULI'PA *Linn.* Tulip.

1. *T. sylves'tris* (L.); st. 1-flowered glabrous, fl. at first drooping, tip of segments of perianth and base of the stamens hairy.—*E. B.* 63. *St.* 29. 11. *R.* x. 446.—Fl. yellow, rarely produced in a wild state.—Chalk-pits in the Eastern Counties. "Meadows near Nottingham and in Yorkshire." P. IV. V. E.

#### 2. FRITILLA'RIA *Linn.* Fritillary.

1. *F. Meleágris* (L.); st. 1-flowered leafy, l. all alternate linear-lanceolate.—*E. B.* 622. *St.* 18. 4. *R.* x. 442.—About a foot high. Fl. dull red with many dark spots, rarely white.—Meadows and pastures in the East and South. P. V. E.

#### 3. LIL'IUM *Linn.* Lily.

[1. *L. Martágon* (L.); l. whorled elliptic-lanceolate, st. downy roughish, fl. nodding, perianth reflexed.—*E. B. S.* 2799. *R.* x. 451.—Height 1—1½ foot. Fl. violet-flesh-coloured with dark-purple spots.—Copses. P. VI. VII. *Turk's-cap Lily.*] E.

[*L. pyrenáicum* (Gouan); l. scattered linear-lanceolate, fl. nodding, perianth reflexed.—*Sy. E. B.* 1517.—About 1 ft. high. Fl. yellow with black dots below.—Between South Molton and Mollond, Devon.] E.

4. LLOYD'IA *Salisb.*

1. *L. serot'ina* (R.); root-l. semicylindrical, st. l. dilated below and sheathing, fl. mostly solitary, nectary a transverse plait.—*E. B.* 793. *St.* 28. 2. *R.* x. 440. *Anthericum* Sm.—Height 5 or 6 in. St. and l. springing separately from the root. St.-l. several, short. Fl. white with reddish lines internally.—Welsh mountains, very rare. Snowdon. Glyder Fawr. P. VI. E.

Tribe II. *Asphodeleæ.*5. ORNITHOG'ALUM *Linn.* Star of Bethlehem.

†1. *O. umbellatum* (L.); fl. corymbose, ped. exceeding the linear-lanceolate bracts, filaments lanceolate simple, l. linear glabrous.—*E. B.* 130. *R.* x. 467.—L. exceeding the stem or filiform and shorter. Height 8—12 in. Fl. white with a broad green longitudinal band externally. Meadows and pastures. P. V. E. S.

2. *O. pyrenæicum* (L.); fl. in an elongated narrow raceme, ped. at first spreading afterwards erect, bracts lanceolate-acuminate, filaments dilated below with a long point, l. soon fading linear grooved.—*E. B.* 499.—St. leafless, 2—3 feet high. Raceme becoming very long. Fl. greenish white; segments of the perianth variable in breadth. L. withering before the stalk appears, rarely contemporaneous.—Woods. Extremely common near Bath. Sussex. Bedfordshire. P. VI. E.

[3. *O. nitans* (L.); fl. few in a lax nodding raceme, ped. falling short of the bracts, filaments flat membranous trifid, the lateral points acute, middle one very short bearing the anther, l. linear-lanceolate.—*E. B.* 1997. *Albuca* R. x. 473.—Height 9—12 in. Fl. large, white, greenish externally.—Fields and orchards, rare. B. IV. V.] E.

6. GA'GEEA *Salisb.*

1. *G. lutea* (Ker); radical l. usually solitary linear-lanceolate flat, bracts 2 opposite, peduncles umbellate simple glabrous, segments of the perianth oblong blunt, bulb ovate solitary.—*E. B.* 21. *R.* x. 477. *Ornithogalum* L.—St. about 6 in. high, shorter than the leaves. Bracts lanceolate, one often exceeding the yellow flowers. Bulb often enclosing many small round offsets.—Woods and thickets, rare. P. III. IV. E. S.

## 7. SCILLA Linn. Squill.

1. *S. autumnalis* (L.); l. linear many, raceme lax, peduncles ascending, *bracts* 0.—*E. B.* 78. *R. x.* 463.—Height 4—6 in. Fl. purplish blue with a green line down the back, in perfection before the l. appear.—Dry pastures in the South and West. P. VIII. *Autumnal Squill.* E.

2. *S. ver'na* (Huds.); l. linear channelled hooded at the end many, raceme few-flowered corymbose, *bracts lanceolate* as long as or longer than the peduncles.—*E. B.* 23. *R. x.* 463.—Height 4—5 in. Fl. blue. L. as long as or longer than the stalk.—Western and Northern coasts. P. IV. V. *Vernal Squill.* E. S. I.

## 8. ALLIUM Linn. Garlic.

\* *Alternate stamens broader and 3-pointed, the middle point alone bearing an anther.* PORRUM Tourn.

† Stem-leaves flat or keeled, not hollow.

\**A. Ampelop'rasum* (L.); st. leafy below, l. linear, spath long, *umbel globose compact*, *stam.* exerted, *anther-bearing point* of 3-pointed filaments as long as the undivided part, bulb compound of 2—4 divisions.—*E. B.* 1657.—Bulb with large off-sets within its coats. St. 2—6 feet high. L. long, linear. Spath parting at the base, and falling off in one piece before the fl. open; horn 1—2 in. long. Fl. pale purple, the keel of the outer submarginate segments greenish and roughish. Germs rather globose; transverse projections at about the middle; lower spaces slightly excavated. *Head-bulbs very rare*, when present small, *the size of peas*.—Steep Holmes Islands in the Severn (remains of former cultivation). Cliffs in Guernsey. P. VIII. E.

†1. *A. Babingtonii* (Borr.); st. leafy below, l. acutely keeled, spath long-pointed, *umbel loose irregular* with hemispherical bulbs, *stam.* exerted, *anther-bearing point* of 3-pointed filaments rather shorter than the undivided part and *with an incurved tip when young*, bulb compound of few (2) divisions.—*E. B. S.* 2906.—St. 4—6 feet high. L. long, linear, broad. *Heads large, with many bulbs about as large as hazelnuts* (a character quite constant in cultivation). Spath usually deciduous in two pieces or persistent; horn 1—2 in. long. Fl. pale reddish purple, rather few, more conical than in the preceding, only slightly opening; outer segments with a green keel, ovate-oblong, with callous points, edges and back rough with minute pellucid points; inner segments slightly emarginate and without points. Lengthened stalks 1—2 in. long, bearing *secondary*

heads, are usually present. Germen rather conical; transverse projection below the middle; lower spaces rather deeply excavated.—Roundstone sparingly, and South Isles of Aran plentifully, Co. Galway. Cornwall. P. VIII. E. I.

2. *A. Scorodop'rasum* (L.!); st. leafy below, l. flat, sheaths 2-edged, *spath* short and broad with a very short point, umbel globose with many spherical small bulbs, *stam.* included or equalling the perianth, anther-bearing point of 3-pointed filaments shorter than the undivided part or the lateral points, bulb with many purple offsets.—*E. B. S.* 2905. *A. arenarium* Sm.—St. 2—3 feet high. Heads small. Fl. few, purple; segments all with a minute apiculus, outer with the edges and keel rough. Head-bulbs deep purple.—Sandy woods and fields in the North. P. VI. VII. E. S. I.

†† Stem-leaves hollow.

3. *A. vineale* (L.); st. leafy below, l. *terete* slightly channelled above, *spath* 1-valved short with a slender long point, umbel globose with many bulbs, *stam.* exerted, *anther-bearing point of 3-pointed filaments equalling the undivided part and half as long as the lateral points.*—*E. B.* 1974. *R. I.* t. 404. *A. arenarium Fries.*—St. 2 feet high. L. faded at the time of flowering. Heads of few pale rose-coloured fl. with green keels and long stalks. Head-bulbs small, oval, acute, greenish.— $\beta$ . *A. compactum* (Thuil.); umbel without fl., head-bulbs with a leaflike point.—Waste ground and dry fields.  $\beta$  is the more common state. P. VII. *Crow-Garlic.* E. S. I.

4. *A. sphaerocephalum* (L.); st. leafy below, l. subcylindrical channelled above, *spath* 2-valved short, *umbel* globose without bulbs, *stam.* twice as long as the perianth, *anther-bearing point of 3-pointed filaments as long as the undivided part longer than the lateral points,* bulb accompanied by stalked offsets.—*E. B. S.* 2813.—St. 1—2 feet high. L. usually faded before the time of flowering. Heads of many rose-coloured or purple fl.; keels darker and rough.—St. Vincent's Rocks, Bristol. Sands in Jersey. P. VII.

\*\* *Stam.* all simple, not 3-pointed, connected at the base. *Spath* 2-valved, one valve with a long point. *St.-l.* narrow.

5. *A. oleraceum* (L.); st. leafy below, l. channelled above ribbed beneath, *spath* with one of the points very long, umbel with bulbs, *stam.* equalling or shorter than the perianth.—*E. B.* 488.—Height 1—2 feet. L. (of the Bristol plant) thick, fleshy, solid, nearly flat but slightly and broadly channelled above, with 4 ribs beneath. Segments of perianth blunt.— $\beta$ . *A. complanatum*

(Bor.); *stam.* shorter than the perianth, l. of equal thickness throughout curved upwards at the sides so as to appear channelled with many ribs on each side. *A. carinatum* Sm., *E. B.* 1658.—Borders of fields, rare.  $\beta$ . Mountains in the North. P. VII. VIII. E. S.

6. *A. carinatum* (L.); *st.* leafy below, l. nearly flat, umbel nearly without bulbs, *stam.* much exceeding perianth.—Height 1—2 feet. L. erect, slightly channelled below, flat in the upper part, slightly furrowed (not keeled) beneath. Perianth-segm. blunt, rose-coloured.—Newark. Seguden, Carse of Gowrie. By Esk above Musselburgh, abundant. P. VIII. E. S.

\*\*\* *Stam.* all simple and distinct. *Spath* 2-valved, short.  
*Leaves* hollow.

7. *A. Schœnoprasum* (L.); *st.* leafless or with one leaf, l. terete or slightly flattened above subulate, *spath* ovate pointed about equalling the flowers, umbel many-flowered globose without bulbs, *stam.* about half the length of the lanceolate segments of the perianth.—*E. B.* 2441.—*St.* about 6 in. high. L. straight, mostly with smooth ribs. Pet. lanceolate. Barren bulbs with leaves. Fl. pink. Forming dense tufts.— $\beta$ . *A. sibiricum* (L.); l. curved and bent downwards with crenulate ribs, pet. lanceolate-attenuate, barren bulbs single-leaved, style longer than the young germen. *E. B. S.* 2934. Height 6 in. to 2 feet. Heads large. Possibly a distinct species.—Meadows and pastures in mountainous situations.  $\beta$ . Rocks and cliffs near the sea. Tintagel and Rill Head, also between Kynance Cove and Mullion, Cornwall. P. VI. VII. Chives. E. S.

\*\*\*\* *Stam.* all simple. *Leaves* flat, all radical.

† 8. *A. triquetrum* (L.); *st.* triquetrous, l. linear acutely folded and keeled, *spath* 2-valved about equalling the erect bulbless lax umbel, *stam.* half as long as the oblong segments of the perianth.—*E. B. S.* 2963. *R. x.* 503.—Bulb ovate. L. angularly folded, acute. Segments of perianth white with a slender green midrib.—Hedges in Guernsey. Helston, Corn. P. V. VI. E.

9. *A. ursinum* (L.); *st.* naked triangular, l. stalked ovate-lanceolate, *spath* 2-valved ovate, umbel level-topped lax bulbless.—*E. B.* 122.—Bulb slender, oblong. L. few, broad, smooth, bright green. Stalk one, as tall as or taller than the leaves. Fl. white. Smelling strongly of garlic when bruised.—Damp woods and hedges. P. V. VI. Ramsons. E. S. I.

[*A. ambiguum* (Sm.); *E. B. S.* 2803.—Rochester. Eye Castle Hill, Suff. Not a native; nor is *A. paradoxum* (Don) at Binnig Craig, Linlithgow.]

Tribe III. *Anthericeæ*.9. *SIME'THIS* Kunth.

1. *S. bicolor* (K.).—*E. B. S.* 2952.—Root of fleshy fibres. L. linear, flat or a little keeled upwards. St. and l. enclosed in sheathing scales and surrounded by brown fibres. Fl. paniced. Pet. purple without, white within. Seed-stalks thick, white. Seeds black.—Sandy heaths. Near Bournemouth, Dorset. Near Derrynane, Kerry. P. V. E. I.

Tribe IV. *Hemerocallideæ*.10. *ENDYM'ION* Dumort. Blue-bell.

1. *E. nitans* (Dum.); l. linear, raceme nodding, fl. bellshaped cylindrical, tip of the sep. revolute, bracts 2.—*Scilla* Sm., *E. B.* 377. *Agraphis* Link.—Scape about a foot high. Fl. blue, rarely white. Stam. united to the perianth halfway up. L. shorter than the scape.—Woods and thickets. P. V. *English Blue-bell.* E. S. I.

11. *MUS'CARI* Tourn. Grape-Hyacinth.

1. *M. racemosum* (Mill.); fl. ovoid nodding crowded upper ones nearly sessile abortive, l. linear channelled flaccid recurved at the end.—*E. B.* 1931.—Scape 1 ft. high. Fl. dark blue, scented. Caps. emarginate. Not *M. neglectum* (Guss.), as supposed in ed. 6. The cultivated plant, *M. botryoides*, has globose flowers.—Sandy fields. Plentiful near Cavenham and Pakenham, Suffolk. About Gogmagog Hills, Cambridge. P. IV. V. E.

## Order LXXXIX. MELANTHACEÆ.

Perianth inferior, 6—7-parted. Stam. 6, on the receptacle or perianth. Anth. attached below their middle. Ovaries superior, 1 of 3 cells, or 3 of 1 cell more or less connected. Ovules many. Styles 1—3. Fr. bursting inwards, of 3 separate 1-celled follicles, or more or less combined into a 3-celled capsule.

1. *COLCHICUM*. Perianth funnelshaped, tube long; limb 6-parted, petal-like. Styles long. Caps. 3, connected throughout, opening at the inner edge, many-seeded.—Tuberous.
2. *TOFIELDIA*. Perianth 6-leaved. Styles short. Caps. 3, connected to above the middle, 1-celled, opening at the inner edge, many-seeded.—A creeping rhizome.

3. NARTHECIUM. Perianth 6-parted. Style undivided. Caps. 3-celled, loculicidal, many-seeded; seed with a long filiform appendage at each end; placenta basal.

1. COL'CHICUM *Linn.* Meadow-Saffron.

1. *C. autumnale* (L.); l. flat lanceolate erect.—*E. B.* 133.—Root-stock large, tuberous. L. a foot long and often an inch broad, dark green, smooth, vernal. Fl. bright purple, radical, with very long tubes, autumnal; germen remaining under ground in winter, rises in spring with the leaves. Rarely greenish spring fl. are found.—Meadows. P. IX. X. E. S.

2. TOFIEL'DIA *Huds.* Scottish Asphodel.

1. *T. palustris* (Huds.); pedicels with a 3-lobed bract at the base but none at the top.—*E. B.* 536. *T. borealis* Wahl. *St.* 78. 8.—*St.* 4—8 in. high. L. swordshaped, about 2 in. long, in 2-ranked radical tufts. Fl. in a short dense spike, at first sessile, afterwards slightly stalked. Mountain bogs. P. VII. E. S.

3. NARTHECIUM *Huds.* Bog-Asphodel.

1. *N. ossifragum* (Huds.); l. swordshaped, pedicels with 1 bract at their base and another above their middle, perianth-segm. linear-oblong exceeding stam. and much falling short of caps.—*E. B.* 535. *St.* 78. 3. *R. x.* 421.—*St.* 6—8 in. high, slightly leafy, decumbent and rooting below. L. mostly in radical 2-ranked tufts, half the height of the stem. Fl. bright yellow.—Turfy bogs. P. VI. VII. E. S. I.

Order XC. JUNCACEÆ.

Perianth of 6 glume-like scarious segments. Stam. usually 6, on the base of the segments; or 3, opposite to the outer series. Anth. 2-celled, attached by base. Ovary 1—3-celled, superior. Ovules 1, 3, or many in each cell. Style 1, stigmas 3. Fr. capsular, 3-valved, loculicidal. Embryo subcylindrical, very minute, within firm albumen, near the hile.—No raphides.

1. JUNCUS. *Perianth glume-like, 6-leaved. Filaments glabrous.* Style undivided. Stigmas 3, filiform. Caps. 3-celled, 3-valved. Seeds attached to the inner edge of the dissepiments.—L. mostly not flat.
2. LUZULA. Caps. 1-celled, 3-valved, with dissepiments. Seeds 3, at the base of the capsule. Otherwise like *Juncus*.—L. flat, grass-like.



## 1. JUN'CUS Linn. Rush.

\* *Barren and fertile stems subulate, with sheathing radical long leaves like the stem, or mucronate sheaths. Seeds with a loose testa forming a sack at each end (appendaged).*

1. *J. marit'imus* (Sm.); sheaths short pale, l. terete sharp-pointed, panicle compound erect, *per.-segm.* equal lanceolate acute *equalling the elliptic mucronate capsule*, seeds fusiform.—*E. B.* 1725. *R.* ix. 402.—St. erect, 1—2 feet high. Panicle long, lax.—Salt marshes near the sea. P. VII. VIII. E. S. I.

2. *J. acútus* (L.); sheaths long shining, l. terete sharp-pointed, panicle very compound mostly compact, *per.-segm.* equal *half the length of the roundish ovoid caps.*, 3 inner ones blunt with a membranous border, seeds broad-ovate.—*E. B.* 1614. *St.* 71. 6. *R.* ix. 401.—St. erect, rigid, with a very sharp rigid point, 3—6 feet high. Panicle dense, corymbose. Fr. twice as large as that of Sp. 1.—Sands on the sea-coast, rare. P. VI. E. I.

\*\* *Barren and fertile stems subulate with sheaths at their bases which are either leafless or bear rudimentary leaves. Seeds with a close testa (not appendaged).*

3. *J. effúsus* (L.); *st. faintly striate soft*, pith continuous, panicle close or diffuse, *caps.* obovate retuse *not mucronate*, *stam.* 3.—*E. B.* 836. *R.* ix. 413.—Height 1—2 feet. L. none or minute and slender at the top of sheathing scales. Panicle diffuse, branched; or more or less dense, globose. Anth. oval, short.—Marshy ground. P. VII. E. S. I.

4. *J. conglomerátus* (L.); *st. faintly striate soft*, pith continuous, panicle close or diffuse, *caps.* obovate retuse *mucronate*, *stam.* 3.—*E. B.* 835. *St.* 71. 3. *R.* ix. 408.—Height 1—2 feet. L. none or minute and slender at the top of the sheathing scales. Panicle globose, dense; or more or less diffuse. Anth. linear. The mucro in the hollowed top of the caps. resembles a little hill bearing the style. Marshy ground. P. VII. E. S. I.

5. *J. glaucus* (Sibth.); *st. deeply striate rigid*, pith interrupted, panicle loose much branched erect, *per.-segm.* lanceolate-subulate rather exceeding the *elliptic-oblong mucronate capsule*, *stam.* 6.—*E. B.* 665. *St.* 71. 5. *R.* ix. 415.—Panicle ascending, diffuse. Fr. black. St. rigid, slender, glaucous, 12—18 in. high. Sheaths dark. L. none or minute and slender at the top of the scales.—Wet places. P. VII. E. S. I.

6. *J. diffúsus* (Hoppe); *st. finely striate rigid*, pith continuous, panicle loose much branched erect, *per.-segm.* lanceolate-subu-

late exceeding the *obovate-blunt mucronate capsule*, *stam.* 6.—*Sy. E. B.* 1562. *St.* 77. 10. *R.* ix. 414.—Very like the preceding, but the caps. much smaller. *St.* green.—In wet places, rare. *P.* VII. VIII. E. S. I.

7. *J. balticus* (Willd.); *st.* very faintly striate rigid, pith continuous, panicle erect slightly branched, per.-segm. ovate-lanceolate acute, caps. elliptic scarcely trigonous blunt mucronate.—*E. B. S.* 2621. *St.* 71. 2. *R.* ix. 411.—Creeping widely. *L.* none or very minute points at the top of the sheathing scales.—Distinguished from *J. articus* by its rounded not trigonous capsules; from *J. glaucus* by its far-creeping rhizome, scarcely striate *st.* and continuous pith.—Sandy and wet sea-coasts of Scotland. *P.* VII. S.

8. *J. filiformis* (L.); *st.* filiform faintly striate, panicle simple of few (about 7) fl. placed near the middle of the *st.*, per.-segm. lanceolate acute, caps. roundish obovate blunt mucronate.—*E. B.* 1175. *St.* 36. 10. *R.* ix. 412.—*L.* none or as in the preceding plants. *St.* remarkably slender, about 1 ft. high; small panicles placed very low. Rhizome creeping.—Stony margins of lakes in the North. *P.* VII. E. S.

\*\*\* *Stems mostly leafy, none barren.*

† *Fl.* capitate or solitary and terminal. Seeds appendaged.

9. *J. castaneus* (Sm.); *st.* with 2—3 channelled *l.*, cymes terminal solitary or 2 or 3, per.-segm. elliptic-lanceolate acute half as long as the ovate-oblong pointed trigonous capsules, creeping.—*E. B.* 900. *St.* 71. 14. *R.* ix. 393.—*St.* 8—12 in. high. Root with lax runners. Leaflike bract exceeding the flowers. Caps. chocolate-coloured, about  $\frac{1}{4}$  inch long. Filaments about twice as long as the anthers.—Micaceous mountain bogs at a great elevation, rare. *P.* VII. VIII. S.

10. *J. triglumis* (L.); *st.* leafless round, *l.* radical subulate channelled bitubular, 2 or 3 fl. terminal erect usually equalling the membranous bract, per.-segm. elliptic-oblong blunt falling rather short of the ovate-oblong blunt mucronate caps., caespitose.—*E. B.* 899. *St.* 28. 2. *R.* ix. 392.—*St.* several from one root, 3—6 in. high, perfectly round. Leaflike bract equalling or falling short of the flowers. Caps. chestnut-coloured.—Boggy places on mountains. *P.* VII. VIII. E. S.

11. *J. biglumis* (L.); *st.* leafless channelled on one side, *l.* radical subulate compressed (not channelled, nor bitubular), fl. 2 unilateral upper which is stalked usually falls short of the leaflike bract, per.-segm. oblong blunt falling rather short of the turbi-

*nate retuse trigonous caps.*, root fibrous.—*E. B.* 898.—St. 2—4 in. high, seldom more than one from each root. Caps. light brown with purple margins.—Boggy spots on mountains, rare. P. VIII. S.

12. *J. trif'idus* (L.); *st. with 1 leaf* on its upper part, basal sheaths awned, upper sheath with a short l., fl. 1—3 *with two setaceous leaflike bracts*, per.-segm. acute, falling short of the rounded elliptic beaked caps., creeping.—*E. B.* 1482. St. 71. 12. *R.* ix. 594.—St. crowded, erect, slender, 2—6 in. high. Occasionally the stem-l. is wanting, and sometimes it has a second head in its axil. Perianth and caps. dark brown.—Damp rocky places on mountains. P. VII. VIII. S.

†† Flowers in 1 terminal cluster or 2 one above the other, or in paniced clusters. Seeds not appendaged.

13. *J. capitatus* (Weigel); *st. naked* erect simple, l. radical filiform, head terminal mostly solitary falling short of the setaceous bract, per.-segm. unequal, outer ovate-lanceolate, *acuminate-aristate twice as long as the truncate apiculate caps.*, *stam.* 3.—*E. B. S.* 2644.—Plant 1—4 in. high. L. half as long as the stems. Heads large, of 3—10 sessile flowers.—Land's End and Lizard district, Corn.; Guernsey and Jersey. A. VI. VII. E.

14. *J. obtusiflorus* (Ehrh.); 1-leaved *st. and internally jointed l. terete*, panicle repeatedly compound spreading divaricate, per.-segm. equal blunt equalling the ovate acute trigonous (pale brown) capsule.—*E. B.* 2144. *R.* ix. 404.—Erect, 2—3 feet high. St. and l. not compressed. Segm. of perianth pale, often purplish, quite blunt or with a small inflexed point.—Marshes, rare. P. VII.—IX. E. S. I.

15. *J. acutiflorus* (Ehrh.); 3—4-leaved *st. and internally jointed l. subcompressed*, panicle repeatedly compound pyramidal, per.-segm. acute inner ones longest all falling rather short of the narrow ovate-acuminate rostrate triquetrous (pale brown) capsule.—*E. B.* 238. *R.* ix. 406.—St. erect, 1½—2 feet high. L. slightly compressed. Clusters 5—6-flowered.—Boggy places. P. VI.—VIII. E. S. I.

16. *J. lamprocar'pus* (Ehrh.); 3—6-leaved *st. and internally jointed l. compressed*, panicle repeatedly compound erect forked, per.-segm. equal acute the inner ones blunt all falling short of the ovate-attenuate mucronate triquetrous (dark brown) capsule.—*E. B.* 2143. *R.* ix. 405.—St. erect, 12—18 in. high. L. compressed; many internal transverse divisions. Clusters 4—8-flowered.—Boggy places. P. VII. VIII. E. S. I.

[*J. alpinus* (Vill.), *R.* ix. 403, having an acute dorsal angle to the sheaths of its l. and blunt per.-segm. mucronate below their

summit,—*J. atrátus* (Krock.), *R.* ix. 407, having “more deeply striate l.” than either of the two preceding species and the inner acute perianth-segm. about equalling the ovate-attenuate capsule,—and *J. nig'ricans* (Drej.), *R.* ix. 407, with unequal acute perianth-segm. and a subglobose capsule will perhaps be found.]

17. *J. nigrítellus* (D. Don); 3—4-leaved *st.* and internally joined *l.* nearly cylindrical, panicle slightly compound erect, *per.-segm.* nearly equal (3 inner rather longer and broader) all acute falling short of the linear-oblong trigonous beaked (black) capsule.—*E. B.* S. 2643, not of *Koch.*—*St.* erect, 6—12 in. high. *L.* scarcely compressed. Clusters of more fl. than in *Sp.* 16. Caps. brown, at length black and glossy, more abruptly pointed than in *J. lamprocarpus.*—Boggy places in the north. P. VII. VIII. E. S. I.

18. *J. supínus* (Moench); *st.* filiform, *l.* setaceous slightly channelled faintly jointed internally, panicle nearly simple irregular long with few distant clusters, *per.-segm.* equal acute (3 inner rather blunt) nearly equalling the oblong very blunt mucronate (pale brown) capsule, anth. as long as their filaments.—*E. B.* 801. *R.* ix. 397. *J. uliginosus* and *J. subverticillatus* Sm.—Extremely variable in size and the direction of its stems, sometimes erect, at others prostrate and rooting at every joint, or floating. Fl. often viviparous. Stam. 3 or 6.— $\beta$ . *Kochii*; caps. shorter, *stam.* 6, filaments nearly twice as long as the elliptic anthers. *J. nigrítellus* Koch. *St.* 78. 2.—Boggy and wet places.  $\beta$ . Ivy Bridge, Devon (in a bog). Connemara, Galway. P. VI.—VIII. E. S. I.

††† Flowers solitary, remote, or corymbose and forming a terminal panicle. Seeds not appendaged. \*

19. *J. squarrósus* (L.); *st.* leafless simple, *l.* linear channelled radical, panicle terminal compound with cymose branches, *per.-segm.* ovate-lanceolate acute or bluntish equalling the obovate blunt mucronate capsule, anth. 4 times as long as their filaments.—*E. B.* 933. *St.* 36. 11. *R.* ix. 400.—*St.* erect, 6—12 in. high. *L.* many, somewhat spreading, rigid, half as long as the stem. Caps. pale brown, shining.—Wet heaths and moors. P. VI. VII. E. S. I.

20. *J. compres'sus* (Jacq.); *st.* with 1 leaf in the middle, *l.* linear channelled, panicle terminal compound subcymose usually falling short of the bract, *per.-segm.* oval-oblong blunt falling short of the roundly ovate shortly mucronate capsule.—*E. B.* 934. *St.* 36. 13. *R.* ix. 399.—*St.* slender, erect, round and leafy below,

naked and compressed above. Floral bracts usually pale. Style half the length of the ovary. Anth. oblong.—Damp places. P. VI.—VIII. E. S. I.

21. *J. Gerar'di* (Lois.); st. with 1 or more leaves, l. linear channelled, panicle terminal compound subcymose usually exceeding the bract, *per.-segm.* oval-oblong blunt *about equalling the oval-oblong strongly mucronate capsule.*—*J. cœnosus* Bich., Sm. *E. B. S.* 2680. *St.* 7l. 8. *R.* ix. 398.—St. trigonous in its upper part. Floral bracts usually shining, brown. Style as long as the ovary. Anth. long.—Salt marshes. P. VI.—VIII. *Mud-Rush.* E. S. I.

22. *J. bufónius* (L.); st. leafy forked, l. setaceous, fl. solitary unilateral scattered mostly sessile, *per.-segm.* unequal lanceolate-acuminate exceeding the *oblong blunt capsule, seeds roundly oval.*—*E. B.* 802. *St.* 36. 12. *R.* ix. 395.—St. 4—8 in. high. Usually with only 1 leaf on the slender stems— $\beta$ . *J. fasciculatus* (Bert.); st. shorter (2—3 in. high) and thicker, fl. 2 or 3 together.—Marshy and wet places. A. VII. VIII. *Toad-Rush.* E. S. I.

23. *J. pygmæ'us* (Rich.); st. nearly or quite leafless, l. subsetaceous, fl. in few small clusters sessile, *per.-segm.* nearly equal narrowly lanceolate exceeding the oblong-acute caps., seeds fusiform-pear-shaped.—*J. of B.* xi. 128.—St. very short. Upper cluster of fl. stalked, naked; lower with a long bract. Seeds much longer than in Sp. 22, apiculate.—Damp spots above Kynance Cove, Cornwall. A. V.  $\rho$  E.

## 2. LU'ZULA Cand. Wood-Rush.

1. *L. sylvat'ica* (Bichen.); l. linear-lanceolate hairy, *panicle subcymose doubly compound,* clusters about 3-flowered on long stalks, *per.-segm.* bristle-pointed equalling the ovoid-mucronate capsule, filaments very short, seeds minutely tubercled at the end.—*E. B.* 737. *St.* 36. 14. *R.* ix. 390. *L. maxima* DC., Koch.—Rhizome woody. *St.* 12—18 in. high. L. broad, shining, striate, with hairy edges. Panicle much exceeding the leaflike bracts.—Sandy places. P. IV.—VI. E. S. I.

2. *L. Fors'teri* (DC.); l. linear hairy, panicle subcymose only slightly branched, *ped.* 1-flowered *erect* with both fl. and fr., style equalling stam., filaments about as long as the anth., *caps. acute* scarcely falling short of perianth, *seeds with a straight blunt crest.*—*E. B.* 2193. *St.* 77. 2.—St. slender about a foot high. Caps. with 3 acute angles, not suddenly contracted above.—Thickets, rather rare. P. V. E.

3. *L. pilósa* (Willd.); l. lanceolate hairy, panicle subcymose only slightly branched, *peduncles* 1—3-fl., upper ones reflexed after flowering, fl. solitary, style (excluding stigmas) exceeding stam., filaments about half as long as the anth., *caps. blunt* exceeding the perianth, *seeds with a falcate crest*.—*E. B.* 736. *St.* 77. 3.—*St.* slender, 6—12 in. high. *Caps.* ovoid, trigonous, suddenly contracted above.—*L. Borreri* (Bromf.) is a sterile form with shorter and acuter capsules.—Thickets. P. V. E. S. I.

4. *L. campes'tris* (Willd.); l. linear hairy, *panicle of 3 or 4* ovate dense *sessile or stalked clusters*, per.-segm. lanceolate-acuminate exceeding the blunt apiculate caps., *filaments much shorter than the anthers*, *seeds nearly globular with a basal appendage*.—*E. B.* 672. *St.* 77. 5.—*St.* 4—6 in. high. *Anth.* linear, about 6 times as long as the filaments. Large forms often mistaken for *Sp. 5.*—Pastures and dry places. P. IV. V. E. S. I.

5. *L. multiflóra* (Lej.); l. linear hairy, panicle of many ovate dense sessile or stalked clusters, per.-segm. narrowly lanceolate strongly acuminate exceeding the blunt apiculate caps., *filaments more than  $\frac{1}{2}$  as long as the anthers*, *seeds nearly twice as long as broad with a basal appendage*.—*E. B. S.* 2718. *St.* 77. 7. *L. congesta* Sm.—*St.* 8—20 in. high. *Anth.* small, rather short. Clusters on elongate drooping stalks, or (*L. congesta* Lej.) subsessile in a rounded lobed head, or a few stalked.—Wet and turfy places. P. VI. E. S. I.

6. *L. spicáta* (DC.); l. narrow slightly channelled hairy, *panicle oblong lobed nodding spikelike*, clusters falling short of their bracts, per.-segm. narrow acuminate bristle-pointed, filaments half as long as the anthers, caps. blunt apiculate, seeds oblong with a very slight basal appendage.—*E. B.* 1176.—*St.* 3—12 in. high. L. short, slender. Spike  $\frac{1}{2}$ —1 in. long, nodding. Partial bracts tapering, bristle-pointed.—Mountains. P. VII. E. S.

7. *L. arcuáta* (Hook.); l. channelled slightly hairy, *panicle subumbellate of few 3—5-flowered clusters on long drooping peduncles*, per.-segm. broadly-lanceolate bristle-pointed, filaments as long as the anthers, caps. roundish-ovate, seeds oblong blunt or apiculate scarcely appendaged below.—*E. B. S.* 2688.—*St.* slender, 2—5 in. long. L. short, curved, narrowly linear. Panicle of 3—5 small clusters, one nearly sessile, the others on long deflexed stalks.—Highest summits of the Cairngorum and Sutherland Mountains. P. VII. ? S.

## Order XCI. ERIOCAULACEÆ.

Fl. capitate, unisexual. Perianth very delicate, 2—6-parted. Stam. 2—6, if in 2 rows the inner row most developed. Anth. 2-celled. Ovary superior, 2—3-celled. Ovules solitary, pendulous. Dehiscence of caps. loculicidal. Seeds coated with wings or rows of hairs. Embryo lenticular, on the outside of farinaceous albumen, at the end remote from the hile.

1. ERIOCAULON. Fl. in a compact scaly head. Barren fl. in the centre. Perianth 4—6-fid, the inner segments united nearly to their top. Stam. 4—6.—Fertile fl. in the circumference. Perianth deeply 4-parted. Stigmas 2—3. Caps. 2—3-lobed, 2—3-celled; cells 1-seeded.

## 1. ERIOCAU'LON Linn. Pipewort.

1. *E. septanguläre* (With.); scapes 6—8-striate exceeding the cellular compressed subulate glabrous l. fl. 4-cleft hairy at the end as well as the scales, stam. 4, caps. 2-cel'ed—*E. B.* 733.—Roots of many white jointed fibres. St. varying in height according to the depth of the water. Fertile fl. 4-parted nearly to the base; 2 lateral divisions keeled, compressed, blunt, fringed, black. Each fl. with a broad blunt black scale in front which is shorter and broader than it.—Hebrides, especially Skye. West coast of Ireland, especially Connemara. P. VIII. S. I.

## Order XCII. TYPHACEÆ.

Fl. monœcious, many, closely placed in cylindrical spikes or in dense globose clusters; barren and fertile on different parts of spike, the males uppermost. Perianth of 3 or more scales or hairs. Stam. 1—6, distinct or monadelphous. Anth. erect.—Fertile fl. Ovary free, solitary, 1-celled; ovule 1, pendulous. Style simple. Stigma unilateral. Fr. dry or spongy. Embryo with a cleft on one side.—Raphides abundant.

1. TYPHA. Spikes long dense cylindrical, upper part male, lower female. Stam. surrounded with hairs. Anth. 3 together on one filament. Ovary surrounded by hairs ultimately stalked.
2. SPARGANIUM. Fl. with a single 3—4-leaved perianth, in distant dense globose heads, the lower bracteate. Stam. free. Fruit dry, sessile.

## 1. TY'PHA Linn. Reed-mace.

1. *T. latifolia* (L.); l. linear nearly flat, *sterile and fertile parts of spikes not separated*, style exceeding the bristles, stig. oblique ovate-lanceolate.—*E. B.* 1455. *R.* ix. 323.—St. 6—7 feet high. L. overtopping the inflorescence, very broad. Spikes very long, sometimes there is a very short space between them; fertile blackish brown.—[In *T. Shuttleworthii* (Koch), *R.* ix. 322, the style equals the bristles.]—Ponds and lakes. P. VI. VII. E. S. I.

2. *T. angustifolia* (L.); l. linear channelled below, *sterile and fertile parts of spikes a little separated*, style exceeding the bristles, stig. long filiform.—*E. B.* 1456. *R.* ix. 321.—St. 5—6 feet high, much slenderer than in the preceding. L. very narrow overtopping the inflorescence. Spike very long, slender, separated by a flowerless interval of about an inch; fertile reddish brown.—[*T. gracilis* (Suhr.), *R.* ix. 320, has a rounded spatulate stigma.]—Lakes and ponds. P. VI. VII. E. S. I.

## 2. SPARGA'NIUM Linn. Bur-reed.

1. *S. ramosum* (Huds.); l. triquetrous at the base with concave sides, *st. branched above*, stigma linear, fl. sessile.—*E. B.* 744. *R.* ix. 326.—St. erect, about 2 feet high; lower branches with several heads, 1—3 fertile, the rest barren. Heads spherical. L. long, linear, erect.—Ditches. P. VI. VII. E. S. I.

2. *S. simplex* (Huds.); l. trigonous at the base, *st. simple*, stigma linear-subulate, fr. slightly stalked subfusiform.—*E. B.* 745. *R.* ix. 325.—St. 1—2 ft. high, long, erect or ascending. L. long, often floating (and then often mistaken for *S. natans*); sheath slightly furrowed, not inflated. Heads many; barren several, sessile; fertile shortly stalked, especially the lowest. Fr. oblong, narrowed into a long beak, elliptic-fusiform.—Ditches. P. VII. E. S. I.

3. *S. natans* (L.); st. simple flaccid, floating l. very long linear flat at the base from a dilated sheathing base, heads many distant, lower fertile heads stalked, *male heads* several sessile, stig. linear-lanceolate, *fr. stalked oblong not longer than its subulate beak*.—*S. affine* Schn. *Sy. E. B.* 1389. *R.* ix. 417.—St. much thicker than in the next plant. L. grass-green, very long. Male heads fewer in our plant than in the Swedish.—Lakes, rare. P. VIII. E. S. I.

4. *S. minimum* (Fr.); st. simple flaccid, l. linear floating



blunt not dilated at the base, heads few racemose or spiked distant, *only 1 male head*, stig. short oblong-lanceolate, *fr. sessile ovoid shortly beaked*.—*S. natans* Sm., *E. B.* 273. *R.* ix. 324.—St. slender. L. pale, pellucid, long. Lowest fertile heads sometimes very shortly stalked. [Often 2 male heads in Ireland (A. G. More). *S. fluitans* Fr.?]—Lakes and ditches. P. VII. VIII. E. S. I.

## Order XCIII. ARACEÆ.

Fl. monœcious or perfect, placed on a spadix and often in a spath. Barren and fertile fl. usually on different parts of the spadix. Perianth none, or of 4—8 scales. Stam. many or definite. Anth. turned outwards. Ovary free, with 1 or more cells. Stigma sessile. Fr. succulent. Embryo slit on one side, —*Arum* has abundant raphides, *Acorus* none.

Tribe I. *ORONTEÆ*. Spath like a continuation of stem. Fl. perfect.

1. *ACORUS*. Fl. on a sessile spadix, appearing lateral. Perianth 6-leaved, inferior, persistent. Stam. 6, filiform.

Tr. II. *AREÆ*. Spath convolute at the base. Fl. monœcious. Perianth 0.

2. *ARUM*. Perianth 0. Male fl. of 1 sessile 2-celled anther. Fem. fl. placed lower, of 1 pistil. Top of spadix naked.

1. *Ac'ORUS*. Sweet Flag.

1. *A. Cal'amus* (L.); st. with a long leaflike prolongation (or spath) beyond the spadix.—*E. B.* 356. *R.* x. 429.—St. 5—6 feet high, resembling the l., swordshaped, flattened. Spadix completely covered by the flowers, 2—3 in. long, lateral. St. and l. sweet-scented when crushed.—In water, rare; except in Norf. and Suff. \*Between Lisburn and Moira, Co. Down.

E. I.

2. *A'RUM* Linn. Cuckoo-pint.

1. *A. maculátum* (L.); *l. vernal* all radical hastate-sagittate with *deflexed lobes*, petiole as long as leaf-limb, spadix club-shaped straight falling short of the spath.—*E. B.* 1298. *R.* vii. 8.—Root tuberous. L. with branching veins, green or spotted with purple. Spath ventricose below and above, constricted in the middle, with inflexed edges when open. *Spadix usually purple* blunt; with ovaries at the base; above them whorls of

stamens; then a few filaments, probably abortive pistils; club naked. Berries scarlet remaining after the rest of the plant has disappeared; seeds mostly 4 or 3, or rarely 2.—Hedge-banks and thickets. P. IV. V. E. S. I.

2. *A. italicum* (Mill.); *l. appearing before the winter* all radical triangular-hastate with *divaricate lobes*, petiole longer than leaf-limb, spadix clubshaped straight falling short of the widely spread spath.—*R.* vii. 11.—*L.* dark blue-green, sometimes with yellowish veins, rarely spotted, blunt. *Spath* ventricose below, opening nearly flat and very broad above, *folding down in front* when fl. are in perfection so as to close the opening like a flap, ultimately bent or folded double over the *yellow spadix*. Abortive pistils very long, both above and below the stamens.—Undercliff, Isle of Wight; Channel Islands; and W. Cornwall. E.

#### Order XCIV. LEMNACEÆ.

Fl. monœcious, 2, in a spath, but without a spadix (rarely found). Perianth 0. Stam. 1—2, distinct. Ovary 1-celled. Style short. Stigma simple. Fr. bladderly, not bursting. Seeds with a coriaceous ribbed testa.—Floating, leaflike, small, proliferous; no distinction of st. or leaf. Fl. very minute.

1. LEMNA. Spath membranous, urnshaped. Fl. from lateral clefts of frond. Stam. 1—2. Anth. 2-celled, didymous (cells bilocular?). Fronds with capillary roots beneath.—Increasing chiefly by offsets.
2. WOLFFIA. Spath 0. Fl. from upper surface of frond. Stam. 1; anth. 1-celled, sessile. Frond very minute, rootless.—Increasing by offsets.

#### 1. LEMNA Linn. Duckweed.

1. *L. trisul'ca* (L.); *fronds* thin pellucid elliptic-lanceolate tailed at one end serrate at the other, roots solitary.—*E. B.* 926. *R.* vii. 15.—Fronds half an inch long, proliferous at right angles. Plants submerged, truly annual, producing autumnal bulblets which survive the winter as in the other species.—In stagnant water. A. VI. E. S. I.

2. *L. minor* (L.); *fronds* obovate compressed opaque, roots solitary blunt.—*E. B.* 1095. *R.* viii. 14.—Fronds 1—2 lines long, nearly flat beneath, of a compact texture.—On stagnant water. A. VI. VII. E. S. I.

3. *L. gib'ba* (L.); *fronds* obovate nearly flat above hemispheri-

cal and spongy beneath, roots solitary blunt.—*E. B.* 1233. *R.* vii. 14. *Telmatophace* Schleid., Endl.—Fronds 1—2 lines long, at first flattish, afterwards remarkably gibbous and cellular beneath.—On stagnant water. *A.* VI.—VIII. *E. S. I.*

4. *L. polyrrhiza* (L.); fronds roundish-obovate, compressed, roots many clustered acute.—*E. B.* 2458. *R.* vii. 15, *Spirodela* Schleid., Endl.—Fronds half an in. long, green above, purple beneath. Fl. not seen in Britain.—On water. *A.* *E. S. I.*

## 2. *WOLFFIA* *Horkel.*

1. *W. arrhiza* (Wimm.); fronds very small subglobular flattish above cellular beneath, solitary, young frond separating immediately from the old one.—*Sy. E. B.* 1398. *R.* vii. 14.—Frond like a grain of sand, subglobular at all ages, green. Offset from within the base of the old frond.—Ponds near London. *A.* *E.*

## Order XCV. POTAMOGETONACEÆ.

Fl. perfect or imperfect. Perianth inferior, 4-parted, or 0. Stam. free, 1, 2, or 4. Ovaries 4, distinct, each with 1 ovule and 1 sessile stigma. Fr. a drupe enclosing a hard nut, or a dry nut, not bursting, 1-seeded. Albumen 0. Embryo with a thin skin having a lateral cleft.

1. POTAMOGETON. Fl. perfect. Perianth 4-parted. Anth. 4, sessile, opposite to the divisions of the perianth. Ovaries 4. Styles 0. Drupes 4, sessile.—Fl. sessile, spiked.
2. RUPPIA. Fl. perfect. Perianth 0. Stam. 2, the cells considerably separated; filaments very short, scalelike. Ovaries 4. Styles 0. Nuts 4, with long stalks.—Fl. about 2 together.
3. ZANNICHELLIA. Fl. monœcious, axillary. Barren with 1 stam., and no perianth. Fertile with a bellshaped perianth, persistent style, and peltate stigma. Nuts 2—5 or more, more or less stalked.

### 1. POTAMOGETON *Linn.* Pondweed.

\* *L. alternate*, floral l. floating and sometimes opposite; stipules free.

1. *P. natans* (L.); l. all stalked, upper coriaceous floating ovate or elliptic folded at the base, lower leafless petioles linear or setaceous, fr. (large) rounded on the back when fresh keeled when dry, peduncle equal.—*E. B.* 1822. *R.* vii. 50.—St. creep-

ing below, simple. *Petioles* plane-concave. L. subcordate below, when pressed flat a ridge is formed on each side of the base, *jointed to their stalks a little below the limb*. Sep. stalked, roundly rhomboidal. Anth.-cells not parallel. *Fr. greenish*, slightly compressed,  $1\frac{1}{2}$ —2 lin. long.—Ponds, ditches, and slow streams. P. VI. VII. E. S. I.

2. *P. polygonifolius* (Pourr.); l. all stalked, upper subcoriaceous floating oblong-elliptic subcuspidate, *no leafless petioles*, lower l. linear-lanceolate, *fr. minute blunt and rounded on the back*, peduncle equal.—*P. oblongus* Viv., *E. B. S.* 2849.—St. creeping below. *Petioles* longer than leaves, convex on both sides, *not jointed below the limb*. Lower l. often very narrow. Spikes rather short and irregular. Sep. transversely elliptic, stalked. *Fr. reddish*, scarcely compressed, 1 lin. long; a faint keel and lateral ridges when dry.— $\beta$ . *pseudo-fluitans* (Syme); submerged l. membranous linear-lanceolate narrowed at both ends.—Ditches, small streams, and ponds.  $\beta$ . Buttermere. Gap of Dunloe, Kerry. P. VII. E. S. I.

3. *P. plantagin'cus* (Ducr.); l. all shortly stalked membranous and pellucid blunt not cuspidate nor plicate, upper elliptic, lower l. oblong, *fr. minute rounded on the back when fresh keeled when dry*, peduncle equal.—*E. B. S.* 2848. *R.* vii. 45.—St. creeping below, branched, sometimes throwing out long runners from its upper axils. L. all beautifully transparent and netted with veins, the upper often almost sessile and nearly orbicular, sometimes slightly cuspidate. *Petioles* plane-concave above. Spikes long, cylindrical. Anth.-cells nearly parallel. Sep. ovate. *Fr. greenish*,  $\frac{3}{4}$  lin. long.—Stagnant peaty water. P. VI. VII. E. S. I.

4. *P. rufes'cens* (Schrad.); *submersed l.* lanceolate narrowed at both ends *sessile* membranous pellucid *entire not apiculate* with chainlike network near the midrib, floating l. subcoriaceous, obovate blunt narrowed into a short petiole, stip. without wings, *fr. acutely keeled*, peduncle equal.—*E. B.* 1286. *R.* vii. 32. *P. fluitans* Sm.—St. simple. Upper l. alone slightly coriaceous, often tinged with purple, longer than their stalks; submerged l. all nearly, if not quite, sessile. Sep. "transversely oval."—Ditches and slow streams. P. VII. E. S. I.

5. *P. lanceolatus* (Sm.); *submersed l.* linear-lanceolate sessile *entire acute not apiculate with chainlike network* near the midrib, floating l. subcoriaceous elliptic-lanceolate shortly stalked, stip. almost subulate uppermost broader with 2 dorsal elevated ribs, *peduncles equal* from uppermost floating opposite leaves.—*E. B.* 1985. *J. of B.* xix. t. 217 (not *R.*).—St. very slender, slightly

branched. Floating l. not always present with the flowers. No fruit even on Sept. 28.—By the bridge at Penrhos Lligwy, Anglesea. Burwell Fen, Cambr. P. VII.—IX. E.

6. *P. sparganiifolius* (Laest. !); *submersed l. linear* narrowed at both ends *very long* sessile entire not apiculate *with many parallel veins next the midrib*, floating l. subcoriaceous lanceolate long-stalked, stip. very long blunt not winged, fr. "acutely keeled" (Fr.), ped. rather slender not clavate.—*Sy. E. B.* 1403.—St. very slender, slightly branched. Floating l. often wanting. Submersed l. sometimes 2 ft. long, very narrow, few-veined with additional parallel veins closely placed next the midrib. L. very much longer than those of *P. lanceolatus* and without the chainlike network, beautifully green when well dried. Exactly the plant of *Fries, H. N.* xii. 75.—In the river at Ma'am, Co. Galway. I.

7. *P. salicifolius* (Wolfg.); *submersed l. narrowly strapshaped* sessile narrowed at both ends entire not apiculate, floating l. subcoriaceous elliptic long-stalked, stip. long bluntish not winged, fr. obliquely obovate 3-keeled.—*P. lonchites* Tuck. *Sy. E. B.* 1404.—St. slightly branched. Submersed l. with a few additional parallel veins next midvein, subsessile. Floating l. often wanting. Ped. long, slightly thickened upwards.—In river Boyne near Navan. P. I.

8. *P. heterophyllus* (Schreb.); *submersed l. lanceolate* narrowed at both ends sessile minutely denticulate and apiculate, floating l. subcoriaceous elliptic stalked, stip. broadly lanceolate blunt *with 2 stout prominent ribs*, lower ones linear-lanceolate, fr. blunt on the back, peduncle thicker than the stem swelling upwards.—*E. B.* 1285. *R.* vii. 41—43. *P. gramineus* Fries, not *Linn.*—St. much branched below. Submersed l. wavy. Lower stip. without the two strong ribs and equally veined, upper ones widely spreading. Dry fr. slightly marked with 3 ridges on the back.—Ponds and ditches. P. VI. VII. E. S. I.

9. *P. nitens* (Web.); *submersed l. lanceolate rounded below and half clasping* wavy at the edge, floating l. "coriaceous elliptic stalked," stipules equally veined, (dry) fr. keeled, ped. swelling upwards.—*R.* vii. 34. *Sy. E. B.* 1407. *J. of B.* ii. t. 23.—St. rather wavy, branched below. Subm. l. pellucid, often recurved; upper l. often not coriaceous. Upper stip. large, persistent, nearly  $\frac{1}{2}$  as long as the peduncle. Dry fr. with 3 ridges on the back.—Lakes near Brandon Mountain, Kerry (*P. curvifolius Hartm.*) Loch Ascog, Bute. River Tay, Perth. Coltfoot, Elgin. P. VII. VIII. S. I.

\*\* *L. all submersed, membranous, lanceolate or broader ; stipules free.*

10. *P. Zizii* (Roth.); *l.* pellucid narrow-lanceolate-oblong cuspidate narrowed downwards minutely denticulate near the end wavy, *stip.* long broad broadly and doubly keeled not winged, "fr. like that of *P. lucens*," *ped.* swelling upwards, very long.—*J. of B.* xvii. t. 204.—*St.* long much branched, lower *l.* usually recurved. *Ped.* 4—5 in. long, thicker than the stem. Spike about 1 in. long.—Cauldshiels Loch, Melrose. *P.* VII. VIII. S.

11. *P. lucens* (L.); *l.* pellucid oval or lanceolate stalked minutely denticulate and cuspidate, *stip.* winged, fr. blunt on the back and slightly keeled when fresh, *peduncle swelling upwards*, spikes cylindrical densely flowered.—*E. B.* 376. *R.* vii. 36.—All the stipules have two prominent wings on their back. Spikes long, about equalling the peduncles.—A form (?) has been found with ovate-lanceolate subacute subcoriaceous floating leaves (var. *coriaceous* Nolte?) which looks different. Sometimes (*P. acuminatus* Schum.) the midrib forms a long prominent point.—Common in deepish water. *P.* VI. E. S. I.

12. ?*P. decipiens* (Nolte); *l.* pellucid oblong rounded below not hooded but blunt mucronate and wavy at the end entire sessile or uppermost *l.* sometimes subsessile, *stip.* not winged<sup>1</sup>, fr. "like that of *P. lucens*," *ped.* equal.—*J. of B.* v. t. 61. *R.* vii. 35.—*St.* long, much branched. *Ped.* 1—2 in. long, thicker than the stem. Spikes 1—2 in. long, dense.—Canal near Bath. Navan Canal. *P.* VII. E. S. I.

13. *P. longifolius* (Bab. not Gay); *l.* pellucid long-lanceolate narrowed below entire cuspidate, *stip.* winged, fr. . . . ., *ped.* long swelling upwards, spikes with few rather whorled distant flowers.—*E. B. S.* 2847.—*St.* long, slender. *L.* distant, not denticulate, nearly sessile, with very short stalks. *Stip.* green, lanceolate, with 2 narrow wings on the back. *Ped.* 6—8 in. long, much thicker than the stem. Spike 1 in. long; fl. rather more whorled than as represented in *E. B. S.* Fr. unknown.—Deep water. Found floating loose in Lough Corrib, Galway. *P.* VIII. I.

14. *P. prælongus* (Wulf.); *l.* pellucid oblong-lanceolate half-clasping blunt and hooded at the end entire, *stip.* not winged, fr. keeled on the back when fresh keeled or winged when dry, *peduncle* very long equal, spikes many-flowered.—*E. B. S.* 2858. *R.* vii. 33.—*St.* long, growing in deep water. *L.* with parallel

<sup>1</sup> I cannot find the wings described by Syme; nor did Mr. Baker. *J. of B.* v. 71.

veins adjoining the midrib. Peduncles 6—12 in. long. Spikes 1—2 in. long.—Rivers and ditches, rare. P. V. VI. E. S. I.

15. *P. perfoliatus* (L.); *st.* round *l.* pellucid cordate-ovate clasping not hooded, *fr.* rounded on the back when fresh keeled when dry, peduncle equal.—*E. B.* 168. *R.* vii. 29.—*St.* long, slightly branched, rather dichotomous at the top. Peduncles rather thick and short. Spikes short.—Lakes and streams. P. VII. E. S. I.

16. *P. crispus* (L.); *st.* compressed, *l.* pellucid linear-oblong blunt sessile serrate wavy, *fr.* with a long beak keeled on the back when dry, peduncle equal.—*E. B.* 1012. *R.* vii. 29—30.—*St.* much branched. *Ped.* long. Spikes few-flowered. Beak as long as the nut. *L.* usually crisped at the edges, occasionally flat (*P. serratus* Huds.).—Ditches and streams. P. VI. E. S. I.

\*\*\* *L.* all submersed, alternute, linear-ligulate; stipules free.

17. *P. zosterifolius* (Schum.); *st.* flattened, *l.* linear-acuminate with 3 principal and many close parallel intermediate veins occupying the whole surface, spikes cylindrical upon long ped., sep. transversely oval, *fr.* obovate keeled, style terminal.—*E. B. S.* 2685. *R.* vii. 27. *P. cuspidatus* Sm., *P. compressus* Fries, Hook.—*Ped.* 2—4 in. long. Spikes 10—15-flowered. Inner edge of *fr.* rounded; faces a little convex. *L.* suddenly acuminate or apiculate.—Rivers and lakes. P. VI. E. S. I.

18. *P. acutifolius* (Link); *st.* flattened, *l.* linear acuminate with 3 principal and many close parallel intermediate veins occupying the whole surface, spikes ovate about as long as the short ped., sep. rhomboidal, *fr.* broadly  $\frac{1}{2}$ -obovate compressed inner edge straight with a tooth near its base keeled, style facial.—*E. B. S.* 2609. *R.* vii. t. 26.—*Ped.* very short. Spikes 4—6-flowered. *L.* gradually acuminate. Styles hooked, continuing the inner edge of the fruit.—Marsh ditches, rare. P. VI. E.

19. *P. obtusifolius* (Koch); *st.* slightly compressed with rounded edges, *l.* linear 3-veined, spikes ovate dense continuous about as long as the short ped., sep. rhomboidal, *fr.* obovate keeled.—*R.* vii. t. 25. *P. gramineus* Sm., *E. B.* 2253.—*St.* slender, much branched. Peduncles very short. *L.* rounded off to a slight point at the end, with oblong network near the midrib, wanting the fine parallel veins of the two preceding species.—Ponds and ditches, rare. P. VI. VII. E. S. I.

20. *P. mucronatus* (Schrad.); *st.* slightly compressed, *l.* linear 5-veined suddenly apiculate, spikes short lax  $\frac{1}{2}$  or  $\frac{2}{3}$  shorter than the compressed clavate ped., sep. transversely oval, *fr.* obliquely

ovate bluntly keeled.—*P. compressus* Sm. *E. B.* 418. *R.* vii. 24, not *Fries*, nor *Koch*.—Lateral veins nearer together and nearer to the margin of the l. than to the midrib (rarely some l. are 3-veined), no intermediate veins.—Like *Sp.* 17 rather than *Sp.* 19.—Ditches. P. VI. VII. E. S. I.

21. *P. pusillus* (L.); st. subterete, l. linear 3-veined, spikes short rather lax  $\frac{1}{2}$  or  $\frac{2}{3}$  shorter than the slender ped., sep. roundish-reniform, fr. obliquely ovate bluntly keeled.—*E. B.* 215. *R.* vii. 22.—St. slender. L. narrow, rather acute, without intermediate parallel veins, the lateral veins equidistant between the midrib and margin. Ped. scarcely compressed, not thickened.—[*P. rutilus* (Wolfg., *Fries*) has 3-veined attenuate-cuspidate l., semiterete st., and oblong-elliptic fr. without a keel. *Fl. Dan.* 2891.]—Ponds and ditches. P. VI. E. S. I.

22. *P. trichoides* (Cham.); st. subterete, l. subsetaceous 1-veined finely pointed, spikes short lax long-stalked, fr. semicircular compressed obscurely keeled, straight inner edge with a tooth near its base.—*Sy. E. B.* 1420. *R.* vii. 21.—St. very slender, a little thickened below the joinings; branches fasciculate. L. not transversely veined. Floral stip. large. Ped. not thickened. Sep. roundish, stalked. Monandrous. Fr. often warted on the back and with a tubercle on each side at its base when fresh.—Framlingham, Norf. Wrotham, Suff. P. VII. VIII. E.

\*\*\*\* L. all submersed, alternate, linear, sheathing.

23. *P. flabellatus* (Bab.); lower l. broadly linear abruptly apiculate or acuminate 3—5-veined with transverse veins, upper l. narrow acute 3-veined, fr. (2 lin. long) broadly  $\frac{1}{2}$ -obovate inner edge nearly straight but gibbous near the top rounded on the back, nut with a prominent keel.—*Sy. E. B.* 1421. *Phytol.* iv. 1158.—Rhizome spreading, from a tuber that has outlived the winter. Floating st. branched, spreading like a fan. Broad lower l. usually decayed at the time of flowering. Lateral veins of upper l. at the margin, of lower l. distant from it. Spikes slightly interrupted. Back of fr. without ridges, rounded when fresh; enclosed nut with faint lateral ridges.—Ponds and ditches chiefly near the sea. B. VI. VII. E. I.

24. *P. pectinatus* (L.); l. formed of 2 interrupted tubes, lower narrowly linear flattened slightly grooved above, upper setaceous, fr. (2 lin. long) broadly  $\frac{1}{2}$ -obovate inner edge rather convex rounded on the back, nut with 2 lateral ridges but (usually) no keel.—*E. B.* 323. *R.* vii. 19.—St. branched, forming linear masses. L. very gradually acute, all 1-veined, no marginal veins and scarcely thickened there; upper l. with an oval section.



Spikes slightly interrupted. Back of fr. without rigidities and rounded when fresh; nut with strong lateral ridges. Varying greatly in length and size of st. and leaves.—*P. marinus* (Huds.) has the stems naked below. *Sy. E. B.* 1423.—Ponds and streams. P. VI. VII. E. S. I.

25. *P. filifor'mis* (Pers.); *l. linear-setaceous* 1-veined with transverse veins, *spikes greatly interrupted, fr.* ( $1\frac{1}{2}$  lin.) obovate rugose rounded on the back without keel or ridges when dry, nut round-backed.—*Sy. E. B.* 1424.—L. like those of the preceding but longer. Fr. smaller. Whorls very distant on the spikes. *Peduncles very long.*—Rare. Scotland. Lough Cullen and Lough Conn, Co. Mayo, Ireland. P. VI. VII. S. I.

\*\*\*\*\* *L. all opposite, submersed; stipules none.*

26. *P. den'sus* (L.); *l. all opposite* pellucid clasping elliptic-lanceolate or lanceolate, spikes shortly stalked ultimately reflexed.—*E. B.* 397. *R. vii. t.* 28.—L. crowded, rather recurved. Spike 4-flowered. Sep. triangular.—Ditches. P. VI. VII. E. S. I.

## 2. RUP'PIA Linn. Tassel-Pondweed.

1. *R. marit'ima* (L.); *ped. long spirally twisted*, anth.-cells oblong  $1\frac{1}{2}$  times as long as broad, nut ovoid rather obliquely erect.—*E. B.* 136. *R. vii.* 17. *R. spiralis* Hartm.—Whole plant stronger than the next. L. very narrowly linear, formed as in *Potamogeton pectinatus*. *Sheaths large, inflated.*—Salt marshes in the South. P. VII. VIII. E. I.

2. *R. rostellata* (Koch); *ped. short not spiral*, anth.-cells squarish as long as broad, nut gibbous at base obliquely ascending when young.—*Sy. E. B.* 1428. *R. vii.* 17.—Whole plant very slender. L. rather filiform than linear. *Sheaths small, close.*—Salt marshes. P. VI.—VIII. E. S. I.

## 3. ZANNICHEL'LIA Linn. Horned Pondweed.

1. *Z. palus'tris* (L.); style at least half as long as the divergent nuts.—*E. B.* 1844.—Floating. L. slender, opposite, filiform. Fl. axillary, sessile.— $\alpha$ . *Z. brachistemon* (Gay); *fr.* 2—4 *sessile* with crenate back, style  $\frac{1}{2}$  length of fr., stigma large crenulate, anth. 2-celled.— $\beta$ . *Z. macrostemon* (Gay); *fr.* 2—4 *sessile* rarely crenate, style  $\frac{1}{2}$  length of fr., stigma small crenulate, *anth. 4-celled.*— $\gamma$ . *Z. pedicellata* (Fr.); *fr.* 2—5 *stalked* muricate, style as long as fr., stigma large crenulate, anth.

2-celled. *Sy. E. B.* 1426.—I doubt the constancy of these characters.—Stagnant water.  $\gamma$ . Brackish water. A. or P. V.—VIII. E. S. I.

2. *Z. polycar'pa* (Nolte); style scarcely  $\frac{1}{2}$  of length of converging sessile nuts, *fr.* 5—6 subsessile crenulate on back, *converging*, style scarcely  $\frac{1}{2}$  length of *fr.*, stigma large repand, anth. 2-celled.—Much like *Sp.* 1.—In water. A. V.—VIII. Orkney. Belfast. S. I.

### Order XCVI. NAIADACEÆ.

Fl. monœcious or dicecious, in a spath. Perianth 0. Stam. free, 1—3. Ovary 1, superior, with 1 ovule. Style 1; stigmas 2—4, filiform. *Fr.* a nut enclosed in the persistent spath. 1-celled, 1-seeded, not bursting. Albumen 0. Embryo with a thin skin having a lateral cleft.

1. ZOSTERA. Fl. imperfect. Stam. and pistils inserted in two rows upon one side of a spadix. Spath linear, leaflike. Fl. naked. Anth. 1. Ovary 1; style 1; stigmas 2.
2. NAIAS. Fl. imperfect, solitary, sheathed; no perianth. Barren fl. of 1 stamen. Fertile: style short; stigmas 2—4 filiform.

#### 1. ZOSTE'RA Linn. Grass-wrack.

1. *Z. nána* (Roth); l. linear 1-veined, ped. filiform, spadix short with 2—5 clasping bands, nuts smooth.—*E. B. S.* 2931. *R.* vii. 2.—Plant small, scarcely 3 in. long. L. slender. Ped. pale-coloured throughout,  $\frac{1}{2}$  as broad and quite as long as the inflated oblong-lanceolate spaths. Ripe seed shining black.—Muddy estuaries. P. ? VII. VIII. E. S. I.

2. *Z. marina* (L.); l. linear 1—7-veined, spadix without bands, nuts striate.—*E. B.* 467. *R.* vii. 4.—Plant long. L. broad. Ped. rather strong. Ripe seed milky white.—*Z. angustifolia*, *Sy. E. B.* 1430, *R.* vii. 3, is a slender form growing upon mud, with fewer veins in its l. and rather longer peduncles which are much compressed and green upwards and narrow gradually into the spadix.—In the sea. P. ? VII. VIII. E. S. I.

#### 2. NA'IAS Linn.

1. *N. flex'ilis* (Rostk.); l. very narrow and very minutely denticulate, sheaths ciliate-denticulate.—*Sy. E. B.* 1432.—L.

ternate or opposite, 1-veined, pellucid, remotely denticulate. Ovary solitary, axillary, sessile, oblong. Style short.—Deep lakes. A. VIII. S. I.

### Division III. GLUMIFERÆ.

(Orders XCVII. and XCVIII.)

Leaves parallel-veined, persistent. Floral envelopes imbricate, bractlike.

#### Order XCVII. CYPERACEÆ.

Fl. perfect or unisexual, each from the axil of a scale (or glume) imbricate on a common axis. Perianth 0, or rarely membranous. Stam. hypogynous (3 or rarely 2 in our plants), with sometimes a row of bristles or abortive filaments. Anth. erect, entire at the apex. Ovary 1-celled, 1 ovule at its base, often surrounded by bristles or enclosed in a bottle-shaped perianth. Style simple, trifid or bifid. Embryo enclosed within the base of the albumen.—L. with entire sheaths.

Tribe I. *CYPEREÆ*. Fl. perfect. Gl. 2-ranked. Perianth 0.

1. *CYPERUS*. Spikelets 2-ranked, many-flowered. Gl. of 1 valve, many, keeled, nearly all with flowers.
2. *SCHÆNUS*. Spikelets 2-ranked, 2—4-flowered. Gl. 6—9; several lower ones smaller, empty. Bristles few or 0.

Tr. II. *SCIRPEÆ*. Fl. perfect. Gl. imbricate on all sides. Perianth 0.

\* *Lowest glumes empty or smaller than the others.*

3. *CLADIUM*. Spikelets 1—3-flowered. Gl. 5 or 6. Bristles 0. *Nut with a thick fleshy coat, tipped with the conical base of the style.*
4. *RHYNCHOSPORA*. Spikelets few-flowered. Gl. 6 or 7. *Bristles about 6. Nut compressed, convex on both sides, crowned with the dilated base of the style.*

\*\* *Lowest glume empty or larger than the others, sometimes all fertile.*

† Bristles scarcely equalling the glume or wanting.

5. *ELEOCHARIS*. One or 2 lowest gl. broader, empty. Bris-  
s 2

gles 3—6. Nut compressed, crowned with the persistent dilated base of the style.

6. SCIRPUS. Glumes nearly equal; or 1 or 2 lowest broader, empty. Bristles about 6 or 0. Nut plane-convex or trigonous; *base of the style filiform, not dilated.*
7. BLYSMUS. Two lowest gl. broader, and empty. Bristles 3—6. Nut plane-convex, tipped with the not dilated base of the style. *Spikelets with bracts, alternate, forming a close distichous compound terminal spike.*

†† Bristles ultimately much exceeding the glumes.

8. ERIOPHORUM. Glumes nearly equal, lowest sometimes empty. Bristles ultimately silky. Nut trigonous.

Tr. III. ELYNEÆ. Fl. unisexual. Perianth 0 or formed of 1 or 2 scales.

9. KOBRESIA. Spikes close together. Lower fl. fem., perianth of 1 scale enclosing the germen and covered by the glume. Upper fl. male, without any perianth.

Tr. IV. CARICEÆ. Fl. unisexual. Nut completely enclosed in the bottle-shaped perianth.

10. CAREX. Fl. in imbricate spikes, each covered by a glume. Female fl. with a single bottle-shaped persistent perianth, 1 style and 2—3 stigmas. Male fl. of 3 stam., without a perianth.

### Tribe I. *Cypereæ.*

#### 1. CYPE'RUS Linn. Galingale.

1. *C. lon'gus* (L.); spikelets linear-lanceolate in twice-compound umbels, peduncles of partial umbels erect unequal, stigmas 3, creeping.—*E. B.* 1309. *St.* 52. 10.—St. triangular, 2—3 feet high. Umbel very large, lax, unequal, its base with 2 or 3 long leaves. Glumes brownish red, with green keels and pale margins.—South of England, rare. P. VIII. IX. E.

2. *C. fus'cus* (L.); spikelets linear-lanceolate in small roundish heads at the extremities of the branches, gl. spreading, stigmas 3, root fibrous.—*E. B. S.* 2626. *St.* 52. 5.—A small nearly prostrate plant. Stems many, 2—5 in. long. Heads with 3

unequal l. at the base. Glumes fuscous with green keels.—Formerly at Little Chelsea, Middlesex; Shalford Common, Surrey. Jersey. A. VIII. IX. E.

## 2. SCHÆ'NUS Linn. Bog-rush.

1. *S. nig'ricans* (L.); st. terete naked, spikelets 5—10 collected into a terminal roundish head overtopped by the lower bract, gl. rough at the keel.—*E. B.* 1121. *St.* 40. 9.—Root of strong black fibres. *St.* 8—12 in. high, clothed at the base with blackish-brown smooth shining scales some of which terminate in setaceous erect leaves which are shorter than the stem. Bristles variable in number, short, rough, with upward spines. Stigmas 3. Anth. ending in a point. Gl. dark brown or black.—Turfy bogs. P. VI. E. S. I.

## Tribe II. *Scirpeæ*.

### 3. CLA'DIUM Pat. Br. Sedge.

1. *C. Maris'cus* (R. Br.); panicles lateral and terminal repeatedly compound, spikelets capitate, st. subterete leafy smooth, l. finely serrate on the margins and keel.—*E. B.* 950.—Creeping. *St.* 3—4 feet high. L. very long, rigid, narrowed and triquetrous towards the end. Fl. in each spikelet 1—3; but usually only one nut is produced.—Bogs and fens, rare. P. VII. *Common Sedge*. E. S. I.

### 4. RHYNCHOS'PORA Vahl.

1. *R. al'ba* (Vahl); spikelets in a compact corymb about as long as the outer bracts, stam. 2, bristles with declining teeth, base of the style without teeth.—*E. B.* 985. *St.* 40. 7.—Slightly creeping. *St.* 6—12 in. high. L. narrowly linear. Spikelets whitish. Bristles 9—12. Filaments slender.— $\beta$ . *sordida*; spikelets brownish, in small oval clusters, often overtopped by the outer bracts.—Turfy bogs. P. VII. E. S. I.

2. *R. fus'ca* (Sm.); spikelets in an oval head considerably shorter than the outer bracts, stam. 3, bristles with ascending teeth, base of the style with erect teeth.—*E. B.* 1575. *St.* 40. 6.—Creeping extensively. *St.* 6—8 in. high. L. nearly filiform. Spikelets brown. Bristles 6. Filaments dilated.—Bogs rare. South-west of England. Ireland. P. VII. VIII. E. I.

## 5. ELEOCHARIS R. Br.

Spikes terminal and solitary in all our species.

1. *E. palustris* (R. Br.); gl. rather acute, *lowest gl.*  $\frac{1}{2}$ -surrounding the spike, stigmas 2, nut roundish plane-convex with rounded edges smooth crowned with the ovate base of the style and falling short of the 4 bristles, base of *st.* clothed with membranous almost transversely truncate sheaths.—*E. B.* 131. *St.* 9. *R.* viii. 297.—Often only slightly creeping, 6—12 in. high. Sheaths with a very blunt point on one side.—Wet and marshy places. P. VI. E. S. I.

2. *E. uniglumis* (Link); gl. rather acute, *lowest gl.* almost surrounding the spike, stigmas 2, nut pearshaped blunt rather compressed with rounded edges very finely punctate-striate crowned with the conical base of the style and falling short of the bristles, base of *st.* clothed with transversely truncate sheaths.—*Sy. E. B.* 1587. *R.* viii. 296.—Far-creeping, 6—8 in. high.—Wet sandy places especially near the sea. P. VI. VII. E. S.

[*E. Watsoni* (Bab.); nut oblong very blunt a little narrowed below exceeding the bristles.—*A. N. H.* ser. 2. x. 19.—It is probably a form of *Sp.* 2.—Taynloan, Argyleshire. *Dr. Balfour.* Murrough of Wicklow. *Dr. D. Moore.*] E. S.

3. *E. multicaulis* (Sm.); glumes blunt, stigmas 3, nut topshaped triquetrous smooth crowned with the broad triquetrous base of the style equalling the six bristles, base of *st.* clothed with obliquely truncate rather acute sheaths.—*E. B.* 1187. *St.* 78. 11. *R.* viii. 296.—Slightly creeping, about 6 in. high.—Marshy places. P. VII. E. S. I.

4. *E. acicularis* (Sm.); glumes blunt, stigmas 3, nut obovate-oblong compressed longitudinally ribbed and transversely striate crowned with the topshaped base of the style, bristles short deciduous.—*E. B.* 749. *St.* 10. *R.* viii. 294.—Root fibrous with slender runners. *St.* many, slender, erect, 3—4 in. high, when growing in water sometimes a foot long. Spikes very small.—In damp places upon heaths. P. A. (Koch) VII. VIII. E. S. I.

## 6. SCIRPUS Linn. Club-rush.

\* *Bristles* 6. *Spikes* many. SCIRPUS.

1. *S. maritimus* (L.); spikes stalked or sessile in a dense terminal cluster, bracts several leaflike, gl. bifid with acute lobes and a point between them, nut obovate trigonous smooth.—

*E. B.* 542. *St.* 13. 3. *R.* viii. 310, 311.—Creeping, sometimes tuberous. *St.* 1—3 feet high, leafy. Spikes large, sometimes solitary. Stigmas 3, or rarely 2.—Salt marshes. *P.* VII.

E. S. I.

2. *S. sylvaticus* (L.); spikes clustered in a large cymose very compound terminal panicle, clusters stalked and sessile, general bracts several leaflike, gl. blunt apiculate, nut obovate bluntly trigonous.—*E. B.* 919. *St.* 36. 8. *R.* viii. 313.—*St.* 2—3 feet high. Spikes very many, small, greenish, ovate. L. broad, flat. Stigmas 3.—[*S. radicans* (Schk.), spikes all stalked, glumes not apiculate, stoles long and rooting, may possibly be found.]—Damp woods and banks. *P.* VII.

E. S. I.

3. *S. triquetér* (L.); *st. acutely triquetrous throughout*, spikelets in a small cymose panicle, gl. notched mucronate *glabrous* fringed with rounded blunt lobes, nut "roundish-obovate plane-convex smooth."—*E. B.* 1694. *St.* 36. 3. *R.* viii. 305.—*St.* 3—4 feet high, with concave faces; 1 or 2 long sheaths at the base, the uppermost ending in a *short broad triquetrous leaf*. Lower bract long and rigid, resembling a prolongation of the stem. Spikelets small, stalked and sessile. Anth. with a short beardless point. Stigmas 2.—Muddy banks of the Thames near London; the Arun, Sussex; and the Tamer near Calstock. Cornwall. *P.* VIII.

E.

[*S. pun'gens* (Vahl); *st. acutely triquetrous throughout*, spikelets few *sessile*, gl. notched mucronate smooth with *acute lobes*, nut roundish-obovate plane-convex smooth.—*E. B.* 2819. *R.* viii. 304. *S. Rothii* Hoppe? *St.* 36. 4. *St.* 6—18 in. high, slender; with several sheaths at the base ending in *long narrow keeled leaves*. Lower bract very long and rigid, resembling a prolongation of the stem. Spikelets large, ovate, blunt, all sessile. Anth. with a subulate fringed point. Stigmas 2.—*St.* Ouen's Pond, Jersey. *P.* VI. VII.]

4. *S. carinátus* (Sm.); *st. terete* below bluntly trigonous upwards, spikes in a small cymose panicle, gl. notched mucronate *slightly asperous* and pilose fringed, nut "convex on the back smooth."—*E. B.* 1983. *S. Duvallii* Hoppe, *St.* 36. 2. *R.* viii. 308.—*St.* 2—4 feet high; with 1 or 2 long sheaths at the base, the uppermost ending in a leaf 3 or 4 in. long. Lower bract much overtopping the panicle. Stigmas 2.—By rivers, near London and in Sussex and Kent. *P.* VI. VII.

E.

5. *S. lacus'tris* (L.); *st. terete*, spikes in a terminal twice-compound panicle, gl. notched mucronate *glabrous* fringed, nut *bluntly trigonous obovate*, stigmas 3.—*E. B.* 666. *St.* 36. 1. *R.* viii. 306.—Far-creeping. *St.* 4—6 feet high, naked; with 1 or

2 long sheaths at the base. *Anth. bearded at the end.* Panicle terminal. The bract sometimes resembles a continuation of the stem. *Sometimes it has long nearly flat floating leaves.*—Rivers and ponds. P. VI. VII. *Bulrush.* E. S. I.

6. ? *S. Tabernæmontani* (Gm.); st. terete, spikes in a terminal compound panicle, *gl.* notched mucronate *asperous* fringed, *nut* compressed *roundish-oblong* smooth, stigmas 2.—*R.* viii. 307. *S. glaucus* Sm., *E. B.* 2321.—Creeping. St. 2 feet high, with 1 or 2 long sheaths at the base. *Anth. not bearded.* Panicle smaller than in the preceding. Lower bract short. Fr. convex on one side.—Rivers and ponds. P. VI. VII. E. S. I.

\*\* *Bristles* 4—6. *Spike solitary, terminal.* BÆOTHRYON Dietr.

7. *S. cæspitosus* (L.); st. terete striate with *many imbricate leafless acute scales* and sheaths with *short subulate l.* below, spike ovate few-flowered, *gl.* ovate membranous pointed, 2 *outer gl.* as long as the spike enclosing it and ending in long rigid leaflike points, *nut* obovate oblong mucronate smooth, bristles longer than the nut with a few erect teeth near the tip.—*E. B.* 1029. *R.* viii. 300.—St. 3—6 in long, many, erect, many of them barren. *Bristles* 6.—Barren turfy heaths. P. VI.—VIII. E. S. I.

8. *S. pauciflorus* (Lightf.); st. terete striate with a few thin narrow leafless scales and one tight abrupt leafless sheath below, spike ovate few-flowered, *gl.* ovate keeled membranous at their edges, 2 *outer gl.* blunt shorter than the spike and enclosing it, *nut* obovate mucronate finely netted, bristles shorter than the nut with declining teeth.—*E. B.* 1122. *St.* 10. *R.* viii. 299.—St. 3—10 in. long, erect, many of them barren, stoloniferous. *Bristles* 6.—Boggy moors and heaths. P. VI.—VIII. E. S. I.

9. *S. parvulus* (R. & S.); st. round many each with one close-pressed leafless sheath, no true leaves, spike oval few-flowered, *gl.* ovate blunt keeled membranous, 2 outer ones rather longer, *nut* ovate-oblong mucronate smooth, bristles twice as long as the nut with declining teeth throughout.—*J. of B.* vi. 85. *Sy. E. B.* 1591. *R.* viii. 299.—Root fibrous with capillary stoles ending in ovate-subulate tubers. St. about an inch high, several, with 2 or 3 longitudinal fibres and more or less perfect transverse lines. *Bristles* 4—6.—Lymington, Hants (extinct). Poole, Dorset. Mouth of river Avoca, Co. Wicklow. Aveton Gifford, Dev. A. VII. E. I.

\*\*\* *Bristles* 0.—† *Spike solitary.* ELEOGITON Link.

10. *S. fluitans* (L.); st. floating branched leafy, l. clustered, fl.-stalks alternate with a sheathing l. at the base, spike terminal



ovate few-flowered, gl. blunt keeled membranous at their edges, 2 outer gl. larger shorter than the spike and enclosing it, *nut obovate*, stigmas 2.—*E. B.* 216. *R.* viii. 298.—St. rooting from the lower joinings and spreading to a great extent in a zigzag manner. On mud it is cæspitose with truly sheathing leaves.—Ditches and ponds. P. VI. VII. E. S. I.

†† St. round, leafy at the base. Spikes 1—3. *ISOLEPIS* R. Br.

11. *S. setaceus* (L.); spikes terminal, lower bract long so as to resemble a short continuation of the st., gl. blunt mucronate, *nut* trigonous obovate *longitudinally ribbed* and transversely striate, stigmas 3.—*E. B.* 1693. *R.* viii. 301.—St. tufted, slender, 3—6 in. high. Spikes small, sessile, considerably shorter than the lower bract. Gl. brown with whitish margins and a green keel.—Wet sandy and gravelly places. P. VII. E. S. I.

12. *S. Sav'ii* (S. & M.); lower bract shorter or slightly longer than the terminal spike, gl. blunt submucronate, *nut* subglobose *rough with minute points*, stigmas 3.—*E. B.* S. 2782. *R.* viii. 301.—Closely resembling *S. setaceus*. Spikes varying considerably in length, sometimes 1 in. long. Gl. scarcely mucronate, greenish, usually with a brown spot on the upper part of each side.—In many places near the coast. A. or P. VII. E. S. I.

††† Spikes many, clustered. *HOLOSCHÆNUS* Link.

13. *S. Holoschæ'nus* (L.); st. round, spikes in dense globular sessile or stalked clusters, lower bract erect long, gl. obovate emarginate mucronate.—*E. B.* 1612.—St. 3—4 feet high, round quite up to the cluster. Upper bract patent or ascending; lower very long, with a flat open white channel. Gl. variegated with fuscous and white, pilose. Anth. with a long entire or toothed point.—Braunton Burrows, Devon. P. IX. E.

## 7. BLYS'MUS Panz.

1. *B. compres'sus* (Panz.); spikelets 6—8-flowered, outer gl. ribbed shorter than the spikelet, bristles 3—6 strong persistent with declining teeth, l. flat rough on the edges and keel.—*Scirpus* Sm., *E. B.* 791. *R.* viii. 293.—St. 6—8 in. high. Outer gl. of lowest spikelet with a subulate leaflike point which often overtops the spike. Glumes reddish brown. Nut lenticular, crowned with the long persistent style, shining.—Boggy pastures. P. VI. VII. E. S.

2. *B. rufus* (Link); spikelets 2—4-flowered, outer gl. smooth as long as the spikelet, bristles 1—6 slender deciduous with patent or ascending teeth, l. channelled not keeled smooth.—*Scirpus* Sm., *E. B.* 1010. *St.* 85. 7. *R.* viii. 293.—St. slender.

Gl. dark brown, polished. Nut ovate, with a long beak, opaque.—Marshes near the sea on the Northern and Western coasts. P. VII. E. S. I.

8. ERIOPHORUM Linn. Cotton-grass.

\* *Bristles* 4—6, at length *crisped*. *Spike solitary*.

[1. *E. alpinum* (L.); st. triquetrous rough, l. very short, spike oblong.—*E. B.* 311 (excl. the leafy shoot). *R.* viii. 288. *St.* 10.—A slender elegant plant.—Moss of Restenet, Forfar (but now lost through drainage). P. VI.] S.

\*\* *Bristles very many, not crisped*.—† *Spike solitary*.

2. *E. vaginatum* (L.); cæspitose, st. *trigonus* above round below, spike oblong, nut *obovoid*, l. long setaceous triquetrous channelled, upper sheath inflated leafless.—*E. B.* 873. *R.* viii. 289.—Bogs and moors. P. V. *Hare's-tail Cotton-grass*. E. S. I.

†† *Spikes more than one*.

3. *E. polystachion* (L.); st. nearly terete, *peduncles smooth*, l. linear channelled their upper half triangular, nut elliptic-acuminate or obovoid triquetrous.—About a foot high and rather slender. L. triangular through more than half their length. Bristles 3 or 4 times as long as the spikes. *E. B.* 564. *R.* viii. 291.—*β. minus*; st. and l. very slender. *E. gracile* Sm., *E. B.* 2402 (not Koch). An alpine form has but one nearly sessile spike. *E. capitatum* Don? *E. B.* 2387.—*γ. elatius* (Koch); st. strong tall, l. 2—3 lines broad the triangular part commencing above the middle. *E. polystachion* Sm.—Bogs. *β* on mountains. P. V. VI. *Common Cotton-grass*. E. S. I.

4. *E. latifolium* (Hoppe); st. triquetrous in its upper half, *peduncles asperous*, l. linear, nearly flat contracted above the middle into a triangular point, nut *pyriform* triquetrous.—*R.* viii. 292. *E. pubescens* Sm., *E. B.* S. 2633. *E. polystachion* *E. B.* 563.—A slender plant, 12—18 in. high. L. about 2 lines broad; triquetrous point short. Several of the elegant spikes upon longish stalks which are asperous. Bristles 2 or 3 times as long as the spikes.—Bogs, rather rare. P. V. VI. E. S. I.

5. *E. gracile* (Koch); st. subtriquetrous, *peduncles downy*, l. *narrowly linear triquetrous*, nut *oblong-linear* triquetrous.—*E. B.* S. 2886. *R.* vii. 290. *E. triquetrum* Hoppe, *St.* 10. 2.—A tall slender plant. Spikes about 4, most of them on downy not asperous stalks. Gl. with many ribs. Bristles about twice

as long as the spike.—Bogs. Halnaby, Yorkshire. Whitemoor Pond, Guildford. P. VI. VII. E.

### Tribe III. *Elyneæ*.

#### 9. KOBRESIA Willd.

1. *K. caricina* (Willd.).—*E. B.* 1410. *Schk.* Rrr. 161. *R.* viii. 193.—St. erect, 6—12 in. high. L. slender, falling short of the stem. Spikes 4—5, collected at the top of the stem, 6—8-flowered. There is an abortive stam. (?) at the base of the nut; but some authors, considering each fl. a separate spike, think that this represents a second flower.—Moors. Yorkshire. Durham. Perthshire. P. VII. E. S.

### Tribe IV. *Cariceæ*.

#### 10. CA'REX Linn<sup>1</sup>.

i. *Monostachyæ*. Spike simple, solitary, terminal. (Sp. 1—5.)

\* *Dicæcious* or *monæcious* with male fl. at the top. *Stigmas* 2.

1. *C. dioica* (L.); usually dicæcious, fertile spike ovate dense, *fr. ascending* ovate many-veined angles rough near the top, nut roundish oval, terete st. and subsetaceous l. roundish, soboliferous.—*E. B.* 543. *Schk.* A. 1. *H.* a. 1. *R.* 194.—About 6 in. high.—Spongy bogs. P. V. VI. E. S. I.

[2. *C. Davalliána* (Sm.); dicæcious, *fr. deflexed* ovate-lanceolate angles rough near the top, nut "linear-oblong," st. and margins of the subsetaceous l. rough, cæspitose.—*E. B.* 2123. *R.* 194.—About 6 in. high.—The true plant did grow on Lansdown near Bath, but is now lost by drainage. P. VI.] E.

3. *C. pulicáris* (L.); half of spike male, *fr.* remote at length *deflexed* oblong-lanceolate compressed, nut linear-oblong plane-convex, l. involute.—*E. B.* 1051. *Schk.* A. 3. *H.* a. 3. *R.* 195.—St. slender, 6—12 in. or more in height, erect, smooth. L.

<sup>1</sup> In the following descriptions, *fruit* means the persistent bottle-shaped perianth, including the ripe nut or true capsule. The *glume* is always taken from the fertile spike unless it is otherwise stated.—*Schk.* refers to the plates of *Schkuhr's Riedgräser*; *H.* to *Hoppe's Caricol. Germ.* in *Sturm Deutschl. Flora*; and *R.* to *Reichenbach Ic. Fl. Germ. Cent.* vii.; *B.* to *Boott's Illust. of Carex*. See also *Andersson's Pl. Scand.*, and *Lang in Linnæa*, xxiv. 481; and for form, &c., of the nuts *Des Moulins Cat. Dordogne, Suppl. Final*.

slender, erect. Glumes deciduous. Fr. dark brown.—Bogs.  
P. VI. E. S. I.

\*\* *Spike male at the top. Stigmas 2.*

4. *C. rupestris* (All.); half of spike male, *fr. obovate* trigonous with a very short beak adpressed scarcely longer than the persistent gl., nut obovoid acutely triquetrous, l. flat.—*E. B. S.* 2814. *H. b.* 4. *R.* 198.—Soboliferous. St. 3—6 in. high, triquetrous, rough upwards. *L. ending in a wavy rough slender triangular point.* Gl. fuscous. Fr. paler.—Lofty mountains. P. VII. S.

5. *C. pauciflora* (Lightf.); 1—3 *terminal fl. male, fr. 2—4 lanceolate-subulate terete* patent or reflexed longer than the deciduous gl., nut linear-oblong trigonous, l. involute.—*E. B.* 2041. *Schk. A.* 4. *H. b.* 1. *R.* 196.—Soboliferous. St. usually about 5 inches high, slender. *L. 2 or 3, much shorter than the stem.* Fr. pale yellow.—Bogs in the North. Pl. VI. VII. E. S.

ii. *Homostachya*. Spikelets in a compound continuous or interrupted spike, male at one or both ends or nearly unisexual. Nut plane-convex or compressed. Stigmas 2. (Sp. 6—24.)

† Spikelets nearly unisexual. Sobole far-creeping.

6. *C. disticha* (Huds.); *spikelets in an oblong interrupted spike upper and lower ones fertile intermediate mostly male, fr. ovate-lanceolate veined narrowly margined bifid with serrate edges above, nut oval, gl. shorter than the fr. acute its midrib not reaching the top, st. with rough angles.*—*C. intermedia* Good., *E. B.* 2042. *Schk. B.* 7. *H. a.* 14. *R.* 210.—Height 1—2 feet. Fr. about as long as gl.; usually abortive and twice as long. Lower bract with a slender leaflike point.—Marshy places. P. V. VI. E. S. I.

7. *C. arenaria* (L.); *spikelets in an oblong interrupted spike upper ones male lower fertile intermediate male at the end, fr. ovate veined winged and serrulate from the middle to the bifid top of the beak, nut bluntly ovoid, glumes longer than the fr. acuminate, st. rough above.*—*E. B.* 921. *Schk. B. & Dd.* 6. *H. a.* 13. *R.* 209.—Height 1 foot. Sobole superficial, very long. Lowermost bracts with slender leaflike points.—Sandy places. P. VI. E. S. I.

†† Spikelets male at the top.—‡ *Sobole far-creeping.*

8. *C. incurva* (Lightf.); spikelets in a roundish head, fr. inflated broadly ovoid acuminate-rostrate, beak split externally,

nut obovoid, st. smooth about equalling the leaves.—*E. B.* 927. *Schk.* Hh. 95. *H. a.* 5. *R.* 199.—St. 2—3 in. high, usually *recurved* so as to bring the large head down to the ground. Beak of fr. usually rather rough.—Sandy shores of the North. P. VI. E. S.

9. *C. divisa* (Huds.); spikelets in a somewhat ovate head, *fr. plane-convex* ovoid many-veined, *beak acutely bifid with finely serrate edges*, nut broadly oblong, gl. with an excurrent rib, *st. roughish at the top*.—*E. B.* 1096.—St. slender, a foot high. Lowermost bract (brown) scarious, often prolonged into a slender green point sometimes overtopping the spike. Fr. veined on both sides. Spike often interrupted below.—Near the sea on the Southern and Eastern coasts. P. V. VI. E. I.

†† *Cæspitose. Spike simply compound. Fr. squarrose, not gibbous.*

10. *C. vulpina* (L.); spikelets compound in a cylindrical oblong crowded spike, *fr. ovoid-acuminate plane-convex* veined, *beak bifid finely serrate, nut ovoid or oval with a beak constricted at the base*, gl. mucronate shorter than the fruit, *st. triquetrous with rough angles, bracts setaceous*.—*E. B.* 307. *Schk.* C. 10. *R.* 217.—Height 2 feet. St. firm. L. broad. Fr. palish green. Spikelets greenish, bracts long, in shady places. *R.* 216.—Wet places. P. VI. E. S. I.

11. *C. muricata* (L.); spikelets contiguous, spike oblong dense or interrupted below, *fr. ovoid-acuminate plane-convex obscurely veined bifid finely serrate* upper spreading, *nut ovoid* its beak extremely short, gl. mucronate shorter than the fruit, *st. triquetrous with rough angles*.—*E. B.* 1097. *R.* 215.—Height 1—2 feet. St. slender but strong. L. narrow. Lowermost spikelets not more than their own length distant from each other. Fr. much larger than that of Sp. 12, with a broad flat beak with very sharp edges.—Gravelly pastures. P. VI. E. S. I.

12. *C. divulsa* (Good.); spikelets distant the upper ones nearer together, *fr. ovoid acute plane-convex obscurely veined bifid* smooth ascending, *beak roughish at the edges, nut ovoid-oblong* compressed its beak extremely short, gl. mucronate shorter than the fruit, *st. triquetrous with rough angles above, bracts setaceous*.—*E. B.* 629. *Schk.* Dd. & Ww. 89. *H. a.* 16. *R.* 220.—Height 1—2 feet. *St. lax, slender, flaccid.* Spikelets greyish, usually distant, 1 or 2 lowest often lengthened into a short branch. Fr. with a thick green margin slightly rough near the top.—Moist shady places. P. VI. E. I.

††† *Cæspitose*. *Spike compound, often panicled. Fr. ascending, gibbous on its back.*

13. *C. teretius'cula* (Good.); *spikelets forming a dense compound oblong spike, fr. ovate with 2—5 central ribs on the convex side (back), beak bidentate serrulate split to its base and overlapping on the back, nut turbinate with a very short beak convex, style not thickened at the base, st. trigonous and rough above.—E. B. 1065. Schk. D. 19. T. 69. H. a. 9. R. 222.—Root forming scattered simple tufts. St. 1—2 feet high, slender.—β. C. Ehrhartiana* (Hoppe); *root more cæspitose, spike long rather loose, st. triquetrous above. Sy. E. B. 620.—Boggy meadows, rare, β near Manchester. P. VI. E. S. I.*

14. *C. paradox'a* (Willd.); *spikes narrowly panicled lower branches rather distinct, fr. ovoid with many short elevated ribs near its base, beak obliquely bidentate serrulate not split nor winged on its back, nut rhomboidal constricted below doubly convex with a short beak, style slightly enlarged at the base, st. trigonous and rough above.—E. B. S. 2896. Schk. E. 21. H. a. 12. R. 222.—Root densely tufted, crowned with the fibrous remains of decayed leaves. St. 1—2 feet high, slender.—Bogs. Near Mullingar, Ireland. Near York. Hoveton, Norfolk. P. VI. VII. E. I.*

15. *C. paniculata* (L.); *spikes panicled with long diverging branches, fr. ovoid faintly many-veined with a bifid fringed beak split to its base and overlapping on the back, nut ovoid blunt narrowed below plane-convex, beak slightly thickened upwards, st. triangular.—E. B. 1064. Schk. D. 20, Ttt. 163. H. a. 19. R. 223.—Root forming dense elevated tufts. St. stout, 2—3 feet high. Panicle usually large and spreading or reduced to a slender compound (or even simple) spike. Bracts all much shorter than the spike.—Bogs. P. VI. E. S. I.*

††† Spikelets male at their base (or at both ends).

‡ *Sobole far-creeping.*

[*C. brizoides* (L.) was probably an escape.] E.

†† *Cæspitose. Bracts leaflike. Lower spikelets distant, simple or compound.*

16. *C. Boeninghauseniána* (Weihe); *spikelets several upper ones simple crowded lower distant alternately branched, fr. lanceolate plane-convex tapering serrulate from below the middle, beak deeply split on one side, nut ovoid elliptic, gl. equalling the fruit, root tufted, lower bract at least equalling the spike.—*

*E. B. S.* 2910. *H. a.* 34. *Kunze Riedg.* 22. *R.* 219.—*St.* 1—2 feet high, triangular with slightly convex faces and rough edges. Inflorescence often a foot long. Rachis straight, with 3 rough edges. *Gl.* ovate, membranous, silvery brown, smooth. Lower spikelets wholly male, male at both ends or at either end. Bracts, except the lowest, short. *L.* channelled.—Marshes and pond-sides, rare. *P. VI.* *E. S.*

17. *C. axillaris* (Good.); spikelets several upper ones simple crowded lower distant densely compound, fr. ovoid-lanceolate plane-convex tapering deeply bifid serrulate above the middle, nut obovoid with a beak, *gl.* shorter than the fruit, root tufted, lower bract as long as or longer than the spike.—*E. B.* 993. *H. a.* 33.—*St.* 1—2 feet high, acutely triangular. Rachis straight, with 3 rough angles. *Gl.* ovate, membranous, brownish; *midrib* often rough, extending to the point. Divisions of the lower spikelets crowded into the axils of the bracts. Bracts, except the lowest, short. *L.* flat. Spikelets male at the base or at both ends.—Marshes, rare. *P. VI.* *E. S. I.*

18. *C. remota* (L.); spikelets several all simple upper ones crowded lower distant, fr. ovoid-acuminate plane-convex notched at the end serrulate above, nut ovoid with a beak, *gl.* shorter than the fr., root tufted, bracts long.—*E. B.* 832. *Schk.* *E.* 23. *H. a.* 35. *R.* 212.—*St.* 1—2 feet high, trigonous. Rachis with 2 rough angles in its upper part. *Gl.* oblong, membranous, greenish white; *midrib* smooth, usually not reaching the point. Several of the bracts long. *L.* channelled. Sometimes with a thick rooting densely proliferous rhizome. *C. tenella* (Sm. not Schkr.) is a starved form.—Damp places. *P. VI.* *E. S. I.*

‡‡‡ *Cæspitose.* Bracts not leaflike. Spikes contiguous or slightly distant.

‡ Spikes and leaves glaucous or dusky.

19. *C. stellulata* (Good.); spikelets about 4 globose rather distant, fr. divergent broadly ovoid acuminate plane-convex striate, beak bifid with serrate edges, nut ovoid abruptly narrowed below, *gl.* shorter than the fruit.—*E. B.* 806. *Schk.* *C.* 14. *H. a.* 28. *R.* 214.—*St.* 6—12 in. high, triquetrous, nearly smooth. Lowermost spikelet often with a short bract. *Gl.* ovate, membranous, reddish with a green keel and white edges. Ripe fr. greenish, stellate.—Boggy places. *P. V. VI.* *E. S. I.*

20. *C. elongata* (L.); spikelets many oblong contiguous, fr. patent oblong-acuminate plane-convex with many ribs on both sides, beak almost entire with rough edges, nut linear-oblong tapering below beak very short style persistent, *gl.* shorter than

the fruit, *bracts none* or one very short.—*E. B.* 1920. *Schk. E.* 25. *H. a.* 32. *R.* 218.—St. 1—2 feet high, triquetrous. Gl. ovate, dark brown with a green keel and whitish edges, blunt, sometimes apiculate. Upper spikelets crowded, lower lax but not distant.—Marshes, rare. P. VI. E. I.

21. *C. canescens* (L.); spikelets 4—6 elliptic contiguous, *fr. erect acute plane-convex faintly striate, beak short notched not split* rough at the edges, *nut elliptic* beak very short style persistent, glumes ovate shorter than the fruit.—*E. B.* 1386. *Schk. C.* 13. *R.* 206. *C. curta*, Good.—St. a foot high, triquetrous, smooth except at the top. Lowermost spikelet often with a setaceous bract. *Glumes membranous, whitish* with a green keel, blunt, apiculate. Fr. whitish.—Bogs. P. VI. E. S. I.

22. *C. vit'ilis* (Fr.); spikelets 4—8 ovate or oblong contiguous, *fr. erect ovate plane-convex faintly striate, beak short split to its base externally* rough at the edges, "nut elliptic," glumes ovate shorter than the fruit.—*C. curta*  $\beta$ . *alpicola* Wahl. *C. Persoonii* Sieb., *R.* 206.—Creeping. St. 6—12 in. high, triquetrous, smooth except at the top. Lowermost spikelet often with a setaceous bract. *Gl. membranous, brown* with a white margin.—Mountains. P. VII. VIII. E. S.

#### ‡ Spikes fuscous.

23. *C. lagopina* (Wahl.); spikelets 3—4 roundish-elliptic contiguous terminal one longer, *fr. erect elliptic acuminate plane-convex narrowed below nearly entire at the point with smooth edges, nut elliptic tipped with the persistent style, gl. ovate nearly as long as the fruit.*—*H. a.* 24. *R.* 205. *C. leporina* Linn. *Fl. Lap.* not *Fl. Suec.*—*E. B. S.* 2815.—St. 4—8 in. high, smooth, triangular. Glumes reddish with the edges paler. Fr. yellow.—Loch na Gar, and Cairn Toul, Aberdeenshire. P. VIII. S.

24. *C. ovális* (Good.); spikelets about 6 oval contiguous, *fr. erect ovate-attenuate plane-convex narrowed below bifid at the point with membranous edges serrulate* above, nut elliptic with a short cylindrical beak ending in a persistent style, gl. lanceolate as long as the fruit.—*E. B.* 306. *Schk. B.* 8. *C. leporina* L. *Fl. Suec. H. a.* 22. *R.* 211.—St. 1—2 feet high, triangular, smooth, or roughish above. Gl. acute, brown with a paler membranous margin. Fr. yellowish.—Meadows. P. VII. E. S. I.

iii. *Heterostachyæ*. One or more terminal spikes wholly or rarely only partially male, others axillary fertile. (Sp. 25—71.)

\* *Beak of fr. short, entire or emarginate or shortly 2-toothed.*

† Nut plane-convex. Stigmas 2—3. Male spikes 1 or more.



† *Stigmas* 2. *Sheaths of leaves webbed. Fertile spikes erect. L. ultimately revolute-edged; lowest sheaths usually leafless.*

25. *C. stricta* (Good.); fertile spikes erect long cylindrical, bracts auricled lowermost leaflike short, *fr. oblong-elliptic acute, compressed veined* closely imbricate, nut roundly obovoid shortly beaked, *sheaths* of l. with *filamentous network*.—*E. B.* 914. *R.* 230. *C. cæspitosa* Gay.—*Densely cæspitose.* St. 2—3 ft. high, triquetrous, rigid. L. short, narrow. Fertile spikes often male at the top. Gl. blunt or acute, equalling or shorter than the fr., narrow, dark purple; keel green. Fr. in 6—9 regular rows.—Marshes. P. VI. E. S. I.

‡‡ *Stigmas* 2. *Sheath not webbed, usually all bearing leaves. L. ultimately revolute-edged. Stoloniferous.*

26. *C. acuta* (L.); male *spikes* 1—3, fertile 2—4 slender cylindrical-acuminate erect with fruit, lowermost bract leaflike often overtopping the stem *with long auricles, fr. oblong lenticular veined*, nut roundish-obovoid with a short slender beak, gl. acute.—*E. B.* 580. *Schk. Ee. & Ff.* 92. *H. a.* 44. *R.* 231, 232.—St. 2—3 feet high, triquetrous, rough at top. L. broad. Gl. narrow-lanceolate, acute, on the male spikes spathulate-lanceolate, purple with a green keel. Fr. pale, blunt or acute, round or oval, spherical or flattened. Fertile spikes usually with a few male fl. at the end, *nodding with flowers*.—Syme mentions a cæspitose form or distinct species (*C. proluxa* Fr.?).—Wet places. P. VI. E. S. I.

27. *C. rigida* (Good.); spikes erect, male 1, fertile oval or shortly cylindrical dense sessile, lower bract leaflike, *fr. elliptic lenticular without veins*, nut roundish blunt with a slender beak, st. triquetrous rough towards the top, l. curving outwards *with deflexed edges*.—*E. B.* 2047. *H. a.* 40. *R.* 225. *C. saxatilis* L. ?—St. 6 in. to a foot or more high. L. broad, flat, keeled, rigid. Bracts without sheaths. Spikes near together, short, rarely narrowed and laxly flowered below. Gl. about exceeding the fr., purple with a green keel. Nut rather longer than broad.—In wet and stony places on mountains. P. VI. VII. E. S. I.

‡‡‡ *Stigmas* 3. *Sheaths not webbed. Fertile spikes erect. Edges of leaves ultimately involute. Glumes narrower than the fruit.*

28. *C. aquatilis* (Wahl.); spikes erect, male 1 or more, fertile 3 or 4 long narrowed below lower ones stalked, bracts leaflike erect overtopping the stem, *fr. elliptic lenticular without*

veins broader than the gl., nut oblong narrowed below with a short slender beak, *st. trigonous smooth*.—*E. B. S.* 2758.—*St.* 1—4 ft. high, with convex faces. Spikes rather distant. Gl. usually short, narrower than the yellowish-green fr., reddish purple with a pale midrib.—Alpine boggy tablelands of the Clova Mountains and in the valley by the bridge at Clova. P. VII. S.

29. *C. vulgaris* (Fries); spikes erect, male 1 or 2, fertile 3—4 subsessile cylindrical, bracts with short auricles leaflike, fr. elliptic plane-convex with many veins below and a short entire beak, nut roundish very blunt, st. triquetrous rough towards the top.—*C. cæspitosa* Sm., *E. B.* 1507. *C. Goodenovii* Gay.—*St.* about a foot high. *L.* slender, not keeled. Bracts without sheaths. Spikes near together, short. Gl. blunt, shorter than fr., purple; keel slender, pale green. Fr. greenish or with a purple tinge. Nut rather broader than long.—Marshes. P. V. VI. E. S. I.

[*C. Gibsoni* (Bab.); fr. lanceolate acute, nut broadly obovoid.—*A. N. H.* xi. t. 5.—*St.* 6—8 in. high. Perianth nearly twice as long as the nut, gradually narrowed from below the middle to the top. Perhaps a montrosity of *C. vulgaris*.—Hebden Bridge, Yorkshire! Not recently found. P. VI.] E.

†† Nut with 3 angles. Stigmas 3, rarely 2.

‡ Fruit glabrous. Terminal spike male at its base.

30. *C. alpina* (Sw.); spikes 1—4 roundish or oblong contiguous nearly sessile, fr. obovoid veinless rough above with a short notched beak, gl. acutish falling short of fr., nut obovoid triquetrous blunt with a short cylindrical beak, bract scarcely overtopping the spikes, st. triangular rough towards the top.—*B. t.* 356. *C. Vahlia* *E. B. S.* 2666. *Schk. Gg.* 94 & Ppp. 154. *R.* 235.—*St.* 6—12 in. high, erect. Gl. brown or black.—Glen Callater and Glen Fiagh, Clova. P. VII. S.

31. *C. Buxbaumii* (Wahl.); spikes 3—4 oblong sessile contiguous, the lowest shortly stalked rather distant, fr. oval blunt compressed (ultimately trigonous) bidentate roughish above veined, gl. cuspidate lower exceeding fr., nut obovoid trigonous blunt apiculate, lower bract leaflike.—*E. B. S.* 2885. *H. b.* 11. *Schk. X.* & *Gg.* 76. *R.* 235. *B.* 438.—*St.* 1—2 feet high, triquetrous, rough. Sheaths of the l. connected by netlike filaments. Gl. nearly black, with a green keel prolonged into a cuspidate point. Fr. glaucous green.—Harbour Island near Toom Bridge, Lough Neagh. P. VI. I.

32. *C. atráta* (L.); spikes 3—4 ovate-oblong shortly stalked contiguous ultimately drooping, the lowest rather distant and with a longer stalk, *fr. elliptic* veinless with a short *terete slightly notched beak*, glumes acute, *nut elliptic* triquetrous blunt apiculate, lower bract leaflike. *E. B.* 2044. *Schk.* X. 77. *H. b.* 8. *R.* 237.—St. 1— $1\frac{1}{2}$  foot high, triangular, smooth. L. flat, broad. Gl. dark purple; midrib slender, pale. Fr. yellowish. Mr. H. C. Watson mentions a form with the term. spike wholly male and the fem. spikes cylindrical and 2—3 in. below it.—Alpine rocks. P. VI. VII. E. S.

‡‡ *Fruit glabrous. Terminal spike wholly male, solitary.*

33. *C. palles'cens* (L.); fertile spikes subpendulous ovate or oblong with exserted stalks *contiguous*, bracts leaflike, gl. mucronate, *fr. ovoid-oblong* convex on both sides veined *blunt, beak 0, nut linear-elliptic* trigonous.—*E. B.* 2185. *Sy. E. B.* 1657. *Schk.* Kk. 99. *H. b.* 44.—St. slender, triquetrous, rough above 1— $1\frac{1}{2}$  foot high. Spikes blunt, *pale green*; the barren one sessile, darker.—Marshy places. P. VI. E. S. I.

34. *C. paníceu* (L.); fertile spikes erect remote subcylindrical on exserted stalks with distant fl., bracts leaflike sheathing, gl. rather acute, *fr. ovoid subglobose inflated* veinless with a short *terete truncate beak*, nut obovoid-oblong trigonous with a cylindrical beak.—*E. B.* 1505. *Schk.* Ll. 100. *H. b.* 33.—St. 1—2 ft. high, erect, smooth. Fertile spikes about 2. Gl. oblong, more or less acute, dark brown with a green keel and membranous pale margins. Lowermost bract about as long as its spike, the rest shorter, sheaths close.—Marshy places. P. VI. E. S. I.

35. *C. vaginata* (Tausch!); fertile spikes erect remote on exserted stalks with distant fl., bracts sheathing scarcely leaflike, gl. bluntish, *fr. ovoid triquetrous glabrous* veinless with a short *terete smooth very obliquely truncate and emarginate beak, nut elliptic* triangular “with a beak slightly thickened upwards.”—*H. b.* 17. *Kunze Riedg.* 15. *B.* 478. *C. phæostachya* Sm., *E. B. S.* 2731. *C. Meilichoferi* Sm., *E. B.* 2293.—St. 5—6 in. high, smooth. Fertile spikes 1—2. Glumes bluntish. Bracts with funnelshaped sheaths.—Highland mountains. P. VII. S.

36. *C. limósa* (L.); fertile spikes 1 or 2 upon very long stalks drooping *ovate densely-flowered* “with occasionally a few male fl. at their top,” bracts auricled slender strongly keeled, gl. ovate mucronate, *fr. roundish-obovoid compressed strongly ribbed* with a very short entire beak, nut obovoid bluntly trigonous with a beak, *l. narrow linear channelled rough at the edges* throughout.—*E. B.* 2043. *R.* 238. *B.* 216.—Creeping. St. 1 ft. high. L. and bracts very slender. Gl. purple, with a green keel, about

as long as the pale fr. which is broadest above the middle. Nut pale.—Spongy bogs. P. VI. E. S. I.

37. *C. irrigua* (Hoppe); *fertile spikes 2 or 3 drooping* upon long stalks oblong densely-flowered *with occasionally a few male fl. at their base, bracts auricled leaflike rather broad nearly flat*, gl. ovoid-lanceolate attenuate acute, *fr. roundish-ovoid compressed faintly ribbed* with a very short entire beak, nut elliptic triangular with a beak, *l. linear flat smooth at the edges* except near the tip.—*E. B. S. 2895. R. 238. B. 219.*—Creeping. St. a foot or more in height. L. and bracts 2 or 3 times as broad as in the preceding. *Gl. wholly purple* usually longer than the pale fr. which is broadest below the middle. Nut pale.—Spongy bogs. Muckle Moss, Northumb. Terregles, Dumfries. Ben Mime near Loch Lomond. P. VI. E. S.

38. *C. rariflora* (Sm.); *fertile spikes 2 or 3 drooping* upon long stalks *oblong few-flowered lax, bracts with very short sheaths*, gl. very broad blunt as long as the fr., fr. oblong with 3 blunt angles, *beak extremely short* entire, nut roundish-oblong, *l. flat rough-edged towards the tip.*—*E. B. 2516. B. 217.*—Creeping. St. 6—8 in. high, smooth. *Gl. folded round the fr., dark brown; midrib pale, terminating in a minute apiculus.* Fr. pale, faintly veined; nut darker.—Elevated bogs in highlands. P. VI. S.

39. *C. capillaris* (L.); *fertile spikes drooping* upon long half-included stalks *few-flowered lax, one bract sheathing several stalks*, gl. blunt, fr. oblong trigonous turgid narrowed below veinless with a slender membranous beak, nut obovoid triquetrous blunt with a short beak.—*E. B. 2069. Schk. O. 56. H. b. 53. R. 241.*—St. very slender, 2—6 in. high, smooth. Peduncles rough, several usually enclosed in one sheath. Gl. short, broad, blunt, midrib not reaching the tip, falling short of the small smooth brown fruit. Nut pale. Root tufted.—Teesdale. Scottish Highlands. P. VI. E. S.

40. *C. strigosa* (Huds.); *fertile spikes drooping* about 4 distant rather long *slender lax* lower with exserted stalks, bracts leaflike sheathing, *fr. oblong-lanceolate* narrowed at both ends trigonous veined *with an obliquely truncate mouth*, nut elliptic triangular punctured, *l. broad.*—*E. B. 994. Schk. N. 53.*—St. 2 feet high. Sheaths nearly covering the peduncles. Gl. elliptic-lanceolate, diaphanous, greenish down the back.—Groves and thickets, rare. P. V. VI. E. S. I.

41. *C. pen'dula* (Huds.); *fertile spikes drooping* about 5 distant *cylindrical very long densely flowered*, bracts leaflike lower ones with sheaths nearly equalling the flowerstalks upper scarcely

sheathing, fr. elliptic subtrigonus *tumid* with short trigonus *emarginate* beak, nut elliptic triangular.—*E. B.* 2315. *Schk.* Q. 60. *R.* 243.—St. 3—6 feet high, round at the angles above. Fertile spikes often 3 or 4 in. long, arched; upper ones frequently with male fl. at the tip. Gl. ovate mucronate, brown with a green keel. Fr. green, ciliate at the mouth.—Damp woods. P. V. E. S. I.

††† Fruit downy, hairy or scabrous. Bracts sheathing. L. in sterile tufts. Male spike solitary.

42. *C. hūmilis* (Leyss.); fertile spikes 2 or 3 remote about 3-flowered enclosed in the membranous leafless bracts, fr. obovoid subtrigonus narrowed below with an entire oblique mouth not obovoid, trigonus, with a short beak.—*H. b.* 15. *C. clandestina* Good., *E. B.* 2124. *Schk.* K. 43. *R.* 239.—St. about 2 in. high, erect, concealed amongst the leaves. Bracts large, wholly membranous, nearly hiding the fertile spikes. L. all radical, linear, channelled, rough.—Limestone hills in Wilts and Somerset and Dorset. P. IV. E.

43. *C. ornithop'oda* (W.); fertile spikes 3 near together suberect exceeding male spike, bracts membranous leafless, fr. pyriform trigonus with a subemarginate mouth exceeding the glumes, nut elliptic subtrigonus shortly stalked with a minute beak.—*J. of B.* xiii. t. 164. *R.* 240.—St. 3—5 in. high, scarcely exceeding the flat bluntish radical leaves. Bracts much falling short of fertile spikes.—Millers Dale, Derbyshire. P. V.—VII. E.

44. *C. digitata* (L.); fertile spikes 2 or 3 distant linear erect lax, bracts membranous obliquely truncate, lowermost with a setaceous leaflike point, fr. obovoid trigonus narrowed below equalling gl., beak short nearly entire, "nut elliptic-oblong triangular shortly-stalked and shortly-beaked."—*E. B.* 615. *Schk.* H. 38. *H. b.* 14.—St. 6—8 in. high, erect, sheathed at the base, leafless, taller than the flat radical leaves.—Woods on limestone, rare. P. IV. V. E.

†††† Fr. hairy or downy or scabrous. Bracts not sheathing (except slightly in No. 49).—† Male spike 1.

45. *C. præcox* (Jacq.); fertile spikes 1—3 oblong-ovoid near together sessile, bracts clasping the lowest leaflike and slightly sheathing, gl. broadly ovate acuminate with excurrent midrib, fr. ovoid-rhomboidal trigonus with an entire mouth, nut obovoid narrowed below trigonus with prominent angles, base of the style surrounded by a cuplike disk, *soboliferous*.—*E. B.* 1099. *Schk.* F. 27. *H. b.* 24. *R.* 261.—St. 3—12 in. high. Lowermost spike often slightly stalked.—Dry places. P. IV. V. E. S. I.

46. *C. ericetorum* (Poll.); *fertile spikes* 1—3 *ovoid* near together sessile, bracts clasping all membranous, *gl. obovate very blunt finely ciliate* midrib not reaching the top, *fr. obovoid trigonous* with a truncate beak, nut subglobose trigonous with no terminal disk or prominent angles, *soboliferous*.—*E. B. S.* 2971. *Schk. J.* 42. *H. b.* 26. *R.* 262. *C. ciliata* Willd.—*Gl. brown* with a broad pale edge, all very blunt. *St.* 3—6 in. high.—On chalk. Norf., Suff., Cambridge. P. IV. V. E.

47. *C. montana* (L.); *fertile spikes* 1—3 *ovoid* crowded sessile, bracts small membranous the lowest with an awlshaped point, *gl. of fertile spikes blunt notched and mucronate, fr. narrowed below oblong-obovoid* trigonous with a short *notched beak*, nut oblong narrowed below with a pyramidal beak, *root fibrous* from a shaggy thick branched rhizome.—*E. B. S.* 2924. *Schk. F.* 29. *H. b.* 21. *R.* 261. *C. collina* Willd.—*St.* about a span long, slender. L. narrow; sheaths of lower l. red. *Gl. very dark*; midrib narrowly yellowish. *Fr. hairy*, its beak purple.—Heaths and banks. P. IV. V. E.

48. *C. pilulifera* (L.); *fertile spikes* about 3 *roundish* near together sessile, bracts small lowest scarcely leaflike awlshaped not sheathing, *gl. broadly ovate mucronate, fr. stalked subglobose* with a short *bifid beak*, nut subglobose subtrigonous narrowed below, persistent base of style recurved, *root fibrous*.—*E. B.* 885. *Schk. J.* 39. *R.* 260.—*St.* 6—12 in. long, slender, at length decumbent.— $\beta$ . *Leesii* (Ridl.); lowest bract very long leaflike, upper bract long and slender, midrib of *gl. excurrent*, nut fusiform. *J. of B.* xix. t. 218.—Wet heaths.  $\beta$ . Plumpton, Yorkshire. Glen Callater, Braemar. On rocks in shade. P. V. E. S. I.

49. *C. tomentosa* (L.); *fertile spikes* 1 or 2 nearly sessile *cylindrical blunt, lowermost bracts leaflike* with a very short sheath, *gl. broadly ovate acute, fr. obovoid* subtrigonous scarcely beaked *slightly emarginate*, nut blunt trigonous narrowed below with a *short beak constricted at its base, soboliferous*.—*E. B.* 2046. *Schk. F.* 28. *H. b.* 28. *R.* 263.—*St.* a foot high, with 3 sharp angles, rough upwards, *erect*. Beak of nut slightly swelling upwards. *Fr. with copious white down*, mouth very broad.—Water-meadows at Merston Measey, Wilts. P. VI. E.

‡‡ Male spikes usually more than 1.

50. *C. glauca* (Scop.); *fertile spikes* 2 or 3 *erect or drooping cylindrical densely flowered long-stalked*, bracts leaflike scarcely sheathing, *gl. ovate acute, fr. blunt elliptic veinless* slightly

rough entire at the small point, nut roundish-ovate triangular.—*H. b.* 67. *Schk.* O. P. 57. *R.* 269. *C. recurva* Huds., *E. B.* 1506.—Soboliferous. St. a foot or more in height. Male spikes variable in number. Fertile spikes often have a male fl. at the top.— $\beta$ . *C. Micheliiana* (Sm.); gl. blunt, fr. smaller. *E. B.* 2236.— $\gamma$ . *C. stictocarpa* (Sm.); fertile spikes ovate, fr. obovate dotted. *E. B. S.* 2772.—Damp places. P. VI. E. S. I.

\*\* *Beak of fr. long, 2-toothed or bifid. Nut with 3 angles. Stigmas 3, rarely 2. (Sp. 51—71.)*

† *Beak of fruit terete. Terminal spike male.*

51. *C. ustulata* (Wahl.); fertile spikes 2 or 4 upon short stalks ovoid densely flowered, bracts scarcely leaflike or sheathing, gl. ovate acute, fr. elliptic compressed rough-edged with a cloven beak, nut elliptic triangular on a long stalk, root fibrous.—*E. B.* 2404. *Schk.* Y. 82. *H. b.* 47.—St. 3—10 in. high. L. very short, broad. Gl. dark purple with a slender pale midrib. Fr. dark purple, paler below. Nut fuscous.—Ben Lawers. *Mr. G. Don.* See *Sy. E. B.* x. 137. P. VII. S.

†† *Beak of fruit plane-convex. Male spike 1, or rarely 2. Fruit glabrous. Bracts sheathing, leaflike.*

52. *C. frigida* (All.); fertile spikes fusiform upper sessile lower ultimately pendulous long-stalked, bracts sheathing leaflike, gl. acute, fr. lanceolate triquetrous gradually narrowed into a ciliate-serrate bifid beak, nut ovate triquetrous, st. triangular smooth.—*Edm. Bot. Tr.* xii. t. 2. *J. of B.* xiii. t. 159.—St. 3—12 in. high. L. broad, parallel-sided when young.—Above Loch Ceann-mor, Aberdeenshire. P. VIII. S.

53. *C. flava* (L.); fertile spikes roundish-oval subsessile, lowest spike with a nearly included stalk, bracts leaflike with short sheaths, gl. blunt, fr. ovate inflated ribbed smooth narrowed into a deflexed rough-edged bifid beak, nut obovoid trigonous punctate, st. trigonous smooth.—*E. B.* 1294. *Schk.* H. 36. *H.* p. 22. *R.* 273.—St. 6—12 in. high. L. broad. Male spike cylindrical, blunt; gl. blunt. Fertile spikes usually near together and near the barren spike, sometimes distant; gl. with a green midrib slightly rough and often excurrent at the end. Beak of the fr. curved downwards parallel-sided when young. Lowest fr. declining.— $\beta$ . *C. lepidocarpa* (Tausch!); barren spikes usually long-stalked, beak of fr. often nearly straight.—*Sy. E. B.* 1673. *Kunze Riedgr.* 13. *R.* 272.—Wet places. P. V. VI. E. S. I.

54. *C. Œderi* (Ehrh.); fertile spikes roundish-ovate subsessile near together, lowest spike with an included stalk, bracts leaflike

with short sheaths, gl. blunt (?), fr. *subglobose* inflated ribbed smooth *suddenly contracted into* an erect narrow rough-edged bifid beak, nut obovoid-trigonus punctate, st. trigonus smooth.—*Sy. E. B.* 1674. *H. b.* 23. *R.* 272.—Much like, but distinct from, *C. flava*. Fertile spikes smaller. Fr. much smaller than those of *C. flava*, more in number, with a shorter and narrower-based beak which is conical when young. Very variable in height.—Bogs. P. VI. VII. E. S. I.

55. *C. exten'sa* (Good.); fertile spikes oblong near together subsessile lower one rather distant with a short included stalk, bracts very long leaflike with short sheaths, gl. *mucronate*, fr. ovoid triquetrous ribbed narrowed into a *straight smooth-edged bifid beak*, nut *oblong-elliptic* triangular smooth.—*E. B.* 833. *Schk. V. Xx.* 72. *H. b.* 32. *R.* 274.—*St.* usually curved, 8—12 in. high, trigonus, *smooth*, usually exceeding the leaves. Barren spike nearly sessile, blunt; gl. blunt. *L. and bracts very narrow*, convolute, long.—Marshes, chiefly near the sea. P. VI. E. S. I.

56. *C. Hornschuchiana* (Hoppe); fertile spikes ovate-oblong distant with stalks exceeding the long sheaths, *glumes ovate* not mucronate, fr. ovoid triquetrous rough-edged also *scarious in the notch*, nut obovoid trigonus nearly smooth.—*H. b.* 40. *R.* 252. *C. speirostachya Sm., E. B. S.* 2770.—Rootstock often creeping. *St.* trigonus, rough-edged, about a foot high. Lowest bract often reaching up to or beyond the barren spike.—Boggy places. P. VI. E. S. I.

[*C. xanthocarpa* (Degl.) "has tufted stems, ovate-acute glumes, spreading fr. not scarious in the notch." It is said to be not uncommon. I do not know it.—*C. fulva* Good., Sm., was an error.—See *J. of B.* xiv. 366.]

57. *C. punctata* (Gaud.); fertile spikes erect cylindrical with slightly exserted stalks particularly the lowest, bracts sheathing, gl. ovate shortly awned, fr. spreading ovoid *tumid obscurely veined pellucidly punctate* with a *linear bidentate smooth beak*, nut *ovoid-rhomboidal* narrowed at both ends triangular *rough*.—*H. b.* 37. *Kunze Riedg.* 6. *R.* 251.—*St.* smooth, 1—2 feet high, slender. Spikes distant or the upper ones near; stalks slightly (the lowest often greatly) exserted, rough; lowest spike often very distant. Gl. pale red with a broad green longitudinal dorsal band. Fr. pale. Nut brown. Gl. of barren spike blunt.—Marshy places near the sea. P. VI. E. S. I.

58. *C. distans* (L.); fertile spikes remote erect oblong, upper with included stalks, bracts with sheaths, gl. mucronate, fr. ascending ovoid trigonus *equally and faintly ribbed* pellucidly punctate smooth, edges of the bifid narrow beak rough, nut triquetrous roughish *obovoid narrowed below*.—*E. B.* 1234.



*Schk.* T. 68. *H. b.* 42. *R.* 253.—St. smooth, seldom exceeding a foot high, slender. Spikes distant, short; lower ped. half-exserted. Gl. brownish. Fr. yellowish brown, rather inflated; nut yellowish. Male spike cylindrical-clavate, with blunt glumes.—Marshy places, especially near the sea. P. V. E. S. I.

59. *C. bimer'vis* (Sm.); fertile spikes remote, upper ones nearer together cylindrical their stalks mostly included, lower long with exserted stalks, bracts sheathing, gl. mucronate, fr. ovoid subtri-quetrous with 2 prominent green submarginal ribs on the back other ribs faint, beak broad bifid rough at the edges, nut obovoid roughish.—*E. B.* 1235. *Schk.* Rrr. 160. *H. b.* 39. *R.* 255.—St. triangular, smooth, a foot high. Spikes often very distant; upper stalks often quite included, never much exserted, lower often greatly exserted. Gl. dark purple; midrib greenish yellow. Fr. brown or deeply tinged with purple, 2 prominent ribs always green; nut brown. Male spike with blunt glumes.—Heaths and moors. P. VI. VII. E. S. I.

60. *C. lævigata* (Sm.); fertile spikes remote cylindrical, stalks more or less exserted, bracts sheathing, gl. acute, fr. ovoid attenuate striate, beak long deeply bifid with rough edges, nut subpyriform narrowed below triangular smooth.—*E. B.* 1387. *Schk.* Bbb. 116 & Sss. 162. *H. b.* 38. *R.* 254.—St. smooth, 2—3 feet high. Spikes distant, erect or drooping. Gl. often acute on the male spike, always so on the others, purple with a paler dorsal longitudinal band. Rarely 2 male spikes. Fr. green; nut yellowish. L. broad.—Marshes and wet thickets, rather rare. P. VI. E. S. I.

61. *C. depauperata* (Good.); fertile spikes erect remote 3- or 4-ft., stalks exserted, bracts sheathing leaflike, gl. acute, fr. large nearly globose, beak long bifid with rough edges, nut elliptic trigonous with bluntish angles.—*E. B.* 1098. *Schk.* M. 50.—St. 1—2 feet high, trigonous, smooth. Gl. of the male spike blunt. Fr. very large and few with many ribs. Spikes very distant.—Dry woods, very rare. P. VI. E.

††† Beak of fr. glabrous, terete or compressed. Male spike 1 (rarely more). Stigmas 3.

62. *C. sylvatica* (Huds.); fertile spikes about 4 distant slightly drooping linear with long half-exserted stalks, bracts leaflike sheathing, fr. elliptic trigonous obscurely veined narrowed into a long cloven smooth beak, nut obovoid-elliptic-triangular, l. narrower than in the preceding.—*E. B.* 995. *Schk.* Ll. 101. *H. b.* 55. *R.* 242.—St. about 2 feet high, smooth, its top and the fr.-beak rarely a little rough. Sheaths scarcely half equalling the

peduncles. Gl. ovate, acute, diaphanous with a green keel.—  
Damp woods. P. V. E. S. I.

63. *C. Pseudo-cyperus* (L.); fertile spikes about 5 drooping cylindrical densely flowered stalked near together, bracts leaflike scarcely sheathing, gl. setaceous rough dilated at the base, fr. ovoid-lanceolate ribbed narrowed into a deeply bifid beak, nut elliptic, st. triquetrous with rough angles.—*E. B.* 242. *Schk.* Mm. 102. *H. b.* 56. *R.* 275.—St. 2—3 feet high. Male spike often with some fertile flowers. Fertile spike  $1\frac{1}{2}$ —2 in. long.—Damp places, rare. P. VI. E. I.

†††† Beak of hairy fr. terete or compressed with patent cusps.  
Male spikes 2 or more.

64. *C. filiformis* (L.); male spikes 2, fertile 3 or 4 remote erect sessile oblong, bracts leaflike lowermost slightly sheathing, gl. oblong-ovate cuspidate and ciliate at the point, fr. downy oblong-ovoid narrowed into an obliquely truncate beak with 2 lateral points, nut narrowly elliptic subtrigonous, l. slender channelled.—*E. B.* 904. *Schk.* K. 45. *B.* 132. *R.* 265.—St. 2 feet high. L. with filamentous sheaths below. Lowest spike rarely stalked.—Peat-bogs, rare. P. V. E. S. I.

65. *C. hir'ta* (L.); male spikes 2 or 3, fertile 2 or 3 remote erect oblong-cylindrical stalked, bracts leaflike the lower with long sheaths nearly equalling the peduncles, gl. elliptic-lanceolate with long slender ciliate points, fr. hairy oblong-ovate narrowed into a deeply divided beak, nut obovate narrowed below triangular, l. flat hairy.—*E. B.* 685. *Schk.* Uu. 108. *H. b.* 58. *R.* 257.—St.  $1\frac{1}{2}$ —2 feet high, leafy. L. and sheaths shaggy, rarely glabrous. Fr. tawny. Occasionally the spikes are compound at the base and very long-stalked, and the gl. long.—Wet places. P. IV. *Hammersedge.* E. S. I.

††††† Beak of fr. glabrous, terete, striate, with patent cusps, or 2-toothed. Male spikes many, rarely 1. Bracts not sheathing.

† Male spike 1. Stigmas 2. Beak of fr. 2-toothed.

66. *C. pul'la* (Good.); male spike 1 (rarely 2), fertile 1—3 roundish-ovoid lower one stalked bracteate sheathless erect, fr. ovoid obscurely veined inflated, beak short, nut roundish mucronate, gl. bluntish.—*E. B.* 2045. *Schk.* Cc. 88. *C. vesicaria* v. *alpigena* Fr.—St. 6—8 in. high. Gl. dark purple tipped with white, midrib dark purple. Fr. dark purple paler at the base, longer than the gl., stalked.—*C. saxatilis* (Linn. Herb.); but

Andersson says that the true plant is *C. rigida*.—Wet parts of the higher Scottish mountains. P. VII. S.

67. *C. Grahámi* (Boott); male spikes 1 or 2 slender acute, fertile 2 or 3 ovate blunt lower one stalked bracteate sheathless, fr. oblong-ovate strongly ribbed inflated narrowed into a short bifid beak, nut oblong compressed triquetrous below not  $\frac{1}{2}$  as long as the perianth, gl. acute.—*E. B. S.* 2923.—St. 1—2 ft. high. Gl. fuscous, with the tip and midrib pale. Fr. pale or darkish brown, nearly twice as long as the gl., with several strong ribs on each side.—Glen Fiadh, Clova. Ben Cruichben, Killin. P. VII. S.

†† Male spikes only. Stigmas 3. Bracts not sheathing.

68. *C. ampullácea* (Good.); fertile spikes 2—4 remote cylindrical erect stalked, bracts leaflike, fr. subglobose inflated suddenly narrowed into a long slender beak, nut obovoid triangular, st. smooth with blunt angles.—*E. B.* 780. *Schk.* Tt. 107. *H. b.* 65. *R.* 277.—St. 1—2 feet high, trigonous. L. glaucous, channelled.—Very wet bogs. P. VI. E. S. I.

69. *C. vesicária* (L.); fertile spikes 2—4 remote cylindrical, bracts leaflike, fr. ovoid conical inflated gradually narrowed into a subulate bifid beak, nut elliptic triangular, st. with acute angles.—*E. B.* 779. *Schk.* Ss. 106. *H. b.* 64. *R.* 276.—St. 2 feet high, triangular, roughish near the top. L. rather broad, green.— $\beta$ . *involuta* (Bab.); l. narrow folded into a  $\frac{1}{2}$ -cylinder, midrib of gl. apiculate, fr. narrower.—*Sy. E. B.* 1681.—Wet bogs.  $\beta$ . Hale Moss, Manchester. *Mr. J. Sidebotham.* Congleton, Chesh. *Mr. E. Wilson.* P. V. E. S. I.

70. *C. paludósa* (Good.); gl. of the barren spikes blunt, anth. apiculate, fertile spikes cylindrical blunt, bracts leaflike, fr. oblong-obovate compressed with a short bifid beak, nut roundish-obovate triangular, st. with acute angles.—*E. B.* 807. *Schk.* Oo. 103.—St. 2—3 feet high, angles rough. L. broad. Fr. sometimes recurved at the tip, stig. sometimes 2.— $\beta$ . *C. spadicéa* (Elw., Roth.); gl. of fertile spikes with a long rough beak.—Wet places. P. V. E. S. I.

71. *C. ripária* (Curt.); gl. of the barren spikes acute, anth. with a long point, fertile spikes acute cylindrical, bracts leaflike, fr. oblong-ovoid convex on both sides narrowed into a short broad cloven beak, nut pyriform triangular, st. with acute angles.—*E. B.* 579. *Schk.* Qq. & Rr. 105. *H. b.* 66.—St. 3—4 feet high, angles rough. L. broader than in the preceding.—Wet places. P. V. E. S. I.

*Index to the Carices.*

- C. acuta*, 26.  
*alpigena*, 66.  
*alpina*, 30.  
*ampullacea*, 68.  
*aquatilis*, 28.  
*arenaria*, 7.  
*atrata*, 32.  
*axillaris*, 17.  
*binervis*, 59.  
*Boeninghausenia*,  
   16.  
*brizoides*, 15\*.  
*Buxbaumii*, 31.  
*cæspitosa*, 25, 29.  
*canescens*, 21.  
*capillaris*, 39.  
*ciliata*, 46.  
*clandestina*, 42.  
*collina*, 47.  
*curta*, 21.  
*curta* v. *alpicola*, 22.  
*Davalliana*, 2.  
*depauperata*, 61.  
*digitata*, 44.  
*dioica*, 1.  
*distans*, 58.  
*disticha*, 6.  
*divisa*, 9.  
*divulsa*, 12.  
   *Ehrhartiana*, 13.  
*elongata*, 20.  
*ericetorum*, 46.  
*extensa*, 55.  
*filiformis*, 64.  
*flava*, 53.  
*frigida*, 52.  
*Gibsoni*, 29\*.  
*glauca*, 50.  
   *Goodenovii*, 29.  
*Grahami*, 67.  
*hirta*, 65.  
*Hornschuchiana*, 56.  
*humilis*, 42.  
*incurva*, 8.  
*intermedia*, 6.  
   *involuta*, 69.  
*irrigua*, 37.  
*lævigata*, 60.  
*lagopina*, 23.  
   *Leesii*, 48.  
   *lepidocarpa*, 53.  
*leporina*, 23, 24.  
*limosa*, 36.  
   *Meilichoferi*, 35.  
   *Micheliana*, 50.  
*montana*, 47.  
*muricata*, 11.  
*Cederi*, 54.  
*ornithopoda*, 43.  
*ovalis*, 24.  
*pallescens*, 33.  
*paludosa*, 70.  
*panicea*, 34.  
*paniculata*, 15.  
*paradoxa*, 14.  
*pauciflora*, 5.  
*pendula*, 41.  
   *Persoonii*, 22.  
   *phæostachya*, 35.  
*pilulifera*, 48.  
*præcox*, 45.  
*prolixa*, 26.  
*Pseudo-cyperus*, 63.  
*pulicaris*, 3.  
*pulla*, 66.  
*punctata*, 57.  
*rariflora*, 38.  
*recurva*, 50.  
*remota*, 18.  
*rigida*, 27.  
*riparia*, 71.  
*rupestris*, 4.  
*saxatilis*, 66.  
   *spadicea*, 70.  
*speirostachya*, 56.  
*stellulata*, 19.  
   *stictocarpa*, 50.  
*stricta*, 25.  
*strigosa*, 40.  
*sylvatica*, 62.  
*tenella*, 18.  
*teretiuscula*, 13.  
*tomentosa*, 49.  
*ustulata*, 51.  
*vaginata*, 35.  
   *Vahllei*, 30.  
*vesicaria*, 69.  
*vesicaria* v. *alpigena*,  
   66.  
*vitis*, 22.  
*vulgaris*, 29.  
*vulpina*, 10.  
*xanthocarpa*, 56\*.

## Order XCVIII. GRAMINEÆ.

Fl. perfect or unisexual, 1, 2 or more seated bifariously on a common axis which is contained within an involucre of 2 (or 1) valves (glumes) or rarely none, the whole forming a locusta or spikelet. Each fl. of 1 or 2 scales (pales) of which the outer or lower is simple and usually keeled, the inner with 2 veins or keels. Hypogynous scales 2, 3 or none. Stam. hypogynous, 1—6, usually 3. Anth. versatile, notched at both ends. Ovary

1-celled. Styles usually 2, rarely 1 or 3. Embryo on the outside of the albumen and at its base.—L. with split sheaths<sup>1</sup>.

### Suborder I. CLISANTHÆ.

Flowers closed. Styles or stigmas long, protruded at or near to the top of the flower.

#### A. *Rachis of inflorescence without lateral excavations.*

Tribe I. *PANICEÆ*. *Spikelets dorsally compressed, 1-flowered, or with 1 fl. and an inferior glumelike rudiment or a neuter flower. Lower gl. much the smaller, often rudimentary.*

1. *DIGITARIA*. *Spikes fingered*. Spikelets in pairs on one side of the flattened rachis, awnless, 1-flowered, with an inferior rudiment. Gl. 2, lower smaller or 0, upper 3-veined. Sterile fl. of one 5—7-veined pale equalling the flower.

[2. *ECHINOCHLOA*. *Spikes compound, secund in the whole and in each part. Spikelets on one side of the flattened partial rachis, 2-flowered, inferior fl. rudimentary. Gl. 2, lower 3-veined, upper equalling fl. 5-veined mucronate or awned. Lower pale of sterile fl. like and equalling upper glume.*]

3. *SETARIA*. *Spike cylindrical, compound. Spikelets surrounded by an involucre of bristles, 2-flowered; inferior fl. rudimentary. Gl. 2, lower 3-veined, upper equalling fl. many-veined. Sterile fl. of 1 pale, like and equalling upper glume.*

Tr. II. *CHLORIDEÆ*. *Spikelets laterally compressed 1-flowered in our plants and sometimes with a superior rudiment, placed in 2 rows on one side of a flattened rachis, or alternate and unilateral.*

4. *CYNODON*. *Spikes fingered, spreading. Spikelets 1-flowered, awnless, with a superior rudiment. Gl. nearly equal, patent. Pales equal; lower boatshaped, compressed, embracing the inner. Styles long, distinct. Stigmas feathery.*

5. *SPARTINA*. *Spikes upright, in a raceme. Spikelets 1-*

<sup>1</sup> Nearly all the genera of this Order are beautifully figured in the *Gen. Fl. Germ. Monocotyl.* vol. i. See also Andersson's *Pl. Scand.* fasc. ii. and Du Mortier in *Bull. Soc. Bot. Belg.* vii. 65.

flowered, awnless. Gl. unequal; upper lanceolate. Pales unequal; lower boatshaped, compressed. *Styles* long, united halfway up. Stigmas feathery.

6. MIBORA. *Inflorescence a somewhat 1-sided raceme.* Gl. not keeled blunt. Fl. 1. Pale 1, scarious, very hairy, blunt, not awned.

Tr. III. PHALARIDÆ. Panicked. *Spikelets laterally compressed, 1-flowered,* with 1 or 2 glumelike inferior rudiments, or 1 or 2 inferior male flowers. Gl. equal covering the flowers. *Styles* long. Stigmas filiform.

7. PHALARIS. Gl. 2, boatshaped, keeled, membranous, nearly equal, exceeding the flower. Pales coriaceous, unequal, closely investing the fruit. Rudimentary fl. 1—2, scalelike.

8. ANTHOXANTHUM. Gl. 2, unequal, membranous; lower small, 1-veined; upper exceeding the fl., 3-veined. Pales scarious. *Stam.* 2. Rudimentary fl. 2, scalelike, bifid, awned on the back.

9. HIEROCHLOE. Gl. 2, nearly equal, membranous, about equalling the spikelet. Fl. 3; 2 lower male, 3-androus, upper pale with 2 keels; upper fl. perfect, 2-androus, upper pale with 1 keel.

Tr. IV. PHELINEÆ. *Inflorescence dense, spikelike.* Spikelets laterally compressed, 1-flowered or with a superior rudiment. Gl. nearly equal, covering the flowers. *Styles* long. Stigmas filiform.

10. PHELEUM. Gl. compressed, keeled, parallel at the midrib, truncate, with a terminal seta, or acute. Fl. 1. Pales 2, membranous; lower 3-veined, blunt, without awns, or with a minute central point.

11. ALOPECURUS. Gl. compressed, connate below, membranous, awnless. Fl. 1. Pale 1, scarious, 5-veined, awned on the back.

Tr. V. SESLERIÆ. Panicle spikelike. Spikelets laterally compressed, with 2 or more flowers. *Styles* 0 or very short. Stigmas very long, filiform.

12. SESLERIA. Spikelets sessile, imbricate all round. Gl. 2—6-flowered, nearly or quite equalling the spikelet. Lower pale keeled, membranous, with a scarious margin, ending in 3 or 5 points; dorsal rib excurrent.

B. *Spikelets sessile in hollows of the rachis.*

Tr. VI. *NARDEÆ*. Spikelets 1-flowered. Gl. 0. Style short. Stigmas filiform.

13. *NARDUS*. Spikelets in 2 rows on one side of the rachis. Lower pale keeled, tapering into a subulate point. Stigmas long.

Suborder II. *EURYANTHÆÆ*.

Flowers open. Style short. Stigmas protruded near to the bottom of the flower.

A. *Inflorescence paniced or racemose; rachis without lateral excavations.*

Tr. VII. *ORYZEÆ*. Spikelets laterally compressed, 1-flowered, Glumes 0. Stigmas feathery. Pales enclosing but free from the nut.—Pales opening only slightly.

14. *LEERSIA*. Pales 2, like parchment, compressed, keeled, awnless; lower much broader.

Tr. VIII. *STIPACEÆ*. Spikelets cylindric, 1-flowered, without any rudiment. Gl. unequal membranous, enclosing the flowers. Pales hardening, enclosing but free from the nut.

[15. *STIPA*. Spikelets stalked. Pales coriaceous; outer cylindrical, convolute, evidently jointed to the kneed twisted feathery awn.]

16. *MILIUM*. Spikelets stalked. Pales like parchment, awnless; lower ventricose, convex.

Tr. IX. *AGROSTIDEÆ*. Spikelets laterally compressed, 1-flowered, or with a superior rudiment, or many-flowered. Gl. and pales membranous.

† *Gl. falling short of the outer pale. Style long. Stigmas often protruded near the middle of the flower.*

17. *PHRAGMITES*. Pan. diffuse. Gl. unequal; lower much smaller. Fl. 1—6, awnless, with silky hairs at their base; lower imperfect. Lower pale acuminate, much exceeding the inner, awnless.

†† *Gl. exceeding the flowers. Style short or none.*

18. *PSAMMA*. Pan. spikelike. Gl. nearly equal; lower rather the shorter. Fl. 1, with silky hairs at its base, or without a superior rudiment. Lower pale with a very short awn.

19. CALAMAGROSTIS. Pan. diffuse. Gl. nearly equal; lower rather the longer. Fl. 1, with silky hairs at its base, with or without a superior rudiment. Lower pale awned.
20. APERA. Pan. loose. Gl. membranous, acute, unarmed; *lower the smaller*. Fl. 1, with hairs at its base, and a pedicel-like rudiment. Pales unequal, scarious; lower with a long subterminal awn.
21. AGROSTIS. Pan. loose. Gl. membranous, acute, unarmed; *upper smaller*. Fl. 1, with hairs at its base; no rudiment. *Pales unequal*, scarious; dorsal awn falling short of the glumes, or 0.—*A. canina* wants the inner pale.
- [22. LAGURUS. Pan. spikelike. Gl. scarious, *ending in a long fringed bristle*. Fl. 1, with a pedicel-like rudiment. Lower pale ending in 2 long bristles and with a dorsal kneed twisted awn.]
23. POLYPOGON. Pan. close, spikelike. Gl. scarious, each *with a long bristle* from just below the notched tip. Fl. 1. Lower pale usually awned from just below the tip.
24. GASTRIDIVM. Pan. close, spikelike. Gl. membranous, acute, awnless, *ventricose* at the base, much exceeding the flower. Fl. 1. Lower pale truncate or toothed at the end, with or without a dorsal awn.
- Tr. X. AVENEÆ. Spikelets with 2 or more flowers; upper often barren. Gl. *equalling or overtopping the flowers*. Lower pale awned. Style short or 0.
25. HOLCUS. Fl. 2; lower perfect, awnless (or very rarely awned); upper usually male, with a dorsal awn. *Pales hardening on the fruit*; tip of lower entire.
26. CORYNEPHORUS. Fl. 2, perfect, awned. Awn straight, jointed in the middle; *the upper portion clavate*; a tuft of hairs at the joining. Tip of lower pale entire.
27. AIRA. Pan. lax. Fl. 2, with or without the rudiment of a third (sometimes perfected in *A. cæspitosa*). Lower pale denticulate or bifid at the tip, terete on the back. Awn dorsal, kneed (in *A. cæspitosa* straight). Ovary glabrous. Fr. not crested.
28. TRIVSETUM. Spikelets crowded. Fl. 2—6. Lower pale with faint lateral veins, ending in 2 acute teeth, awned. Awn dorsal, kneed and twisted. Ovary glabrous. Fr. *neither crested nor furrowed*.



29. AVENA. Fl. 2 or more. Lower pale with lateral veins, awned, ending in 2 points. Awn dorsal, kneed and twisted. Ovary hairy at the top. *Fr. crested and furrowed.*

30. ARRHENATHERUM. Fl. 2, with a rudiment; *lower fl. male* with a long kneed and twisted awn from below its middle; upper with a short straight awn from near the tip. Pales herbaceous, ending in 2 points. Ovary hairy at the top. *Fr. oblong, terete, downy, not furrowed.*

Tr. XI. FESTUCEÆ. Spikelets with 2 or more flowers, upper often barren. *Gl. falling short of the lowest flower. Style short or 0.*

† *Lower pale with nearly parallel veins which do not join to form an awn. Awn 0. Styles terminal.*

‡ Lower pale 2—3-fid.

31. TRIODIA. Fl. 2—4. Lower pale rather coriaceous, rounded on the back, bifid with an intermediate broad point. Nut free.

‡‡ Lower pale nearly or quite entire.

32. KÆLERIA. *Gl. unequal; upper 2—3-ribbed. Spikelet compressed, 2—5-flowered. Lower pale keeled, acuminate, or with a straight subterminal bristle. Nut free.*

33. MELICA. *Gl. nearly equal, with lateral ribs, nearly as long as the ovate spikelet consisting of 1 or 2 flowers rounded on the back and a clublike rudiment. Pales hardening on the free nut.*

34. MOLINIA. *Gl. unequal, without lateral ribs, falling short of the lanceolate spikelet of 2 or 3 semicylindrical flowers and a subulate rudiment. Pales hardening on the free nut.*

35. POA. *Gl. rather unequal. Lower pale 3—5-veined, scarious at the tip, compressed, keeled. Nut elliptic trigonous, slightly furrowed within, free.*

36. GLYCERIA. *Gl. unequal, submembranous. Lower pale with 5—7 strong prominent ribs and a scarious margin, sub-cylindrical. Nut oblong, convex on back, furrowed within, free.*

37. SCLEROCHLOA. *Gl. unequal, membranous. Lower pale with 5 faint veins, cylindrical below, often keeled at the tip*

or with a very minute mucro. Nut oblong, convex on back, not furrowed within, free.

38. BRIZA. Gl. nearly equal, broad, 3-ribbed. Fl. 3—8, densely imbricate in a short distichous spikelet. Lower pale boatshaped, heartshaped, blunt, rounded on the back. Gl. and pales membranous with a scarious margin. Nut free.—Lower pale with 7—9 faint veins.

‡‡‡ Lower pale truncate and denticulate at the tip.

39. CATABROSA. Gl. unequal, very short, 1-veined. Flowers usually 2, rounded on the back, distant. Lower pale membranous, with 3 veins ending in teeth which do not quite extend to the edge of the scarious margin. Upper gl. often with 2 short and faint lateral veins.

†† Lower pale with converging veins, all or 1—3 of which combine in the awn.

40. CYNOSURUS. Gl. nearly equal, scarious, strongly keeled, 1- or more flowered. Lower pale rounded on the back with a terminal bristle. *Each spikelet with a comb-like bract at its base.* Panicle spikelike, 1-sided. Nut closely coated with the pales.

41. DACTYLIS. Gl. unequal, herbaceous, many-flowered; lower keeled. *Lower pale compressed, keeled, 5-veined; dorsal vein fringed and excurrent just below the tip as a short awn. Spikelets crowded, subsecund.* Nut free.

42. FESTUCA. Gl. unequal, herbaceous, many-flowered. *Lower pale rounded on the back, very acute, or with the dorsal vein excurrent at or just below the tip as a short awn; lateral veins slightly converging and vanishing below the tip.* Upper pale minutely ciliate on the ribs. *Styles terminal.* Nut furrowed, adhering to the pales.—Rachis with acute angles. Sheaths of the leaves divided to the base.

43. BROMUS. Gl. unequal, herbaceous, many-flowered; *lower 1-veined, upper 3—5-veined.* Fl. lanceolate, compressed. Lower pale with a long awn, (usually) founded on 3 veins, from below the tip. *Styles below the top of the ovary.* Nut furrowed, adhering to the pales.—Sheaths of the l. divided halfway down.

44. SERRAFALCUS. Gl. unequal, herbaceous, many-flowered, *lower 3—5-veined, upper 7—9-veined.* Fl. oblong, turgid. Lower pale with a short awn, (usually) founded on 3 veins, from below the tip. *Styles below the top of the ovary.* Nut

furrowed, adhering to the pales.—Sheaths of the 1. scarcely divided halfway down. Spikelets narrower upwards.

B. *Inflorescence spike-like ; bearing the spikelets in hollows of the rachis.*

Tr. XII. *HORDEIÆ*. Spikelets solitary or 2 or 3 together, subsessile on opposite sides of a channelled and toothed jointed rachis. Uppermost fl. often barren. Style very short. (Lateral spikelets often stalked in *Hordeum*.)

† *Spikelets very shortly stalked or subsessile.*

45. *BRACHYPODIUM*. Gl. opposite, unequal, many-flowered, their edges towards the rachis. Upper pale coarsely fringed on the ribs.—The unequal gl. distinguish this from *Triticum*.

†† *Spikelets quite sessile.*

‡ Glumes 2.

46. *TRITICUM*. Spikelets solitary. Gl. opposite, nearly equal, many-flowered, their edges towards the rachis. Inner pale minutely ciliate on the ribs.

47. *ELYMUS*. *Spikelets 2 or 3 together. Gl. 2, both on the same side of the spikelet, without awns or bristles, with 2 or more perfect flowers.*

48. *HORDEUM*. Spikelets in threes, often partially barren. *Gl. 2, ending in long bristles ; 1 perfect flower and a stalk-like rudiment.*

49. *LEPTURUS*. Spikelets solitary, imbedded alternately on opposite sides of the rachis. Gl. 1—2, both on the same side of the spikelet, cartilaginous, covering the one fl. and superior rudiment. Pales scarious. Stigmas feathery.

‡‡ Glume solitary, bractlike, or a very small upper one.

50. *LOLIUM*. Spikelets solitary, placed edgewise on the rachis. Gl. solitary, or that next the rachis very small, with 3 or more flowers.

Suborder I. *Clisanthææ*. Tribe I. *Paniceæ*.

1. *DIGITA'RIA Scop.*

[*D. sanguinalis* (Scop.); l. and sheaths hairy, fl. oblong-lan-

ceolate glabrous with downy margins (?).—*E. B.* 849. *P.* 70.—  
St. ascending, a foot long.—Not a native. *A.* VIII.] *E.*

1. *D. humifusa* (Pers.); l. and sheaths glabrous, fl. ovate  
downy with glabrous veins.—*E. B. S.* 2613. *P.* 71.—St. mostly  
procumbent, 4—8 in. long. Spikes usually 3 or 4, springing  
from nearly the same point. Spikelets in pairs, one on a longer  
stalk than the other.—Sandy fields, rare. *A.* VII. VIII. *E.*

## 2. ECHINOCH'LOA *Pal. de Beauv.*

[*E. Crus-galli* (Beauv.); spikes alternate or opposite, spike-  
lets near together, upper gl. and sterile floret awned or mucro-  
nate hispid, rachis hispid.—*E. B.* 876. *P.* 67. *Panicum* L.—  
Near London. *A.* VII.] *E.*

## 3. SETA'RIA *Pal. de Beauv.*

1. *S. viridis* (Beauv.); pan. spikelike, involucrel bristles with  
forward teeth, lower pale smooth.—*Panicum* L., *E. B.* 875.  
*P.* 68.—London and Norwich. *A.* VII. VIII. *E.*

[*S. verticillata* (Beauv.); pan. spikelike, involucrel bristles  
with declining teeth, lower pale smooth.—*E. B.* 874. *P.* 69.  
—London and Norwich. *A.* VII. VIII.] *E.*

[*S. glauca* (Beauv.); pan. spikelike, invol. bristles with as-  
cending teeth, pales transversely rugose.—Weybridge, Surr.  
*A.* IX.] *E.*

## Tribe II. Chlorideæ.

### 4. CYN'ODON *Rich.*

1. *C. Dactylon* (Pers.); spikes 3—5 digitate, pales smooth  
their edges and keel slightly ciliate, l. downy beneath, barren  
shoots prostrate.—*E. B.* 850. *P.* 72.—Creeping. Flowering st.  
4—6 in. high, ending in a cluster of spreading many-fl. slender  
spikes. Spikelets purplish. L. on the long branched barren  
shoots flat, spreading; on the others usually folded.—Sandy  
shores. Dorset. Devon. Cornwall. *P.* VIII. *E.*

### 5. SPARTI'NA *Schreb.*

1. *S. stricta* (Roth); l. jointed to their sheaths falling short  
of the spikes, spikes 2—3, rachis scarcely extending beyond the  
last spikelet, outer gl. hairy.—*E. B.* 380.—St. 1—2 feet high.

L. narrowing to the base where they easily separate from their sheaths. Spikes pressed close together. A remarkably rigid plant.—Muddy salt marshes. P. VIII. E.

‡2. *S. alterniflora* (Loisel.); l. continuous with their sheaths equalling or exceeding the spikes, spikes many, *rachis produced beyond the spikelets* and flexuose, outer gl. glabrous.—*E. B. S.* 2812. *P.* 75.—St. 2—3 feet high. L. broadest at the base and not separating from the sheaths more easily than at any other part. Spikes pressed close together.—Mud-flats in the river Itchin at Southampton. P. VIII. E.

#### 6. MIBORA *Adans.*<sup>1</sup>

1. *M. minima* (Desv.).—*Knappia Sm. E. B.* 1127. *P.* 73. *Sturmia Hoppe in St.* 7. 1. *Chamagrostis Borkh.*—An elegant but very small grass. Root small, fibrous. St. many. L. short, rough. Spikes slender; spikelets 5—10, sessile. Pale shorter than the glumes, hairy, truncate, ragged.—Sandy southwest coast of Anglesea. A. IV. E.

### Tribe III. *Phalarideæ.*

#### 7. PHALARIS *Linn.* Reed-grass.

[*P. canariensis* (L.); pan. ovoid spikelike, *gl. winged on the keel*, wing entire, rudimentary fl. 2 half as long as the fertile fl., pales pilose.—*E. B.* 1310. *P.* 9.—St. 1—2 feet high, ending in a compact compound panicle. Gl. large, pale yellow variegated with green lines and remarkably winged at the back.—Scarcely naturalized. A. VII. *Canary-grass.*]

[*P. paradox'a* (L.), pan. spikelike, gl. of fertile fl. with a blunt toothed *wing on the keel* many-veined, rudimentary fl. several much shorter.—St. decumbent below, then ascending, 1—3 ft. high, branched. Lower part of pan. usually barren; branches with about 6 spikes.—Swanage, Dorset. A. VII.] E.

1. *P. arundinæa* (L.); pan. upright with spreading branches, fl. clustered, *gl. not winged*, rudimentary fl. 1 or 2 small hairy.—*E. B.* 402. *P.* 9.—St. 4—5 feet high. Creeping. Pan. 3—4 in. long. Gl. keeled. L. sometimes variegated with white lines.—By water. P. VI. VII. E. S. I.

#### 8. ANTHOXANTHUM *Linn.* Vernal-grass.

1. *A. odoratum* (L.); pan. spikelike dense oblong, gl. about

<sup>1</sup> *Mibora Adans.* 1763, *Chamagrostis Borkh.* 1790.

equalling awns, st. nearly simple below.—*E. B.* 647. *P.* 8.—About a foot high. Panicle lanceolate, dense, or rather interrupted below.—There are two forms or species: (1) with purple anth. in meadows, (2) with dull yellow anth. in woods.—Very common in pastures. *P. V.* VI. E. S. I.

2. *A. Puel'lii* (Lec. & Lam.); pan. spikelike lax, gl. much falling short of awns, st. many branching from the base.—*J. of B.* xiii. t. 157.—Pan. narrowed to the top. St. 6—12 in. high.—Sandy pastures. *A.* VII.—IX E.

#### 9. *HIERO'CHLOE Gmel.* Holy-grass.

1. *H. borealis* (R. & S.); pan. divaricate, pedicels glabrous, fl. awnless, l. flat.—*E. B. S.* 2641. *P.* 31.—About a foot high. Spikelets ovate, brown.—Forfarshire (but now lost). Thurso, Caithness. *Mr. R. Dick.* *P. V.* VI. S.

### Tribe IV. *Phleineæ.*

#### 10. *PHLE'UM Linn.* Cat's-tail-grass.

[*P. as'perum* (Jacq.); l.-sheaths slightly inflated, pan. cylindrical, gl. wedgeshaped truncate swelling upwards, keels rough, rudimentary fl. subulate.—*E. B.* 1077. Probably an escape, now lost. *A.* VII.] E.

1. *P. Boeh'meri* (Wibel); l.-sheaths slightly inflated, pan. cylindrical, gl. linear-lanceolate obliquely truncate mucronate, keel ciliate above, rudimentary fl. subulate.—*E. B.* 459. *P.* 80.—St. leafy below, naked upwards, with sterile leafy shoots. Lower pale entire, not awned. Anth. linear-oblong.—Dry chalky fields, rare. *P.* VII. E.

[*P. Michélli* (All.) was probably a mistake.]

2. *P. arenarium* (L.); l.-sheaths inflated, pan. oblong somewhat narrowed below, gl. lanceolate acuminate, keel ciliate above, rudimentary fl. subulate minute.—*E. B.* 222. *St.* 29. 1. *P.* 7.—St. varying greatly in height. Lower pale notched at the summit,  $\frac{1}{3}$  the length of the glumes. Anth. very small.—Sandy places chiefly near the sea. *A.* V.—VII. E. S. I.

3. *P. pratense* (L.); l.-sheaths not inflated, pan. cylindrical, gl. oblong truncate with an awn of less than half their length, keel ciliate above, no rudimentary flower.—*E. B.* 1076. *P.* 77 & 78.—Slightly creeping (or slightly tuberous, *P. nodosum* L.). Pan. 1—5 in. long. Awns sometimes longer. Lower pale jagged at the summit.—*a. P. pratense* (L.); st. from a prostrate base, l. broad, pan. usually long, anth. purplish.—*β. sto-*

*lomiferum* (Bab.); many barren leafy prostrate or erect shoots, l. broad, pan. rather short, anth. purplish.—*γ. P. præcox* (Jord.); st. from an ascending base, l. slender, pan. rather short, anth. pale yellow.—Meadows and pastures. P. VI. *Timothy-grass*.  
E. S. I.

4. *P. alpinum* (L.); l.-sheaths inflated, pan. oblong, gl. truncate glabrous ciliate on the back with a scabrous awn nearly equalling their length, keel ciliate, upper sheath inflated.—*E. B.* 519. P. 6. *P. commutatum* Gaud.—Somewhat creeping. St. 6—12 in. high.—Wet alpine moors. P. VII. S.

### 11. ALOPECURUS Linn. Fox-tail-grass.

1. *A. pratensis* (L.); st. erect smooth, pan. cylindrical blunt, gl. acute connected below ciliate downy, pale equalling the glumes, awn projecting more than half its own length beyond the pale.—*E. B.* 759. St. 8. l. P. 4.—Scarcely creeping.—St. 1—3 feet high. In salt marshes the base of the st. becomes fleshy. Upper sheath slightly inflated. Ligule short, blunt. Pan. 1—3 in. long, branches 4—6-flowered. Anth. yellow. Styles combined.—Rich pastures. P. IV.—VI. E. S. I.

2. *A. alpinus* (Sm.); st. erect, smooth, pan. oblong, gl. acute connected below hairy, pale equalling the glumes, awn projecting  $\frac{1}{2}$  of its length beyond the pale.—*E. B.* 1126. P. 4.—Somewhat creeping. St. decumbent at the base, then erect, 9—12 in. high. Ligule short, blunt. Uppermost l. usually (not always) short and broad,  $\frac{1}{3}$  of the length of its inflated sheath. Gl. connate through about  $\frac{1}{4}$  of length. Awn from about the middle of pale, sometimes wanting. Styles combined. Pan. not exceeding an inch, sometimes rather lax; silky branches 4—6-flowered.—High on mountains. Loch-na-Gar. Ben Lawers. Clova. P. VII. S.

3. *A. geniculatus* (L.); st. ascending bent at the knots smooth, pan. cylindrical, gl. blunt connected below ciliate rather exceeding the pale, awn from near the base of the pale and projecting half its length beyond it, anth. linear.—*E. B.* 1250. P. 5.—Root fibrous. St. about a foot long, branching below. Knots generally (in dry places) oval and fleshy. Upper sheath inflated. Ligule oblong. Gl. membranous at the top except the midrib, often villose or hairy below. Pale when laid open oblong, blunt, slightly notched. Anth. ultimately violet-yellow. Styles mostly combined. Pan. 1—2 in. long.—*A. pronus* (Mitten) is a prostrate form of this.—Wet places. P. VI. VII. E. S. I.

4. *A. fulvus* (Sm.); st. ascending bent at the knots smooth, pan. cylindrical, gl. connected below ciliate rather falling short

of the pale, awn from just below the middle of the pale and scarcely extending beyond it, *anth.* short and broad.—*E. B.* 1467. *P.* 5.—*St.* 1—2 feet long, procumbent below. Ligule oblong. *Pan.* 2—3 in. long. *Anth.* at first white, afterwards orange-coloured.—Wet margins of ponds. *P.* VI.—IX. E. S.

5. *A. bulbosus* (L.); *st.* smooth, *pan.* cylindrical acuminate, *gl.* distinct abruptly acute downy exceeding the pale, awn from near the base of the pale and projecting half its length beyond it.—*E. B.* 1249. *P.* 76.—*St.* 1 foot long, ascending or decumbent, in a circular tuft, kneed, the lowermost knots forming *ovate fleshy knobs*. Upper sheath inflated. Ligule oblong. Pale when laid open truncate, emarginate, with 2 small teeth in the middle. Styles combined. *Pan.* about 1 in. long, less decidedly racemose than in our other species; pedicels usually 1-flowered.—Salt marshes in the South, rare. *P.* V. VI. E.

6. *A. agrestis* (L.); *st.* erect roughish above, *pan.* tapering slender, *gl.* acute connected below nearly glabrous, awn from near the base of the pale and projecting half its length beyond it.—*E. B.* 848. *P.* 3.—*St.* 1—2 feet high, slender. Sheaths roughish. Ligule prominent, blunt. *Gl.* glabrous, but with a row of *fine short cilia on the back*, connate nearly to the middle. Styles combined.—A very troublesome weed. *A.* IV.—XI. *Black grass.* E.

### Tribe V. *Seslerieæ*.

#### 12. *SES'LERIA Scop.* Moor-grass.

1. *S. cærulea* (Scop.); *pan.* ovoid slightly 1-sided, outer pale ending in 4 teeth, midrib rough with a short excurrent point, l. abrupt with a minute rough point.—*E. B.* 1613. *P.* 27.—Roots tufted. *St.* 6—12 in. high. *Pan.* about  $\frac{1}{2}$  in. long, bluish purple. *Anth.* purple-tipped. *Stig.* very long, linear.—Mountains. Banks of the Shannon. *P.* IV. V. E. S. I.

### Tribe VI. *Nardeæ*.

#### 13. *NAR'DUS Linn.* Mat-grass.

1. *N. stricta* (L.).—*E. B.* 290. *P.* 2.—Tufted. *St.* and l. erect, slender, rigid. Height 5—8 in. Spike close, slender. Lower pale with a short rough awn, coriaceous, often purplish; upper membranous.—Moors and heaths. *P.* VII. E. S. I.



Suborder II. *Euryantheæ*. Tribe VII. *Oryzæ*.14. LEER'SIA *Soland.* Cut-grass.

1. *L. oryzoides* (Sw.); pan. patent with wavy branches, spikelets 3-androus half-oval, keel ciliate.—*E. B. S.* 2908.—Creeping. St. 1—2 ft. high. L. broad, rough-edged; uppermost horizontal at the flowering-season. Pan. mostly enclosed in the sheath of the uppermost leaf. The included fl. alone are usually fertile.—Marsh-districts in Sussex, Surrey, and Hampshire. P. VIII. IX. E.

Tribe VIII. *Stipaceæ*.15. STI'PA *Linn.* Feather-grass.

[*S. pennata* (L.); awn very long twisted feathery its base glabrous.—Awns remarkably long.—Not a native. Common in gardens. P. VI.] E.

16. MIL'IUM *Linn.* Millet-grass.

1. *M. effusum* (L.); pan. diffuse, pales acute, st. smooth, l. lanceolate-linear.—*E. B.* 1106. P. 17.—Stoloniferous. St. 3—4 feet high. Branches of the panicle long, in distant alternate tufts, in flower horizontal, afterwards deflexed.—Damp shady woods. P. VI. E. S. I.

Tribe IX. *Agrostideæ*.17. PHRAGMI'TES *Trin.* Reed.

1. *P. communis* (Trin.); pan. diffuse, spikelets 1—6-flowered, fl. exceeding the glumes.—*Arundo* L., *E. B.* 401. P. 29.—St. 5—6 feet high, erect. Pan. large, purplish. Spikelets usually 3—6-fl., or 1—2-fl. in *P. nigricans* Dum. L. flat, broad. Soboliferous; soboles rarely aerial and 20—40 ft. long (*Phytol.* i. 146).—Marshes. P. VIII. E. S. I.

18. PSAM'MA *Pal. de Beauv.* Marram. Bent.

1. *P. arenaria* (R. & S.); pan. cylindrical rather thicker at the middle, gl. and pales linear-lanceolate acute, hairs  $\frac{1}{2}$  the length of the pale.—*Arundo* L., *E. B.* 520. *Ammophila* (Host) P. 8.—St. erect, stiff, 2—3 feet high. L. rigid, involute, acute, glaucous. Panicle straw-coloured.—Sandy sea-shore, binding the shifting sands. P. VII. E. S. I.

2. *P. baltica* (R. & S.); pan. lanceolate, gl. and pales lanceolate-prolonged very acute, hairs  $\frac{1}{2}$  the length of the pale.—*J. of B. x. t. 127.*—Known by its more lax sublanceolate pan. and very acute glumes.—Ross Links, Northumb. P. VII. E.

### 19. CALAMAGROS'TIS Adans. Small Reed.

1. *C. lanceolata* (Roth); pan. loose erect, awn very short from the bottom of the notch at the end of the lower pale and scarcely extending beyond it, hairs longer than the pales, no rudimentary fl.—*P. 84.*—*Arundo Calamagrostis* L., *E. B. 2159.*—St. slender, 3—4 feet high.—Wet places, rare. P. VII. E.

2. *C. epigejos* (Roth); pan. rather close lobed, straight awn from about the middle of the lower pale, hairs longer than the pales, no rudimentary fl.—*E. B. 403. P. 16.*—St. 3—5 feet high.—Damp shady places. P. VII. E. S. I.

3. *C. stricta* (Nutt.); pan. close, gl. lanceolate rough on the keel one (at least) 3-ribbed, lower pale nearly as long as the upper gl. deeply notched at the top longer than the hairs, straight awn from below the middle of the pale and scarcely extending beyond it, a rudimentary flower, l. of the barren shoots slender.—*E. B. 2160.*—St. erect, 2—3 ft. high. L. broad; on the barren shoots much narrower. Uppermost ligule short, blunt; longer and acute in the Irish plant, which has shorter pan.-branches.—Bogs. Oakmere, Cheshire. Formerly found near Forfar. Lough Neagh. P. VI. VII. E. S. I.

### 20. APE'RA Adans. Wind-grass.

1. *A. Spica-ven'ti* (Beauv.); pan. spreading, anth. linear-oblong.—*Agrostis* Sm., *E. B. 951.* *Anemagrostis* (Trin.) *P. 17.*—St. 1—2 feet high. Pan. very light and elegant; branches spreading horizontally with flowers. Awn 3 or 4 times exceeding the pale. Rudimentary fl. like a pedicel. A tuft of hair on each side of the inner pale.—Sandy fields, rare. A. VI. VII. E. S.

2. *A. interrupta* (Beauv.); pan. close, anth. oval.—*E. B. S. 2951.*—St. 1—2 feet high. Pan.-branches dividing from their base, never spreading. Awn 3 or 4 times exceeding the pale.—Sandy fields. Pampisford and Chippenham, Cambridgeshire. Thetford, Suffolk. Dirleton, S. A. VI. VII. E. S.

### 21. AGROS'TIS Linn. Bent-grass.

1. *A. setacea* (Curt.); panicle close oblong, branches and pedicels rough, gl. unequal acute, lower pale jagged at the top 4-

ribbed, lateral ribs ending in short setæ, kneed and twisted awn from near the base of the pale and twice its length, *l. setaceous involute, sheaths rough*, ligule oblong acute.—*E. B.* 1188. *P.* 83.—Root tufted. *L.* short, almost capillary. *Pan.-branches short.* Midrib of lower glume rough in its upper half, slightly excurrent. Upper pale very minute, a tuft of hairs at its base.—Dry heaths in the South-west. *P.* VII. E.

2. *A. canina* (L.); *pan.* spreading when in flower otherwise close, branches and pedicels rough, gl. unequal acute, lower pale jagged at the top 4-ribbed, kneed and twisted awn from below the middle of and exceeding the pale, lower *l. setaceous tufted*, stem-*l.* narrow flat, *sheaths smooth*, ligule oblong acute.—*E. B.* 1856. *P.* 15.—Trailing leafy shoots. *St.* decumbent below, then erect. *L.* narrow; radical involute. *Pan.-branches* long, slender. *Fl.* green or purplish. Lower gl. not jagged at the top; midrib rough from rather below the middle. Upper pale 0, or very minute. Awn sometimes very short or rarely absent.—Peaty heaths. *P.* VII. VIII. E. S. I.

3. *A. vulgáris* (With.); *pan.* spreading during and after flowering, gl. nearly equal, lower toothed on the upper part of the keel, *ligule short truncate.*—*E. B.* 1671. *P.* 12 & 13.—*St.* long, ascending or decumbent below and rooting at the knots, sometimes with long prostrate stoles. Sheaths mostly smooth. Pedicels toothed. *Fl.* rarely awned.— $\beta$ . *A. pumila* (Lightf.); cæspitose, *st.* 2—3 in. high, *fl.* often awned usually infested with smut.—Rather dry places. *P.* VII. E. S. I.

4. *A. al'ba* (L.); *pan.* compact after flowering, glumes nearly equal, lower toothed throughout its keel, *ligule long, acute.*—*E. B.* 1189. *P.* 13 & 14.—*St.* procumbent and rooting below, then erect, often with long prostrate stoles. Sheaths roughish. *Pan.* spreading with flowers, afterwards close. Pedicels very much toothed. Florets rarely awned.— $\beta$ . *A. stolonifera* (L.); *st.* procumbent and rooting, panicle lobed. *E. B.* 1532.—Fields &c.  $\beta$ . Sea-sands. *P.* VII. *Fiorin-grass.* E. S. I.

## 22. LAGU'RUS Linn. Hare's-tail-grass.

[*L. ovátus* (L.).—*E. B.* 1334. *P.* 88.—*St.* 4—12 in. high. *L.* broad, lanceolate. Spikes ovate, soft, with long protruded awns.—Sandy places in Guernsey. *A.* VI. VII.]

## 23. POLYPO'GON Desf.

1. *P. monspeliensis* (Desf.); awns more than twice as long as the bluntly and shortly lobed glumes.—*E. B.* 1704. *P.* 11.—Root fibrous. *St.* a foot or more high. *Pan.* dense, lobed, pale,

silky, often 2 in. long. Gl. linear, hairy. A most beautiful grass.—Salt marshes, rare.—A. VI. VII. E. S.

2. *P. littoralis* (Sm.); awns as long as the acute glumes.—*E. B.* 1251. *P.* 81. *R.* vii. 75.—Somewhat creeping. St. a foot or more high. Pan. close, lobed, purplish. Gl. linear-lanceolate.—Muddy salt marshes, rare. *P.* VI. VII. E.

#### 24. GASTRID'IUM *Pal. de Beauv.* Nit-grass.

1. *G. lendig'erum* (Gaud.); gl. lanceolate acuminate, lower pale awned, awn rather exceeding the glumes.—*E. B.* 1107. *P.* 86.—St. 3—12 in. high. L. roughish at the edges. Ligule oblong. Pan. close, almost spiked, lobed. *Gl. remarkably ventricose and shining at the base.* Pales very small.—Damp places especially near the sea, rare. A. VI.—IX. E.

### Tribe X. *Aveneæ.*

#### 25. HOL'CUS *Linn.* Soft-grass.

1. *H. lanátus* (L.); upper gl. blunt apiculate, awn smooth except near the tip ultimately curved like a fish-hook and included within the glumes, *sheaths and knots villose.*—*E. B.* 1169. *P.* 21.—Cæspitose. Root fibrous. Height 1—2 feet. Knots not hairy. Inflorescence paniced, often pinkish. Gl. rough. Lower fl. awnless, quite smooth or slightly rough at the point.—Meadows and pastures. *P.* VII. *Yorkshire Fog.* E. S. I.

2. *H. moll'is* (L.); upper gl. acute, awn rough throughout ultimately kneed protruding beyond the glumes, l. rough, sheaths glabrous, *knots bearded.*—*E. B.* 1170. *P.* 21 & 22.—Creeping. Height 1—2 feet. St. and l. subglabrous or slightly hairy. Inflorescence not so compact as in the preceding, whitish. Gl. smooth. Lower fl. awnless; but sometimes it has an awn; rarely the upper is perfect. Occasionally the spikelets are much smaller and the plant only 12—18 in. high.—Thickets or open places on a light soil. *P.* VII. E. S. I.

#### 26. CORYNE'PHORUS *Pal. de Beauv.*

1. *C. canes'cens* (Beauv.); pan. rather dense long, gl. exceeding the fl. acuminate, awn from near the base of the pale, l. setaceous.—*Aira Sm., E. B.* 1190. *P.* 110.—St. tufted, slender. 6—8 in. high. L. many. Pan. close, spreading with flowers;

branches short. Spikelets variegated with purple and white. Anth. dark purple. Lower portion of the awn dark yellow, straight, cylindrical, striated lengthwise and slightly twisted; upper part *clavate*, white tinged with purple.—Sandy coasts of Norf., Suff., and Jersey. P. VI. VII. E.

27. AIRA Linn. Hair-grass.

\* *Lower pale truncate, jagged. Nut free, not furrowed on the back.*

DESCHAMPSIA Beauv.

† Awn straight.

1. *A. cæspitosa* (L.); pan. spreading, l. flat, *gl. slightly rough at the midrib, awn from below the middle of the pale and scarcely extending beyond its tip, ped. of second fl. downy or hairy.*—*E. B.* 1453. *P.* 23.—Root tufted. St. 1—4 feet high. L. rigid, roughish; their margins involute when dry. Pan.-branches rough. Lower pale with 4 veins in addition to that which ends in the rough awn. Rudiment of third fl. often scarcely, if at all, distinguishable; or half the length of the upper fl. and somewhat clavate.— $\beta$ . *brevifolia* (Parn.); radical l. short, sheaths and st. smooth, panicle small. *P.* 106. Viviparous states are often called *A. alpina*.— $\gamma$ . *longiaristata* (Parn.); awns exceeding the fl., sheaths rough. *P.* 105.— $\delta$ . *A. alpina* (L.); pan. close, l. mostly involute, *gl. smooth on midrib, awn from above the middle of the pale.* Height 6—12 in. L. narrower. Fl. often viviparous. *E. B.* 2102. *P.* 23.—Meadows, thickets, &c.  $\beta$ ,  $\gamma$ , and  $\delta$  on mountains. *P.* VII. E. S. I.

†† Awn bent, twisted at the base.

2. *A. flexuosa* (L.); pan. spreading triply forked with wavy branches, l. very narrow subsetaceous, *awn from near the base of the pale and exceeding it, pedicel of the second fl. less than  $\frac{1}{4}$  of its length, ligule short truncate.*—*E. B.* 1519. *P.* 107.—St. erect, slender, about a foot high. L. solid, nearly terete. *Upper sheaths rough from above downwards.* Lower pale notched at the tip.—*A. montana* (Huds. not L.) is a form with more slender and shorter leaves.—Heathy places. *P.* VII.

E. S. I.

3. *A. set'acea* (Huds.); pan. spreading drooping at the end, l. filiform, *awn from near the base of the pale and exceeding it, pedicel of second fl. quite equalling  $\frac{1}{2}$  its length, ligule linear-lanceolate.*—*Sy. E. B.* 1733. *A. uliginosa* Weihe.—St. erect, slender. L. folded. Sheaths smooth.—Wet turfy bogs. *P.* VII. VIII. E. S. I.

\*\* *Lower pale bifid. Nut adnate to the pales, furrowed on the back. No rudiment of a third flower.* AIROPSIS Fries.

4. *A. caryophyll'lea* (L.); pan. spreading triply forked, spikelets rounded below, awn from below the middle of the pale and extending considerably beyond its attenuate deeply bifid tip.—*E. B.* 812. *P.* 24. *Avena* Koch.—St. 6—12 in. high. L. short and narrow. Sheaths roughish from below upwards. Spikelets small, rounded below, chiefly collected at the ends of the branches which are sometimes divaricate with seed.—Dry gravelly places. A. V. E. S. I.

5. *A. præ'cox* (L.); pan. spikelike oblong, spikelets scarcely rounded below, awn from below the middle usually near the base of the pale and extending considerably beyond its attenuate deeply bifid tip.—*E. B.* 1296. *P.* 25. *Avena* Koch.—Height 1—6 in. Pan. close, oval or oblong. L. very narrow. Sheaths smooth.—Dry and sandy places. A. IV. V. E. S. I.

## 28. TRISE'TUM Pers.

1. *T. flaves'cens* (Beauv.); pan. much branched diffuse equal, gl. very unequal about 3-flowered.—*P.* 54. *Avena* L., *E. B.* 952.—St. about a foot high. Radical l. and sheaths hairy. Spikelets yellowish. Upper gl. oblong-lanceolate, acuminate. Floral axis hairy, hairs short.—Fields. P. VII. E. S. I.

## 29. AVE'NA Linn. Oat.

\* *Upper gl. 5—9-veined. Spikelets ultimately drooping. Root annual. No lateral clusters of leaves. L. alike on both sides.*

1. *A. fat'ua* (L.); pan. erect, spikelets of about 3 fl., fl. falling short of the gl. hairy at the base, lower pale bifid at the end.—*E. B.* 2221. *P.* 37.—Height 3 feet. Fl. with long fulvous hairs at their base by which it may be distinguished from *A. sativa*, the cultivated Oat.—Corn-fields. A. VII. *Wild Oat.* E. S. I.

†2. *A. strig'osa* (Schreb.); pan. secund, spikelets of about 2 fl., fl. equalling the gl., lower pale ending in 2 long straight bristles.—*E. B.* 1266. *P.* 26.—Height 3 feet. Very like *A. sativa* but readily distinguished by the bristles at the end of the fl.—Corn-fields. A. VII. E. S.

\*\* *Upper gl. 3-veined. Spikelets erect. Root perennial. Lateral clusters of l. barren. L. with raised ribs.*

3. *A. pratens'is* (L.); pan. erect with simple or slightly divided

branches, *fl. erect* 3—6 exceeding the glumes, *l. rough*.—*E. B.* 1204. *P.* 52.—Root fibrous. Height nearly 2 feet. St. usually nearly round. L. usually short, narrow, acute. Pan.-branches usually simple with only one spikelet.—*β. longifolia* (Parn.); *l.* much longer.—*γ. A. alpina* (Sm.); st. often compressed and sheaths keeled, pan.-branches often with several 5—6-fl. spikelets, upper pale less acute, *l.* broader. I believe that none of these characters are permanent. *E. B.* 2141. *P.* 53.—*A. planiculmis* of *E. B. S.* 2684 may belong to this species; it differs by its greatly compressed st., strongly keeled sheaths and more branched panicle.—In *α* and *β* the lowest fl. sometimes slightly falls short of the longer gl., in *γ* exceeds it.—Dry pastures and mountainous places. *P.* VI. E. S. I.

4. *A. pubescens* (L.); pan. erect nearly simple, fl. erect 2 or 3 scarcely exceeding the glumes, lower *l. and sheaths hairy*.—*E. B.* 1640. *P.* 53.—Creeping slightly. Height 1—2 feet. L. short, rounded behind the tip.—Chalky and limestone districts. *P.* VI. E. S. I.

### 30. ARRHENATHERUM *Pal. de Beauv.* Oat-grass.

1. *A. elatius* (M. & K.); *l.* flat.—*Holcus* Sm., *E. B.* 813. *P.* 25.—Height 2—3 feet. Root fibrous. Knots of the st. glabrous, sometimes downy. Pan. long, ultimately close. Spikelets greenish.—*β. A. bulbosum* (Lindl.); base of the st. with swollen knobs, knots downy. *P.* 26.—Hedges and pastures. *P.* VI. E. S. I.

## Tribe XI. Festuceæ.

### 31. TRIO'DIA *R. Br.* Heath-grass.

1. *T. decumbens* (Beauv.); pan. racemose, spikelets few oval, fl. about 4 scarcely extending beyond the glumes without awns.—*E. B.* 792. *P.* 30.—St. 6—12 in. high. L. flat. Sheaths rather hairy. Ligule a tuft of hairs. Spikelets few, 1—7. Gl. smooth, coriaceous. Lower pale with 3 points, 5-ribbed, hairy at the base.—Dry places and heaths. *P.* VII. E. S. I.

### 32. KÆLERIA *Pers.* Crested Hair-grass.

1. *K. cristata* (Pers.); pan. compact spikelike interrupted below, lower pale acute, *l.* narrow rough at the edges ciliate.—*Aira* L., *E. B.* 648. *P.* 19.—Root crowned with the undivided sheaths of the old leaves. St. 6—18 in. high, downy particularly in the upper part. L., gl. and pales downy or glabrous. Gl. finely toothed on the keel. Lower pale finely toothed on the

midrib. Sometimes the l. become convolute (*K. albescens* DC.P). In dry places the l. fall short of the st., in damper they are long and often nearly equal it.—Dry pastures. A large form on Ben Bulben, Co. Sligo. P. VI. VII. E. S. I.

### 33. MEL'ICA Linn. Melic.

1. *M. uniflora* (Retz); pan. branched slightly drooping, spikelets erect with 1 perfect glabrous fl., gl. equalling pales, l. flat, *ligule* short blunt with a slender acuminate lobe on one side.—*E. B.* 1058. P. 18.—Shady and rocky woods. P. V. VI. E. S. I.

2. *M. nitans* (L.); pan. a nearly simple lax secund raceme, spikelets drooping with 2 perfect glabrous fl., gl. falling short of pales, l. flat, *ligule* short blunt.—*E. B.* 1059. P. 18.—Damp shady woods in hilly districts. P. V. VI. E. S.

### 34. MOLIN'IA Schrank.

1. *M. cærúlea* (Moench); pan. erect long narrow, spikelets 1—3-fl., lower pale 3-veined awnless, upper part of the st. naked.—*E. B.* 750. P. 20.—St. 1—2 feet high, with only one knot placed near to its base. L. long, linear, attenuate, all from near the base of the stem. Panicle purplish, close.— $\beta$ . *M. depauperata* (Lindl.); spikelets 1-fl. few. P. 19.— $\gamma$ . *major*; pan.-branches long, spikelets distant of about 3 fl., st. 3—4 feet high.—Wet heaths.  $\beta$ . Alpine places. P. VII. VIII. E. S. I.

### 35. PO'A Linn. Meadow-grass.

\* *Root fibrous, annual. Base of stem sometimes prostrate and rooting. Pan.-branches solitary or in pairs.*

1. *P. an'nua* (L.); pan. spreading erect with a triangular outline, spikelets ovate-oblong of 5 or 6 free fl., lower pale with 5 veins, upper sheath longer than its leaf, *ligule* oblong acute.—*E. B.* 1141. P. 40, 41.—St. ascending or prostrate. L. flaccid, often wavy, broad. Spikelets subsecund with patent or divaricate branches.—Very common. A. III.—IX. E. S. I.

\*\* *Root fibrous, perennial.*

† Lower pan.-branches solitary or in pairs. Dorsal and marginal veins of the lower pale hairy.

2. *P. bulbósa* (L.); pan. close erect, spikelets ovate of 3 or 4



acute webbed <sup>1</sup> fl., lower pale with 3 silky veins, upper sheath below the middle of the st. much longer than its leaf, ligule prominent acute.—*E. B.* 1071. *P.* 89.—Root fibrous. Base of the st. and offsets swollen, bulblike. L. with a narrow white serrate edge. The st. soon wither, and the tubers lie loose until the autumn.—Sandy seashore of the South and East. *P.* IV. V. E.

3. *P. minor* (Gaud.); pan. oblong subovate, spikelets of 3 or 4 webbed fl., lower pale with 5 veins but only 3 hairy, upper sheath longer than its leaf which is folded and slightly incurved but tapering at the tip, uppermost knot covered, upper ligule long acute, lower ones short rather blunt.—*P. flexuosa* Sm., *E. B.* 1123, not of others. *P. laxa* Fr. ?—Root fibrous. St. 6—8 in. high. Fl. rarely viviparous.—Lofty mountains. Loch-na-Gar. *P.* VII. VIII. S.

4. *P. lax'a* (Haenke!); pan. lax slightly drooping, spikelets oblong-ovate of 3 free fl., lower pale with 3 hairy veins, upper sheath longer than its leaf which is flat and taper-pointed, uppermost knot covered, ligules all long acute.—*P.* 38. *P. stricta* Sy. *E. B.* 1763. The synonymy of this and Sp. 3 is very doubtful.—Root fibrous. St. 6—12 in. high. Fl. often viviparous.—Lofty mountains. Loch-na-Gar. *P.* VII. VIII. S.

5. *P. alpina* (L.); pan. erect spreading when in flower, spikelets ovate of 3 or 4 free fl., lower pale with 3 hairy veins, upper sheath longer than its leaf which is folded and rounded behind the tip, uppermost knot exposed, ligule long pointed.—*E. B.* 1003. *P.* 37 & 94. *P. stricta* Lindb.—Root fibrous, tufted. St. 6—12 in. high, covered with decayed basal sheaths common to it and the tuft of leaves. Fl. often viviparous.—Lofty mountains. *P.* VI. VII. E. S. I.

6. *P. glauca* (Sm.); pan. erect slender, spikelets ovate of 2 or 3 acute free fl., lower pale with 5 veins but only 3 hairy, upper sheath about as long as its leaf which is folded and slightly incurved but tapering at the tip, uppermost knot near to the base of the stem, ligule blunt.—*E. B.* 1720.—Root-stock rather creeping. St. 6—12 in. high. Lowest fl. longer than the large glume.—*P. cæsia* Sm. *E. B.* 1719 is a very doubtful plant—Mountains. Ben Lawers. Ben Nevis. Clova. Snowdon *P.* VII. E. S.

<sup>1</sup> That is, connected together by fine cottony fibres growing from the base of each flower: when these are wanting, the fl. is said to be free.

†† Lower pan.-branches in fives or 2 or 3 together. Dorsal and marginal veins of the lower pale hairy.

7. *P. nemoralis* (L.); pan. rather drooping slender, spikelets ovate-lanceolate of 3 or 4 webbed fl., lower pale with 5 veins but only 3 hairy, *upper sheath not longer than its leaf*, uppermost knot at about the middle of the st. exposed, *ligule extremely short truncate*.—*E. B.* 1265. *P.* 36.—Slightly creeping. St. slender, 1—2 ft. high. Sheaths smooth.—*α*; st. slender weak, pan. lax.—*β. angustifolia* (Parn.); st. and pan. very slender, l. long and narrow, uppermost knot near the pan., spikelets few 1—2-flowered.—*γ. P. coarctata* (Gaud.); st. rigid, pan. close, spikelets 3—5-flowered.—*δ. glaucantha*?; st. slender, pan. with many long-stalked spikelets, plant glaucous.—Shady places. *γ.* On walls. *δ.* Mountains. *P.* VI. VII. E. S. I.

8. *P. Parnelli*'ii (Bab.); pan. suberect large rather close oblong, spikelets ovate of 2 or 3 acute *free fl.*, lower pale with 5 veins but only 3 hairy, *upper sheath usually longer than its leaf*, upper knot at about the middle of the st. exposed, *ligule very short truncate*.—*E. B. S.* 2916. *P.* 36.—St. ascending, 1 ft. or more high, compressed; knots 5 or 6, uppermost not above the middle of the stem. Ligule 6 times as broad as long, but longer than that of *P. nemoralis*, to which this plant is perhaps too nearly allied. Occasionally the fl. are slightly webbed.—High Force and other parts of Upper Teesdale. *P.* VII. E.

9. *P. Balfour*'ii (Parn.); pan. erect rather spreading, spikelets ovate of 3 or 4 webbed fl., lower pale with 5 veins but only 3 hairy, upper sheath about as long as its leaf, upper two-thirds of the stem without knots, *ligule prominent blunt*.—*P.* 66. *E. B. S.* 2918.—Creeping. St. 3—15 in. high; knots about 3, uppermost within the lower third of the stem. Lower fl. as long as the larger glume. *Fl. Dan.* 964? Combined with *P. glauca* by Syme.—*β. P. montana* (Parn.); spikelets few of 2 or 3 free fl., knots 4 or 5 rather above the middle of the stem. *P. dissitiflora* (R. & S.)?—Tops of mountains. *P.* VII. E. S.

††† Lower pan.-branches subverticillate, quinate. Dorsal vein of the lower pale hairy or glabrous; *marginal glabrous*.

10. *P. trivialis* (L.); pan. diffuse, spikelets ovate of 2 or 3 acute webbed fl., lower pale with 5 veins, dorsal vein hairy, upper sheath much longer than its leaf, *ligule acute long*.—*E. B.* 1072. *P.* 35.—Root tufted. St. 1—2 feet high. Sheaths usually slightly rough.—*β. parviflora* (Parn.); spikelets small 1—2-flowered, plant slender.—Moist and shady places. *P.* VI. E. S. I.

[\**P. sudetica* (Haenke); pan. diffuse, spikelets oval of 3 or rarely 5 acute *not webbed fl.*, lower pale with 5 glabrous veins, upper sheath very much longer than its leaf, *ligule blunt very short*.—*R. i.* 90.—Rhizomatous, not creeping. *St.* 2-edged, 2—3 feet high. Sheaths, edges, and midribs of l. rough. L. hooded and apiculate.—In deep shade in several places near Kelso. *P.*]

\*\*\* *Creeping by long soboles.*

11. *P. pratensis* (L.); pan. diffuse, spikelets ovate of 3 or 4 webbed fl., lower pale with 5 prominent veins but only 3 hairy, upper sheath much longer than its leaf, *ligule prominent blunt*.—*E. B.* 1073. *P.* 31—34.—Very variable in size. *St.* and sheaths nearly always smooth. *Fl.* strongly webbed.— $\beta$ . *P. subcærulea* (Sm.); spikelets broader, l. broad and short, upper l. compressed rounded at the end behind. *E. B.* 1004.— $\gamma$ . *angustifolia* (Parn.); spikelets small, l. slender long, lower l. involute.—Common. *P.* VI. VII. E. S. I.

12. *P. compressa* (L.); pan. erect or slightly unilateral spreading when in flower otherwise close, spikelets ovate or oblong-ovate of 5—7 blunt slightly webbed fl., lower pale 3-veined, veins hairy, upper sheath about as long as its leaf, *uppermost knot at about the middle of the stem, ligule short truncate*.—*E. B.* 365. *P.* 37 & 90.—*St.* decumbent at the base, then erect, very much compressed, 1—1½ ft. high.— $\beta$ . *P. polynoda* (Parn.); fl. free, pales with 2 faint interm. veins, uppermost knot higher, *ligule rather more prominent*. *P.* 91—93.—Dry situations. *P.* VII. E. S. I.

### 36. GLYCERIA *R. Br.*

1. *G. aquatica* (Sm.); pan. erect much branched spreading, branches rough, spikelets oblong of 5—10 fl., lower pale blunt, l. smooth with terete sheaths.—*E. B.* 1315. *P.* 44. *G. spectabilis* Koch.—Creeping. *St.* 3—6 ft. high, smooth slightly compressed. Sheaths very long. L. long, rough on the edges and keel, never floating. *Ligule short*. Pan. large; branches angular, slender, branched.—Watery places. *P.* VII. E. S. I.

2. *G. fluitans* (R. Br.); pan. secund slightly branched very long, branches nearly simple roughish, spikelets linear of 7—12 adpressed lanceolate-oblong acute fl., lower pale nearly thrice as long as broad, sheaths compressed.—*E. B.* 2975. *P.* 95.—*St.* ascending, rooting below, or floating. Sheaths nearly smooth, striate. L. pale green, acute, often floating. *Ligule*

long. Pan. very long, often nearly simple; branches without callosities, ascending, lowermost usually in pairs. Spikelets adpressed. Lower pales rather the shorter, with a triangular central point. *Anthers about 5 times as long as broad*, purple, pale yellow when empty.—β. *G. pedicellata* (Townsend); pan.-branches simple roughish, spikelets of 13 blunt flowers. L. more acute. Lowermost pan.-branches about in threes. *Anth.* about 4 times as long as broad, yellow when young. *A. N. H.* ser. 2, v. 105. *Curt. Fl. Lond.* i. 18.—Watery places. P. VI.—IX. *Flote-grass.* E. S. I.

3. *G. plicata* (Fries); pan. compound, branches compound nearly smooth erect with flowers *divaricate with fruit*, spikelets linear of 7—20 oval-oblong rather acute fl., *lower pale twice as long as broad*, sheaths compressed.—*R.* vii. 79. *G. fluitans* Sm., *E. B.* 1520. P. 45.—St. ascending, rooting below. Sheaths rough, furrowed. L. glaucous, bluntish, plicate when young. Ligule shorter. Pan. much branched; branches with callosities at the base, lowermost about in fives. Lower pales with 3 teeth at the end. *Anth.* about 3 times as long as broad, cream-coloured, fuscous when empty.—Stagnant water and wet places. P. VI.—VIII. E. S. ? I.

### 37. SCLEEROCH'LOA *Pal. de Beauv.*<sup>1</sup>

\* *Glumes with 3 veins.*

† *Panicle unilateral.*

1. *S. maritima* (Lindl.); pan. branched, lowermost branches in pairs or single, branches ultimately erect, spikelets linear adpressed 4—8-flowered, rachis terete, *lower pale blunt apiculate, midrib reaching the tip, stoloniferous.*—*Glyceria* Sm., *E. B.* 1140. P. 42.—Root fibrous, with ascending prostrate or rarely rooting leafy stoles. L. involute; the central ridge on their upper surface strongly marked the others faint. *Anth.* about 6 times as long as broad. Lower pale with involute edges. Ligule bluntish.—β. *hispidula* (Parn.); st. compressed, rachis furrowed on one side and as well as the pan.-branches rough. P. 99.—Sea-coast, in damp places. P. VI. VII. E. S. I.

2. *S. Borreri* (Bab.); pan. branched, branches ultimately erect-patent lowermost generally in fours, spikelets linear 4—7-fl., rachis terete, *lower pale with a rigid apiculus formed by the tip of the dorsal vein, cæspitose.* *Glyceria*, Bab., *E. B. S.* 2797 (1837). P. 98. *G. conferta* Fries (1839).—St. 6—12 in. high.

<sup>1</sup> See *Crepin, Notes Fl. Belg.* v. 155—214.

No stoles. L. short, flat, with very long sheaths. Ligule short, truncate. Edges of the lower pale not involute. Spikelets and fl. half the size of those of *S. maritima* and *S. procumbens*. Pan.-branches short, scarcely elongated after flowering, hispid.—Muddy salt marshes. P. VI.—VIII. E. I.

3. *S. procumbens* (Beauv.); pan. ovate-lanceolate compact distichous rigid, spikelets linear-lanceolate of about 4 fl., rachis angular, lower pale blunt with an apiculus formed by the tip of the dorsal vein, root fibrous.—*Glyceria* Sm., *E. B.* 532. P. 42.—St. procumbent (rarely erect when growing in water), rigid. L. flat, with inflated sheaths. Pan. about 2 in. long, with very short rigid branches spreading in 2 rows. Fl. large.—Muddy sea-shores. A. VI. VII. E. S. I.

†† Panicle regular.

4. *S. distans* (Bab.); pan. branched, branches long ultimately spreading or deflexed lowermost in fours or fives, spikelets linear 3—6-flowered, rachis semiterete rather flat on one side, lower pale blunt, midrib not reaching to the tip, root fibrous.—*Glyceria* Sm., *E. B.* 986. P. 41.—Without rooting stoles. St. decumbent below. L. flat, short, with 8—10 equally prominent ridges upon their upper surface. Ligule short and truncate. Edges of lower pale not involute. Spikelets and fl. half the size of those of the preceding.—β. *obtusa* (Parn.); pan. more compound, spikelets about 7-flowered, lower pale truncate and broader, ligule shorter. P. 96 & 97.—Sea-shores and waste sandy places. β. Leicestershire. P. VI.—VIII. E. S. I.

\*\* Glumes 1- (rarely 3-) veined.

5. *S. rigida* (Link); pan. lanceolate rigid distichous, spikelets linear acute of 7—10 fl., lower pale blunt with a mucro, upper gl. reaching to the base of the third fl., root fibrous.—*Glyceria* Sm., *E. B.* 1371. P. 43.—St. slender, wiry, erect. L. nearly flat, acute. Pan. 1—2 in. long, nearly simple. Lower pale faintly veined. Fl. small.—Dry places. A. VI. E. S. I.

6. *S. loliacea* (Woods); pan. racemose narrow rigid secund, spikelets oblong of 8—12 fl., lower pale blunt with a mucro, upper gl. reaching to the base of the fourth fl., root fibrous.—*Triticum* Sm., *E. B.* 221. P. 43.—St. stout, slightly curved, ascending, 2—6 in. long. L. flat, convolute when dry. Spikelets usually solitary, alternate, all directed to one side; footstalks very short and stout. Lower pale with well-marked marginal veins.—Sandy sea-coasts. A. VI. VII. E. S. I.

38. *BRI'ZA* Linn. Quaking-grass.

1. *B. minor* (L.); spikelets triangular of about 7 fl., *gl. exceeding the lowest fl.*, pan. diffuse, *ligule long lanceolate acute*.—*E. B.* 1316. *P.* 101.—St. very slender, about 1 foot high. Spikelets pale green. Lower pale roundish cordate, cartilaginous, very gibbous in the middle of the back.—Dry and sandy fields in the South-west. A. VII. E.

2. *B. média* (L.); spikelets broadly ovate of about 5 fl., *gl. falling short of the lowest fl.*, pan. diffuse, *ligule truncate very short*.—*E. B.* 340. *P.* 30.—St. slender erect, 1—1½ foot high. Panicle light and elegant, with slender branches. Spikelets usually purplish. L. linear-acuminate. Lower pale roundish oval, cartilaginous, not gibbous.—Pastures. P. VI. E. S. I.

39. *CATABROSA* Pal. de Beauv.

1. *C. aquatica* (Beauv.); pan. long-pyramidal with half-whorls of patent branches, lower pale 3-ribbed, l. broad linear blunt.—*E. B.* 1557. *P.* 20.—Creeping. St. long, procumbent or floating below. L. flat. Pan.-branches in alternate threes or fives. Spikelets usually 2- (or 3—5-) flowered. Gl. very thin, often purplish. Fl. distant.—*β. minor*; st. 2—3 in. high, spikelets mostly 1-flowered.—Ponds and ditches. *β.* Wet sea-sands. P. VI. VII. E. S. I.

40. *CYNOSURUS* Linn. Dog's-tail-grass.

1. *C. cristatus* (L.); raceme spikelike linear, fl. with a very short awn.—*E. B.* 316. *P.* 28.—St. 12—18 in. high. Spike unilateral, plane-compressed. Spikelets closely placed. Bract comblike.—Pastures. P. VIII. E. S. I.

[*C. echinatus* (L.); raceme contracted close ovate, awns about as long as the pales.—*E. B.* 1333. *P.* 28 & 129.—St. erect, 1—2 feet high. Bract comblike with long points.—Sandy places in Guernsey and Jersey. A. VII.]

41. *DAC'TYLIS* Linn. Cock's-foot-grass.

1. *D. glomerata* (L.); pan.-branches with ovate clusters of spikelets, st. erect, linear flat with rough margins, root cæspitose.—*E. B.* 335. *P.* 29.—A coarse grass. Pan.-branches long, spreading or divaricate with fl., afterwards adpressed, distant; each bearing an ovate cluster of spikelets; or panicle reduced to one cluster.—Meadows. P. VI. VII. E. S. I.

## 42. FESTUCA Linn. Fescue-grass.

\* *Root-leaves very narrow. Ligule with round auricles. Awn terminal.*

† Awn longer than the pale. Gl. very unequal. VULPIA.

‡ Usually triandrous.

1. *F. uniglumis* (Sol.); uppermost sheath far distant from the erect close 2-ranked *simple panicle*, fl. compressed keeled, *gl. very unequal often only one*, larger gl. very long and very acute.—*E. B.* 1430. *P.* 112.—St. 6—12 in. high, erect, leafy nearly to the top. Panicle close, short. Lower gl. usually scarcely distinguishable. Upper gl. setigerous nearly equalling 1st fl. on same side.—Sandy sea-shores. A. VI. E. I.

‡‡ Usually monandrous.

2. *F. sciuroides* (Roth); uppermost sheath far distant from the *erect-patent oblong pan.*, lowermost pan.-branch about equaling  $\frac{1}{2}$  pan., fl. terete rough, *gl. unequal as 2 to 1*, larger gl. about equalling lowest flower.—*F. bromoides* Sm., *E. B.* 1411.—Slender, 6—12 in. high. L. linear, involute.—Walls and sandy places. A. ? VI. VII. E. S. I.

3. *F. ambigua* (Le Gall); uppermost sheath very nearly reaching to the *long narrow erect close pan.*, lowest pan.-branch equaling more than  $\frac{1}{3}$  of pan., fl. terete rough, *gl. unequal as 3—6 to 1*, larger gl. about equalling  $\frac{1}{3}$  of lowest flower.—*E. B. S.* 2970.—Slender, 8—12 in. high. L. involute.—Sandy places. Isle of Wight. Dorset. Suffolk. A. V. VI. E.

4. *F. Myurus* (L.); uppermost sheath reaching to or partly covering the *long narrow nodding interrupted pan.*, lowest pan.-branch equaling  $\frac{1}{4}$  of pan., fl. terete rough, *gl. unequal as 3 to 1*, larger gl. about equalling  $\frac{1}{2}$  lowest flower.—*E. B.* 1412. *F. pseudo-myurus* (Soy.-Will.).—About a foot high, slender. Pan. very long and narrow.—Walls and sandy places, rare. A. ? VI. VII. E. S. ? I.

‡‡ Triandrous. Awn shorter than the pale.

5. *F. ovina* (L.); pan. narrow subsecund close with fr., spikelets 4—6-fl., fl. mostly awned, *l. all setaceous*, sheaths glabrous, *cæspitose*.—*E. B.* 585. *P.* 56, 57.—Very variable. L. short, slightly curved, densely tufted. Fl. glabrous or hairy. Spikelets sometimes changed into leafy shoots.— $\beta$ . *F. tenuifolia* (Sibth.); l. very long setaceous, fl. awnless.— $\gamma$ . *F. duriuscula* (L.); pan.

pyramidal, branches spreading, l. filiform channelled, st.-l. broader. St. and l. stouter. Fries thought it distinct.—Dry hilly pastures.  $\beta$ . On mountains.  $\gamma$ . Damper places. P. VI. *Sheep's Fescue-grass*. E. S. I.

6. *F. rubra* (L.); pan. broadish below subsecund, spikelets 4—10-fl., l. involute-setaceous, st.-l. flat, lowest sheaths hairy, *soboliferous*.—*F. duriuscula* Sm. (in part), *E. B.* 470, *P.* 58—60.—Very variable. Fl. shortly awned, glabrous, hairy or villose. L. variable in length and breadth and the fl. in size. Creeping but *cæspitose*.—Common in dry sandy, rarely in wet places. P. VI. E. S. I.

7. *F. oraria* (Dum.)<sup>1</sup>; pan. secund, spikelets 4—10-fl., l. all involute-setaceous, lowest sheaths hairy, far creeping not *cæspitose*. *F. rubra* Sm., *E. B.* 2058.—Near *F. rubra*, but mode of growth very different.—Sandy sea-shores. P. VI. E. S.

\*\* *Root-leaves broad and flat. Ligule not auricled. Awn 0, or dorsal.* SCHEDONORUS Beauv.

† Uppermost ligule prominent. Lower pale 3-veined.

8. *F. sylvatica* (Vill.); pan. erect diffuse much-branched, branches rough, spikelets of 3—4 *awnless acute fl.*, lower pale rough, *dorsal rib serrulate throughout*, l. lanceolate-linear with rough margins.—*Poa P.* 44 & 100. *F. Calamaria* Sm., *E. B.* 1005.—Scarcely creeping. St. 2—4 feet high, covered at the base with imbricate broad acute leafless sheaths, tufted. L. very long, broad, roughish on both sides; uppermost l. smaller. Lower pale very acute; midrib extending nearly to the tip or slightly beyond it. Ovary pilose at the top.— $\beta$ . *F. decidua* (Sm); l. narrower, fl. about 2. *E. B.* 2266.—Woods in mountainous districts. P. VII. E. S. I.

†† Uppermost ligule very short. Lower pale 5-veined. BUCETUM Parn.

9. *F. gigantæa* (Vill.); pan. open drooping branched, spikelets of about 5 *awned fl.*, *dorsal rib of lower pale nearly smooth* not extending to the tip but *ending in a rough awn twice as long as the pale*, l. linear-lanceolate.—*E. B.* 1820. *P.* 47.—St. 3—4 feet high. L. very long, broad, roughish on both sides, except near the base on the underside. Ligule unequal, auricled. Lower pale roughish, membranous, often bifid at the tip. Top of the

<sup>1</sup> *F. oraria* Dum. Agr. Belg. 105 (1823), *F. sabulicola* Duf. (1825), *F. arenaria* Godr. (1855, not Osb.). See *Bull. Belg.* vii. 367.



ovary glabrous.— $\beta$ . *F. triflora* (Sm.); pan. smaller and more erect, spikelets scattered of about 3 flowers. *E. B.* 1918.—Moist woods and thickets. P. VII. E. S. I.

10. *F. arundinacea* (Schreb.); panicle diffuse patent, branches mostly in pairs each bearing 2 or more ovate-oblong spikelets divaricate with fl. or afterwards, spikelets very many of 5—6 closely placed fl., dorsal rib of lower pale ending at or just below the tip or forming a short awn, l. linear-lanceolate.—*F. elatior* Sm., *E. B.* 1593. P. 46, 47.—St. 2—6 ft. high, forming large tufts. L. broad.— $\alpha$ . *F. arundinacea* (Schreb.); pan.-branches divaricate with fl. and fruit. A very large plant, 3—6 ft. high.— $\beta$ . *F. elatior* (L. ?); pan.-branches shorter “divaricate with fl. afterwards ascending.”— $\alpha$ . Banks near the sea.  $\beta$ . Damp pastures. P. VI. VII. E. S. I.

11. *F. pratensis* (Huds.); pan. close subsecund, branches in pairs one bearing a single spikelet the other several never divaricate, spikelets linear-oblong of 5—10 rather distant fl., dorsal rib of lower pale ending at or just below the tip or forming a very short awn, l. linear-lanceolate.—*E. B.* 1592. P. 46. *F. elatior* Koch.—A smaller plant than the preceding. Rachis triangular. Pan.-branches ascending; one of each pair nearly always reduced to a single spikelet. In this and the preceding the pale is blunt or acute according as the midrib is or is not attached up to the tip.— $\beta$ . *F. loliacea* (Huds.); spikes solitary alternate long slender truly distichous, lower ones stalked, rarely in pairs, upper nearly sessile, fl. distant, lower gl. 5—8-ribbed, veins of lower pale parallel, no awn, rachis flattish.—*E. B.* 1821. P. 45, 113 & 114.—Wet meadows. P. VI. VII. E. S. I.

#### 43. BRO'MUS Linn.<sup>1</sup>

\* Spikelets broader upwards when in flower, not afterwards.  
Rib of upper pale finely fringed.

1. *B. erectus* (Huds.); pan. erect nearly simple, spikelets linear-lanceolate, fl. remote subcylindrical, lower pale indistinctly 7-veined, lowest fl.  $\frac{1}{2}$  exceeding the upper gl. and longer than its awn, root-l. very narrow ciliate.—*E. B.* 471. P. 51.—St. 2—3 ft. high, erect. Root-l. convolute; upper l. broadest; sheaths somewhat hairy with upward hairs.— $\beta$ . *villosus*; lower pale hairy.—On dry sandy and chalky soil. P. VI. VII. E. S. I.

2. *B. asper* (Murr.); pan. drooping with long slightly divided branches, spikelets lanceolate, fl. remote linear-lanceolate, lower pale more or less hairy 5—7-ribbed, lowest fl. twice exceeding the upper gl. and longer than its awn, l. broad hairy.—*E. B.*

<sup>1</sup> Synonymous with the genus *Schedonorus* of Fries, not of Beauv.

1172. *P.* 51. *B. serotinus* (Benek.).—St. 4—5 feet high. L. flat; lower ones broadest; sheaths with downward hairs.—The *B. Benekenii* (Syme) appears to have a glabrous upper sheath, much more hairy pales, more nearly glabrous glumes. I have not seen any native specimen according with it.—Damp woods and thickets. *P.*? VII. E. S. I.

\*\* *Spikelets always broader upwards. Ribs of upper pale strongly fringed. Awn long.*

3. *B. sterilis* (L.); pan. drooping, branches long slightly divided, spikelets linear-lanceolate, fl. remote, lower pale glabrous shorter than its awn with 7 distinct equidistant ribs, l. pubescent.—*E. B.* 1030. *P.* 50.—Height 1—2 feet. L. broad, flat.—Waste places. A. VI. E. S. I.

4. *B. madritensis* (L.); panicle erect, branches short scarcely divided, spikelets lanceolate, fl. linear remote subcylindrical, lower pale about as long as its awn 7-ribbed, 2 lateral ribs close together, interm. rib faint.—*E. B.* 1006. *P.* 50.—*a. B. diandrus* (Curt.); st. glabrous, rachis and pedicels rough. St. 6—12 in. high. Remarkable for its erect panicle. Upper pale but little shorter than the lower.—[*β. B. rigidus* (Roth); pan. compact, pedicels very short, upper part of st., pedicels, rachis and gl. pubescent.]—Dry sandy places, rare. [*β.* Channel Islands.] A. VI. VII. E. S. I.

[*B. tectorum* (L.); like *B. madritensis*, pan. secund drooping, upper pale much the shorter.—Introduced.]

[*B. maximus* (Desf.); pan. erect lax at length nodding, branches slightly divided lengthened after flowering, spikelets downy, lower pale 7-ribbed about half as long as its awn.—*E. B. S.* 2820.—Height 1—2 feet. A most beautiful grass.—Sandy places. Channel Islands. A. VIII.]

#### 44. SERRAFAL'CUS *Parlatore*<sup>1</sup>.

\* *Fl. at first loosely imbricate, afterwards distinct and cylindrical.*

\*1. *S. secalinus* (Bab.); pan. loose drooping in fr. slightly compound, simple peduncles about as long as the oblong glabrous spikelets, fl. about as long as the straight awn, lower pale not overlapping the next fl. uniformly rounded at the sides, l. hairy.—*E. B.* 1171. *P.* 49, 121 & 122.—With seed the fl.

<sup>1</sup> Corresponds to the genus *Bromus* of Fries; to part of *Bromus* of Sm., Hook., &c.

spread and the spikelets droop. *Top of upper gl. ½-way between its base and the top of fourth fl.* (second on the same side). Lower pale not twice as long as broad, longer than the upper, 7-ribbed.—*β. Bromus velutinus* (Sm.); pan. nearly simple, fl. larger downy. *P.* 123.—Corn-fields. *A.* VI. VII. E. S.

\*\* *Fl. closely imbricate even with fruit.*

2. *S. racemósus* (Parl.); pan. long erect usually simple, spikelets ovate rather compressed glossy, fl. imbricate about as long as the straight awn, *lower pale uniformly rounded at the sides*, l. and sheaths slightly hairy.—*E. B.* 1079.—*Top of the upper glume ½-way to the top of the fourth flower.* Lower pale longer than the upper. Anth. 4 times as long as broad. Pan. close with fruit.—Common. *B.* VI. E. I.

3. *S. commutátus* (Bab.); pan. loose slightly drooping compound, simple peduncles as long as or longer than the oblong-lanceolate spikelets, fl. loose imbricate about as long as the straight awn, *sides of lower pale bluntly angular* above the middle, l. and sheaths hairy.—*P.* 124. *E. B.* 920.—Lower pale only slightly overlapping the next fl. at the base when in fruit. *Top of upper gl. ½-way to top of fourth fl.* Lower pale twice as long as broad, longer than the upper, glabrous or downy, 7-ribbed. Anth. 6 times as long as broad.—*β. multiflorus* (Parn.); more numerous fl., top of upper gl. rather higher. *P.* 125.—Common. *B.* VI. VII. E. S. I.

4. *S. mollis* (Parl.); pan. close erect compound or rarely simple, spikelets ovate rather compressed pubescent, fl. closely imbricate about as long as the straight awn, *sides of lower pale bluntly angular about the middle*, l. and sheaths hairy or downy.—*E. B.* 1078. *P.* 116.—*Top of the upper gl. ½-way to the top of the sixth flower*; or a little higher (*ovalis* Parn. 117, with short oval spikelets); or about ½-way to the top of the eighth (*pratensis* Parn. 118, with longer spikelets). Lower pale longer than the upper. Simple ped. not longer than the spikelets. Anth. about thrice as long as broad. Rarely the spicules are glabrous, when it is *B. racemosus* Parn. 119.—A maritime plant with nearly or quite prostrate st., nearly simple pan., and nearly glabrous pales, seems to be the *B. hordeaceus* (Fries). Another with very short ped. and densely downy spikelets is also found near the sea.—Common. *A.?* V. VI. *Lop-grass.* E. S. I.

\*5. *S. arven'sis* (Godr.); pan. spreading compound its branches ultimately horizontal, spikelets linear-lanceolate, *pales equal in length lower 7-ribbed* with two prominent ribs near each margin

and its sides bluntly angular above the middle, anth. 4 times as long as broad.—*E. B.* 1984. *P.* 126.—Top of the upper gl.  $\frac{1}{2}$ -way to the top of the fourth flower.—Naturalized in various places from Fife southwards. *A.* VII. VIII. E. S.

[*S. pat'ulus* (Parl.); pan. spreading compound its branches ultimately deflexed, spikelets lanceolate, pales unequal, lower 7-ribbed with two lateral prominent ribs and its sides bluntly angular above the middle, anth. twice as long as broad.—*P.* 127.—Not naturalized. *A.* VI.] E.

[*S. squarrosus* (Parl.); pan. drooping simple, spikelets ovate-lanceolate subcompressed, fl. nearly glabrous imbricate compressed, lower pale 9-ribbed and its sides bluntly angular above the middle, awn twisted divaricate, l. pubescent.—*E. B.* 1885. *P.* 128.—Not naturalized. *A.* VI. VII.] E.

## Tribe XII. *Hordeieæ*.

### 45. BRACHYPO'DIUM *Pal. de Beauv.*

1. *B. sylvat'icum* (R. & S.); spike drooping, spikelets (at first) terete alternate distichous, awns of the upper fl. longer than their pales, l. flat linear-lanceolate flaccid, root fibrous.—*E. B.* 729. *P.* 61.—St. usually solitary or 2 or 3 from the same root, erect, 1—2 feet high. Sheaths hairy. Ligule short, blunt, notched or torn. L. ciliate. Pales usually hairy.—Woods and hedges. *P.* VII. E. S. I.

2. *B. pinnatum* (Beauv.); spike erect, spikelets (at first) terete alternate distichous, awns of the upper fl. shorter than their pales, l. flat linear-lanceolate rigid, creeping.—*E. B.* 730. *P.* 132—137.—St. several, erect, 1—2 feet high. Pales rough or hairy. Sheaths subglabrous. Ligule short, truncate. L. not ciliate.—Sometimes the l. are very narrow and involute, st. very many, spikelets small smooth.—On dry limestone soil. *P.* VII. E.

### 46. TRIT'ICUM *Linn.* Wheat-grass.

1. *T. caninum* (Huds.); spike rather close, spikelets 2—5-fl., 3—5-ribbed gl. and lower pales awned, axis and edges of the rachis hispid, l. flat rough on both sides, root fibrous.—*E. B.* 1372. *P.* 62.—St. erect. Ribs on the upperside of l. very slender. Gl. round on the back, its ribs reaching the tip and joining to form the short awn. Lower pale shorter than its awn; or in an alpine form longer than it.—Banks, rare. *P.* VII. E. S. I.

2. *T. répens* (L.); spike rather close, gl. 5—7-ribbed equalling at least  $\frac{2}{3}$  of the 4—5-fl. spikelet rough on the keel, lower pale acuminate, axis asperous, rachis with rough angles not brittle, l. mostly flat the many slender ribs each bearing a row of deciduous hairs above, soboliferous.—*E. B.* 909. *P.* 62.—L. at first involute afterwards flat, ribs on upperside not much raised nor nearly hiding the interm. surface of the leaf. Rachis glabrous or downy with forward prickles on the angles. Gl. scarcely keeled, acuminate-subulate; ribs reaching the tip. Pales rarely awned.—Common. *P.* VI. *Couch-grass.* E. S. I.

3. *T. pun'gens* (Pers.); spike close, gl. with 7—9 thick ribs not exceeding  $\frac{1}{2}$  the 5—12-fl. spikelet rough on the keel, lower pale acute, axis asperous, rachis nearly or quite smooth not brittle, l. with involute edges the many thick closely-placed ribs slightly rough and each bearing a row of acute points above, upper part of l. wholly involute (subulate and rigid), soboliferous.—*Sy. E. B.* 1811.—*St. erect*, like a corn-field. Ribs on upperside of l. so broad and so elevated as nearly to hide the interm. part of the leaf. Gl. keeled; ribs reaching the tip. Lower pale of our plant often awned. Producing erect barren leafy clustered shoots.—Sea-shores. *P.* VII. E. I.

4. *T. acútum* (DC.); spike rather lax, gl. with 5—7 slender elevated ribs blunt or apiculate not exceeding  $\frac{2}{3}$  of the 5—8-fl. spikelet, lower pale blunt mucronate, axis downy, rachis smooth or slightly rough at the angles not brittle, l. flat or with involute edges the many thick closely placed ribs rough with minute sharp scattered points (asperous) above, soboliferous.—*Sy. E. B.* 1812. *T. laxum* Fr.—*St.* prostrate or ascending. Ribs of the l., on each of which there is usually a deciduous row of hairs, not so completely hiding the intervening hollows as in Sp. 3. Gl. keeled; keel often with forward bristles, reaching the tip or forming a slight mucro. Lower pale rarely shortly awned. Producing decumbent and ascending, barren, leafy, clustered shoots.—Sandy sea-shores. *P.* VII. VIII. E. S. I.

5. *T. jun'ceum* (L.); spike rather loose, gl. with 9—11 slender scarcely elevated ribs blunt equalling at least  $\frac{2}{3}$  of the 4—8-fl. spikelet smooth on the keel, lower pale blunt rarely mucronate, axis smooth or slightly downy, rachis brittle smooth, l. involute with many thick ribs with much spreading hair above, soboliferous.—*E. B.* 814. *P.* 63.—*St.* prostrate. The short hairs on the ribs of the l. spread so as to cover the interm. spaces. Rachis easily separating above each spikelet. Gl. rounded or truncate at the tip; ribs not reaching the tip. Producing decumbent barren leafy shoots.—Sandy sea-shores. *P.* VII. VIII. E. S. I.

47. *EL'YMUS* Linn.

1. *E. arenarius* (L.); spike upright close, rachis flat not winged, gl. lanceolate downy not longer than the spikelets.—*E. B.* 1672. *P.* 64.—Closely resembling *Psamma arenaria*, but readily distinguished by its broad l. and short ligule. Soboliferous. St. 3—4 feet high.—Sandy sea-shores. *P.* VII. E. S. I.

[*E. geniculatus* (Curt.); spike lax, rachis winged, gl. awl-shaped glabrous longer than the spikelet.—*E. B.* 1586. *P.* 131.—St. 3—4 feet high. Spike 1—2 feet long, usually remarkably bent downwards at the second or third spikelet.—In a salt marsh near Gravesend. *Mr. Dickson!* *P.* VII.] E.

48. *HOR'DEUM* Linn. Barley.

1. *H. sylvaticum* (Huds.); gl. all awlshaped not ciliate rough, lateral fl. perfect, middle fl. often barren, lower pale with an awn of twice its length.—*P.* 130. *Elymus europæus* Linn., Sm., *E. B.* 1317.—About 2 ft. high. Spike subcylindrical. Middle fl., if barren, with shorter gl. having involute edges, thus appearing setaceous. The spikelets have a second fl. occasionally.—Woods and thickets on a calcareous soil, rare. *P.* VII. VIII. E.

2. *H. pratense* (Huds.); gl. all setaceous not ciliate rough, lateral fl. imperfect, lower pale of fertile middle fl. with an awn of about its length.—*E. B.* 409. *P.* 11.—Often 2 ft. high. Spike compressed, erect. Gl. of lateral fl. shorter.—Damp meadows. *P.* VII. E. S. I.

3. *H. murinum* (L.); gl. of the middle spikelet linear-lanceolate ciliate of the lateral ones setaceous rough, lateral fl. imperfect.—*E. B.* 1971. *P.* 10.—Spike often slightly nodding, compressed. Height 12—18 in. Awn longer than the lower pale. Lateral gl. sometimes slightly ciliate.—*β. arenarium* (Bab.); lower part of the st. buried, lengthened, branched and rooting, thus appearing to creep.—Waste places. *β.* Loose sand. *B.* VI. VII. E. S. I.

4. *H. maritimum* (Wither.); gl. rough, inner gl. of the lateral fl. half-ovate the rest setaceous, lateral fl. imperfect.—*E. B.* 1205. *P.* 10.—Spike thick, erect, subterete.—Pastures and banks near the sea. *A.* VI. E. S.

49. *LEPTU'RUS* R. Br.

1. *L. filiformis* (Trin.); spike cylindrical-subulate, gl. 2 equalling or slightly exceeding the flowers.—*Rottbœllia* Sm., *E. B.* 760. *P.* 2.—St. 2—6 in. long. Spike long, slender

straight or slightly curved. [*L. incurvatus* (Trin.) is apparently only found as a ballast plant.]—Gravelly and waste places near the sea. A. VII. E. S. I.

50. LO'LIUM Linn. Rye-grass.

1. *L. peren'ne* (L.); with leafy barren shoots, *edges of young l. simply folded*, spikelets 3—11-flowered, gl. equalling the lowest fl., lower pale usually awnless.—*E. B.* 315. *P.* 65.—*St.* 12—18 in. high, usually bent at the lower knots. Whole plant rather dark green.— $\beta$ . *L. aristatum*; lower pale with a long awn.— $\gamma$ . *L. tenue* (L.); spikelets few-flowered, l. slender.—Sometimes the spikelets become converted into branches; or the rachis is so much shortened as to produce a broad ovate close distichous spike.—Common. *P.* VI. *Rye-grass.* E. S. I.

[*L. ital'icum* (A. Braun); with leafy barren shoots, *edges of young l. involute*, spikelets 9—14-flowered, lower pale with a long awn.—*R.* vii. 77. *P.* 138—141.—*St.* many, straight, in close tufts,  $1\frac{1}{2}$ —3 ft. high. Whole plant, especially the spikelets, paler than in the preceding. Ligule short, abrupt.—*L. multiflorum* (Lam.), perhaps confounded with this, has no barren shoots and is annual.—Cultivated fields. *P.* VI. *Italian Rye-grass.*] E.

[*L. linic'ola* (Sond.); no barren shoots, spikelets 7—11-flowered exceeding the gl., lower pale longer than its awn or awnless, fl. tumid with fruit.—*E. B.* 2955.—*St.* erect. Spike slender. Lower pale cartilaginous below, narrower than the upper, tumid in fruit.—Cultivated fields. A. VI. VII.] E.

‡2. *L. temulen'tum* (L.); no barren shoots, *spikelets about 6-flowered equalling or shorter than the gl.*, lower pale awned, fl. tumid with fruit.—*E. B.* 1124. *P.* 142.—*St.* erect. Ligule short. Upper gl. usually present, often bifid.— $\alpha$ ; awns as long as or longer than the pale.— $\beta$ . *L. arvense* (With.); fl. 4—5 without or with short awns. *E. B.* 1125.—Cultivated fields. A. VI.—VIII. *Darnel.* E. S. I.

## FLOWERLESS PLANTS.

## Class III. CRYPTOGAMEÆ.

Substance of the plant of cellular tissue or with a few ducts. No woody fibre. No true flower with stamens and pistils. No distinct embryo, nor cotyledons.

A. Plants with a few ducts amongst the cellular tissue. Producing spores which develop into a prothallus which bears antheridia and archegonia<sup>1</sup>.

## Order XCIX. EQUISETACEÆ.

Leafless branched plants with a striate hollow stem; each joint ending in a sheath which conceals the joining and encloses the base of the next joint. Sporules surrounded by elastic clavate filaments and enclosed in capsules arising from the peltate scales of terminal cones or spikes.—Rhizome creeping. Branches whorled. Cuticle abounding in silex. Only one genus.

1. *EQUISETUM* Linn. Horse-tail.

\* *Fertile stems mostly unbranched and succulent; barren stems with solid whorled branches, appearing later.*

1. *E. arven'sis* (L.); sterile st. with 6—19 furrows slightly rough, branches rough with 3 or 4 simple angles, *teeth of sheaths long acute 1-ribbed at the tip*, fertile st. simple with few lax distant sheaths.—*E. B.* 2020. *S. 1.* *H. F.* 60. *N.* 77.—Sterile st. many, procumbent or ascending; with many whorls of roughish solid usually simple branches with deep furrows and 3—4-toothed sheaths, their *lowest joint* (including its terminal sheaths) *exceeding the st.-sheath*; general outline narrowed upwards, usually naked at the end. Fertile st. short, with few

<sup>1</sup> (*N.*) refers to *Newman's British Ferns*, ed. 2 (1844); (*S.*) to *Sowerby's Ferns and Fern allies*; (*H. F.*) to *Hooker's British Ferns*. *Moore's Handbook of British Ferns*, ed. 3, may be consulted with much advantage, and *Milde in Nov. Act. Soc. Nat. Cur.* vol. xxxii.



(4—5) sheaths, appearing before the sterile ones. Sterile and fertile st. distinct.—*β. alpestre* (Wahl.); sterile st. short (2—3 in.) prostrate with ascending terminal point and second sub-erect branches.—Damp meadows. *β.* Mickle Fell, Teesdale. P. IV. E. S. I.

2. *E. pratense* (Ehrh.); sterile st. with about 20 striæ very rough with prominent points particularly above, branches simple with 3 or 4 simple angles, teeth of sheaths 1-ribbed but not to the tip, fertile st. simple with many crowded deeply toothed sheaths.—*E. umbrosum*, *H. F.* 59. *S.* 2. *N.* 63. *E. Drummondii* *E. B. S.* 2777.—Sterile st. 1—1½ ft. high, nearly naked below; with many whorls of slender solid branches in the upper part, having 3- or 4-toothed sheaths, their lowest joint and sheath falling short of the st.-sheath; general outline remarkably blunt. Branched fertile st. with larger sheaths and whorls of about 6 branches; simple fertile st. short (4—6 in.), with many loose and still larger yellowish-white sheaths with black prominent ribs upwards and 12—20 teeth.—Wet places, rare. P. IV. E. S. I.

3. *E. maximum* (Lam.); sterile st. nearly smooth with about 30 striæ and branches, branches rough doubly angular simple, teeth of sheaths 2-ribbed, fertile st. simple with many crowded large deeply-toothed sheaths.—*E. Telmateja*, *H. F.* 58. *S.* 3. *N.* 67. *E. fluviatile* Sm. not Linn., *E. B.* 2022.—Sterile st. 3—6 ft. high; occasionally bearing a small terminal spike; furnished from top to bottom with whorls of slender solid branches with 4 longitudinally furrowed angles and 4-toothed sheaths, their lowest joint and sheath exceeding the st.-sheath. Fertile st. stout, 1 foot or more high, with many very long pale-brown sheaths with 30—40 teeth; spike large.—Wet places. P. IV. E. S. I.

\*\* Sterile and fertile st. subsimilar, contemporaneous, branched.

4. *E. sylvaticum* (L.); sterile and fertile st. with about 12 furrows and many whorls of slender compound spreading or deflexed solid branches, sheath lax ending in 3 or 4 blunt lobes.—*E. B.* 1874. *H. F.* 61. *S.* 4. *N.* 59.—St. 12—18 in. high. Sheaths of the branches with 3 long acute teeth each 1-ribbed to its tip. Fertile st. occasionally simple. Spike blunt. General outline of sterile st. pyramidal, of fertile abrupt.—Wet shady places. P. IV. V. E. S. I.

\*\*\* Stems of one kind, with or without simple hollow whorled branches.

5. *E. limosum* (L.); stem nearly smooth with many slight

furrows, teeth of sheaths short rigid acute, branches simple whorled or none.—St. 2—4 ft. high. Sheaths rather short. Spike blunt.—*a. E. limosum* (L.); st. smooth, barren st. narrowing gradually upwards, branches short rigid slightly tapering upright and equalling st.-joints from green sheaths often wanting. *H. F.* 62.—*β. E. fluviatile* (L.); st. subglabrous. barren st. with a lax whiplike end, branches long slender tapering lax exceeding st.-joints from dark-brown sheaths rarely wanting.—In stagnant water. P. VI. VII. E. S. I.

6. *E. palustre* (L.); st. with 4—8 deep furrows branched throughout, sheaths loose pale with acute wedged shaped teeth tipped with brown and membranous at the edges, branches simple.—*E. B.* 2021. *H. F.* 63. *S.* 6. *N.* 43, 47 & 49.—St. slightly rough. Barren st. whipshaped at the end. Spike blunt. Sheaths coloured like the st. or paler; teeth brown with nearly transparent edges, ribs furrowed on the back. Branches usually barren or (*β. polystachium*) each ending in a spike, hollow, with shallow furrows, lowest joint of branch (often reduced to its sheath) falling short of the st.-sheath. Occasionally (*γ. nudum* DC.) the angles and teeth are fewer and the st. nearly or quite simple and dwarf.—Spongy bogs. *γ.* Sandy places. P. VI. VII. E. S. I.

7. *E. hyemale* (L.); st. simple very rough with 14—20 slender furrows, sheaths close whitish but the top and bottom black, teeth with slender black-brown very deciduous tips.—*E. B.* 915. *H. F.* 64. *S.* 8. *N.* 17.—St. 1—2 ft. high, simple, biennial, its central hollow equalling at least  $\frac{2}{3}$  of diameter. Spike apiculate. Sheaths widest at their top, at first green with black crenate rim after the teeth have fallen, then entirely black, and ultimately pale in the middle and black above and below.—*β. E. Moorei* (Newm.); st. annual very rough with about 12 furrows, sheaths loose white with the base black, teeth black-based rather persistent. *Phytol.* v. 19 (1853). *S.* 12. St. 1—2 ft. high, its central hollow equalling about  $\frac{2}{3}$  of diameter. Sheaths loose, pearly white; teeth long, usually light brown, whitish above, more persistent and longer.—Damp banks and woods. *β.* Wicklow. P. VII. VIII. E. S. I.

8. *E. trachy'odon* (A. Br.); st. simple or very slightly branched very rough with 8—12 furrows, sheaths close ultimately wholly black, teeth slender persistent.—*H. F.* 65. *E. Mackayi* *N.* 24. *S.* 9.—St. 1—3 feet high, simple or with solitary distant branches, biennial, its central hollow equalling  $\frac{1}{3}$  of diameter. Spike apiculate. Sheaths quite cylindrical, pale green with a black band beneath the teeth but ultimately wholly black.

Teeth much more persistent than in the preceding, usually black.—Near Belfast. Chevy Chase near Gort. P. VII. VIII. I.

9. *E. variegatum* (Schleich.); st. simple or slightly branched very rough with 4—10 furrows, *sheaths slightly enlarged upwards green below black above*, teeth blunt each tipped with a deciduous bristle.—*H. F.* 66. *N.* 31.—St. about a foot high, erect, usually simple except at the base or irregularly branched, its central hollow equalling  $\frac{1}{3}$  of diameter. Lower half of the sheaths green like the stem, upper part black; teeth persistent ovate, black in the centre, with a white membranous margin. Spike apiculate.—*E. Wilsoni* (Newm. 39. *S.* 10) is a large form.—*β. arenarium*; st. procumbent, usually more slender, teeth of the sheaths wedgeshaped. *E. variegatum* Sm., *E. B.* 1987. *S.* 11.—Wet places, or in water. *β.* Sandy places near the sea. P. VII. VIII. E. S. I.

### Order C. FILICES.

Leafy plants with a rhizome or trunk. L. or fronds usually circinate when young (Tribe VII. excepted), simple or divided. Fructification springing from the veins on the underside or at the edge of the l., of 1-celled capsules (thecae) which are stalked and have an elastic ring or sessile and without a ring.

\* *Capsules with an elastic marginal ring.*

#### Suborder I. POLYPODIACEÆ.

Capsules in dorsal or marginal clusters, opening transversely or irregularly. Young fronds circinate.

† *Clusters dorsal. Ring vertical, usually incomplete. Caps. opening transversely.*

Tribe I. *POLYPODIEÆ.* Clusters nearly circular, without an indusium, seated upon the back of the lateral veins.

1. *CRYPTOGRAMME.* Clusters circular, at length confluent, concealed by the reflexed margin of the frond.—Barren and fertile fronds dissimilar.

2. *POLYPODIUM.* Clusters circular, naked; edge of the frond flat, not reflexed.

3. *WOODSIA.* Clusters circular, with an inferior involucre divided at the edges into many capillary segments.

Tr. II. *ASPIDIEÆ*. Clusters nearly circular, covered by an indusium, seated upon the back of the lateral veins.

4. *LASTREA*. Indusium reniform, attached by the notch. Veins distinct after leaving the midrib, not uniting with those of the adjoining lobe.
5. *POLYSTICHUM*. Indusium circular, attached by the centre. Veins distinct after leaving the midrib.
6. *CYSTOPTERIS*. Indusium attached by its broad hooded case under the clusters, with a long fringed free extremity at first covering the capsules.

Tr. III. *ASPLENIEÆ*. Clusters oblong or linear, usually covered by an indusium opening longitudinally on one side, placed on the side of the lateral veins.

7. *ATHYRIUM*. Clusters oblong-reniform. Indusium opening towards the central vein or midrib, margin fringed.
8. *ASPLENIUM*. Clusters long, straight. Indusium opening towards the central vein or midrib, nearly flat.
9. *SCOLOPENDRIUM*. Clusters long, straight, 2 together. Indusia of each pair opening towards each other.
10. *CETERACH*. Lateral veins anastomosing; clusters attached to their middle on the side next the midrib, except in the lowest. Indusium (?) a narrow nearly erect membrane on the back of the vein. Whole back of the frond covered with chaffy scales.

[11. *GYMNOGRAMME*. Clusters oblong or linear on both branches of the forked vein, becoming confluent and covering the back of the frond. Indusium wanting.]

Tr. IV. *ADIANTEÆ*. Capsules covered by a marginal or submarginal elongated part of the frond, or by a separated portion of the cuticle resembling an indusium.

12. *BLECHNUM*. Capsules in a continuous line parallel to the midrib upon a longitudinal anastomosing part of the transverse veins, covered by a continuous scarious indusium.—Barren and fertile fronds dissimilar in our plant.
13. *PTERIS*. Capsules in a continuous marginal line covered by a continuous indusium formed of the reflexed margin.
14. *ADIANTUM*. Capsules marginal, oblong or roundish,

covered by distinct reflexed portions of the margin of the frond.

†† *Capsule opening irregularly, seated on a receptacle which ends a vein at the edge of the frond. Ring oblique, transverse, complete.*

Tr. V. *HYMENOPHYLLÆ.*

15. *TRICHOMANES.* Capsules on a long filiform receptacle within a cupshaped involucre of the same texture with the frond.

16. *HYMENOPHYLLUM.* Capsules on a narrow subclavate receptacle within a two-valved involucre of the same texture with the frond.

\*\* *Capsules without an elastic ring.*

Suborder II. *OSMUNDACEÆ.*

Young fronds circinate. Rachis woody. Capsules regularly 2-valved, stalked, in clusters at the extremity of the frond.

Tr. VI. *OSMUNDEÆ.*

17. *OSMUNDA.* Capsules clustered, arranged in a branched spike terminating the frond.

Suborder III. *OPHIOGLOSSACEÆ.*

Young fronds straight. Rachis succulent. Capsules regularly 2-valved, sessile, in clusters on a separate branch of the frond.

Tr. VII. *OPHIOGLOSSEÆ.*

18. *BOTRYCHIUM.* Capsules distinct, disposed in a compound spike attached to a pinnate or bipinnate frond.

19. *OPHIOGLOSSUM.* Capsules connate, disposed in a simple distichous spike attached to an undivided frond.

Suborder I. *Polypodiaceæ.* Tribe I. *Polypodieæ.*

1. *CRYPTOGRAMME R. Br.* Rock-brake.

1. *C. cris'pa* (R. Br.); barren fronds 2—3-pinnate, leaflets wedgedshaped or linear-oblong often bifid at the end, leaflets of the fertile fronds oblong.—*H. F.* 39. *Allosorus* Bernh., *N.* 103. *E. B.* 1160.—Fertile frond nearly triangular. Veins alternate,

often forked and each branch ending in a cluster having no indusium but concealed by the reflexed edge of the leaflet. Height 6—12 in. St. slender, very brittle.—Amongst loose stones on mountains. P. VII. *Parsley Fern*. E. S. I.

## 2. POLYPO'DIUM Linn. Polypody.

\* *Clusters at the end of a veinlet, other veinlets knobbed at the end and not reaching the edge. Rhizome without fronds at its end. Stipes jointed to rhizome.*

1. *P. vulgäre* (L.); fronds deeply pinnatifid, lobes linear-oblong somewhat serrate all parallel upper ones gradually smaller.—*E. B.* 1149. *H. F.* 2. *N.* 111.—Rhizome brown, densely scaly creeping. Fronds strapshaped. Clusters large, on the upper part of the frond. Lateral veins of the pinnæ with 4 branches. Pinnæ occasionally bifid at the end, sometimes deeply serrate or even (*P. cambricum* L.) doubly pinnatifid.—Shady banks, walls and old trees. P. VIII.—X. *Common Polypody*. E. S. I.

\*\* *Veinlets not knobbed but reaching the edge, all usually fertile, clusters near the end of each. Rhizome with fronds at its end. Stipes not jointed to rhizome.*

2. *P. Phegop'teris* (L.); fronds pinnate, pinnæ linear-lanceolate united at the base pinnatifid with linear-oblong blunt lobes, lowest pair of pinnæ turned downwards and forwards the rest upwards, clusters marginal.—*E. B.* 2224. *H. F.* 3. *N.* 115.—Rhizome nearly black, wiry, slightly scaly, creeping extensively. Fronds triangular. Pinnæ very acute, pointing upwards, rather hairy, connected by their whole width with the rachis; lowest pair quite distinct, with a minute stalk, standing forwards and pointing from the others.—Damp places, loving the spray of waterfalls. P. VII.—IX. E. S. I.

3. *P. Dryop'teris* (L.); fronds ternate glabrous, divisions pinnate, pinnæ pinnatifid blunt uppermost nearly entire, clusters marginal.—*E. B.* 616. *H. F.* 4. *N.* 123.—Rhizome black, wiry, creeping, slightly scaly. Stipe slender, brittle. The three divisions of the frond loosely spreading, the middle one rather the largest. Very young fronds resemble 3 little balls on wires. Not glandular.—Shady mountainous places. P. VI. VII.

E. S. I.

4. *P. Robertianum* (Hoffm.); fronds triangular subternate glandular-mealy, lower branches pinnate, pinnæ pinnatifid blunt uppermost nearly entire, clusters marginal.—*H. F.* 5. *P. cal-*

*careum* Sm., *E. B.* 1525. *N.* 131.—More erect and rigid than the preceding, always covered with very minute stalked glands giving a mealy character to the surface. Frond scarcely 3-fid, the lower branches being much smaller in proportion to the middle one; all the 3 erect, rigid.—On broken limestone ground. P. V.—VIII. E.

\*\*\* *Veinlets not knobbed, scarcely reaching the edge, simple or branched, each bearing a cluster below its end or that of its anterior branch. Rarely there is an oblique curved false indusium. Stipes not jointed to rootstock.*

5. *P. alpes'tre* (Hoppe); fronds lanceolate bipinnate, *pinnæ narrow-lanceolate with a broad base contiguous*, pinnules widest at their base acute pinnatifid with serrate lobes and branched veinlets.—*H. F.* 6. *S.* 49.—Fronds suberect, 1—3 ft. high. Much like *Athy. Filix-fœmina*.—High mountain-valleys. P. VII. VIII. S.

6. ? *P. flex'ile* (Moore); fronds linear-lanceolate bipinnate, *pinnæ ovate-lanceolate distant*, pinnules narrow at their base obovate bluntish serrate and with unbranched veinlets.—*N. ed.* 3. 203.—*P. rheticum* Fl. Dan. 2607 ?—Stipe short. Fronds much less divided, narrow, elbowed, spreading horizontally. Clusters crowded at the base of the frond, fewer upwards.—Abundant in Glen Prosen, Forfarshire. P. VII. VIII. S.

### 3. WOODS'IA R. Br.

1. *W. ilven'sis* (R. Br.); frond lanceolate hairy and chaffy beneath pilose above, *pinnæ oblong or ovate* pinnatifid, lobes very blunt nearly entire.—*E. B. S.* 2616. *H. F.* 8. *N.* 137.—Rhizome tufted. Spike jointed. Frond 1—5 inches long; *pinnæ* 4—6 lines long, mostly opposite.—Exposed alpine rocks. Glyder Fawr, N. Wales. Falcon Clints, Durham. White Coombe, Dumfries. P. VII. E. S.

2. *W. hyperborea* (R. Br.); frond linear-lanceolate or oblong pinnate glabrous or slightly hairy only beneath, *pinnæ triangular* pinnatifid or lobed, lobes 3—7 very blunt nearly entire.—*E. B.* 2023. *H. F.* 7. *N.* 143. *Acrostichum alpinum* Bolt. t. 42.—Rhizome tufted. Stipe jointed. Frond 1—3 inches long; *pinnæ* mostly alternate, a little longer than broad; pinnules 2—3 lines long.—Exposed alpine rocks. Ben Lawers, Perthshire. Clogwyn y Garnedd, Snowdon. P. VII. E. S.

Tribe II. *Aspidiæ*.

## 4. LAS'TREA Presl.

\* *Lateral veins simple or forked. Clusters on the simple veins or either or both branches.*

1. *L. Thelypteris* (Presl); rhizome slender far-creeping, fronds pinnate, pinnæ linear-lanceolate pinnatifid slightly downy but *without glands*, lobes oblong, clusters submarginal.—*H. F.* 13. *N.* 183.—Fronds lanceolate; earlier barren with flat lobes; later fertile with revolute-edged lobes; 2 or 3 lowest pairs of pinnæ decreasing in size. Lateral veins alternate, forked, extending to the edge. Clusters at length confluent, midway between the midrib and edge. Height 6—8 inches.—Marshy and boggy places. *P.* VII. VIII. *Marsh-Fern.* E. S. I.

2. *L. Oreopteris* (Presl); rhizome thick short, fronds pinnate, pinnæ linear-lanceolate pinnatifid *glandular* beneath gradually decreasing from about the middle of the frond to near the root, lobes oblong flat, clusters marginal.—*H. F.* 14. *E. B.* 1019. *N.* 187.—Fronds remarkably narrowed downwards, rising in a circle from a tufted rhizome, fragrant when bruised. Lobes blunt, entire; lateral veins simple or forked. Height 2—3 feet. Indusium often scarcely distinguishable.—Mountain heaths. *P.* VII. *Sweet Mountain-Fern.* E. S. I.

\*\* *Lateral veins branched or forked. Cluster upon the first upper lateral veinlet.*

3. *L. Filix-mas* (Presl); fronds lanceolate *subbipinnate*, pinnæ linear-lanceolate lowermost pair rather smaller than the second, pinnules oblong blunt or acutish serrate (not spinulose) attached by their whole width or often connected below, *clusters near the midvein*.—*E. B.* 1458. *N.* 198.—Fronds only slightly narrowed downwards and the lowest pinna of considerable size, rising in a circle from a tufted rhizome. Stipe and rachis nearly glabrous, yellow, or densely clothed with purple scales. Indusium very persistent, convex, with no marginal glands. Height 2—4 ft.—*α*; pinnules crowded linear-oblong blunt slightly confluent broad-based adpressed-serrate.—*β. L. Borreri* (Newm.); stipe and rachis very scaly, pinnules truncate subentire at the sides.—*γ. L. affinis* (Fisch.); pinnules less crowded longer acutish narrow and often slightly auricled at their base, teeth patent lower ones notched. Var. *incisa* Moore.—*δ. L. abbreviata* (DC.); clusters near the base of the confluent



crowded very blunt pinnules, about 1 ft. high.—Woods and banks. δ. Cumberland. Yorkshire. Wyck, Glouc. Glen Isla, Forfar. P. VI. VII. *Male Fern*. E. S. I.

4. *L. remo'ta* (Moore); fronds narrowly lanceolate, pinnæ triangular-lanceolate lowermost pair slightly smaller, pinnules acute with a narrow attachment deeply cut, clusters near the midvein.—*H. F.* 22.—Rhizome tufted. Scales ovate-acuminate and subulate. Much resembles *L. spinulosa*.—Marshy places. Windermere. P. VIII. IX. E.

5. *L. rig'ida* (Presl); fronds triangular-lanceolate bipinnate glandular, pinnæ triangular-lanceolate lowermost pair not smaller than the second, pinnules oblong blunt lobed and serrate with a narrow attachment, segments 2—5-toothed *not spinulose*, indusium persistent fringed with stalked glands, *stipe clothed with long-pointed 1-coloured scales*.—*H. F.* 16. *E. B. S.* 2724. *N.* 191.—Fronds erect, lanceolate with the lower pinnæ rather short and triangular, or triangular with the lower pinnæ long; upper pinnæ narrow; all pinnate. Pinnules truncate below. Covered with minute stalked glands. Height 1—2 feet.—Ingleborough, Arnside Knot, and near Settle. P. VII. VIII. E.

6. *L. crist'ata* (Presl); fronds linear-lanceolate or narrower subbipinnate glabrous, pinnæ short triangular-oblong pinnatifid or pinnate lowermost pair not smaller than the second, *pinnules oblong blunt or rarely acute serrate attached by their whole width and connected below* lowermost lobed and subtripinnatifid and superior and inferior nearly equal, stipe with broad ovate acute 1-coloured (pale) scales.—Fronds erect, 2 ft. high. Indusium without marginal glands.—*a*; fronds nearly linear, pinnæ pinnatifid, pinnules blunt; barren fronds broader. *H. F.* 17. *S.* 10. *N.* 203.—*β. L. uliginosa* (Newm.); fronds linear-lanceolate, pinnæ pinnate, pinnules acute, barren and late fertile fronds lanceolate, pinnæ subpinnate, pinnules blunt.—Bogs and boggy heaths, rare. P. VIII. E.

7. *L. spinul'osa* (Presl); fronds oblong-lanceolate bipinnate glabrous, pinnæ triangular-oblong or -lanceolate lowermost pair scarcely smaller than the second, *pinnules ovate-oblong acute incise-serrate with a narrow attachment inferior* lowermost often largest, stipe with ovate acute 1-coloured (pale) scales, rhizome stout creeping.—*S.* 12. *N.* 203. *H. F.* 18.—Height 3—4 ft.; fronds nearly erect. Upper pinnules narrowed and decurrent below. Indusium without marginal glands.—*a*; pinnæ triangular oblong lowermost pair not smaller, pinnules with a stalklike attachment lowermost nearly equal, stipe with ovate acute scales. "Spores wharted."—Marshy places and wet woods. P. VIII. IX. E. S. I.

8. *L. dilatata* (Presl); frond triangular-lanceolate or -ovate bipinnate, lower pinnæ unequally triangular lowest pair not shorter than the second, pinnules oblong with a narrow attachment pinnatifid or pinnate inferior ones largest, segments spinous-serrate, *stipe clothed with long pointed scales with a dark centre and diaphanous margin*, rhizome tufted.—*H. F.* 19. *L. multiflora* N. 215.—Caudex usually erect. Fronds 2—4 ft. high, arched, often drooping, convex, more or less clothed with stalked glands when young; on young or starved plants often triangular, never so on older and perfect ones. Indusium with marginal glands. The largest scales of the full-grown plant should be examined. “Spores winged and crested.”—*L. collina* (Newm.) has a triangular-ovate-prolonged frond and ovate blunt bluntly mucronate-serrate pinnules. *N.* 223.—*L. glandulosa* (Newm.) has a broad lanceolate frond covered with stalked glands beneath and the scales on the stipe often nearly without the dark centre. *Deak. Fl. Brit. f.* 1612.—Woods, banks, &c. P. VIII. IX. E. S. I.

9. *L. æmula* (Brack.); frond triangular or triangular-ovate bipinnate, lower pinnæ unequally triangular lowest pair longest, pinnules prolonged-triangular with a narrow attachment pinnatifid or pinnate inferior lower ones largest, segments spinous-serrate, *stipe clothed with long narrow laciniate 1-coloured scales*.—*H. F.* 20. *Nephrodium fœniseeii* Lowe! *L. recurva* N. 225.—*Frond* 1—2 feet long; the lower pinnæ much the largest. *Pinnules and segments concave above*. Stipes, rachis, and fronds with many *globose sessile glands*. Sweet-scented.—Rocky shady places. P. VIII. IX. E. S. I.

### 5. POLYS'TICHUM Roth.

1. *P. Lonchitis* (Roth); *fronds rigid* linear pinnate, pinnæ not lobed serrate spinous their base auricled above oblique below.—*N.* 163. *H. F.* 9. *E. B.* 797.—*Stipe* very short. Fronds narrow, very rigid and leathery. Pinnæ overlapping and twisted (most in the Irish, much less so in the Welsh plant), lower ones usually auricled both above and below.—Young simply pinnate fronds of the next species are often much like this plant.—Alpine rocks. P. VII. *Holly Fern*. E. S. I.

2. *P. aculeatum* (Roth); *fronds rigid* linear or lanceolate bipinnate, *pinnules obliquely decurrent*.—*N.* 169. *H. F.* 11. *Aspidium lobatum* Kunze. *A. aculeatum* and *A. lobatum* Sm.—*Stipe* usually short. Frond 1—2 feet high. First upper pinnule of each pinna longer than the others, its lower side (next the main rachis) usually nearly straight, its upper acutely auricled and forming an acute angle with the lower and with

the partial rachis at the point of attachment. In young plants the pinnæ are serrate or pinnatifid or with one or more pinnules distinct. A few of the lowest pinnules are often slightly stalked, but very differently from those of *P. angulare*.—*A. lobatum* (Sw.) has the pinnæ less divided than in the type of the species and the fronds linear-lanceolate and more rigid. *H. F.* 10.—Hedge-banks. *P.* VII. VIII. E. S. I.

3. *P. angulare* (Newm.); fronds lax drooping lanceolate bipinnate, pinnules truncate or obtuse-angled below distinctly stalked. *N.* 173. *H. F.* 12.—Stipe usually long. First upper pinnule scarcely longer than the others, its lower side rounded below, its upper with a large bluntish auricle and forming an obtuse angle with the lower, at the top of the short stalk which is nearly at right angles with the partial rachis; all short, broad and bluntish; or first upper pinnule longer and deeply pinnatifid, all more acute; or pinnules all narrower and acute.—Sheltered woods and hedge-banks, chiefly in the West. *P.* VII. VIII. E. S. I.

#### 6. CYSTOPTERIS Bernh. Bladder-Fern.

1. *C. fragilis* (Bernh.); frond lanceolate bipinnate, pinnæ ovate or ovate-lanceolate, pinnules oblong-ovate or cordate-ovate pinnatifid or cut.—*a. vera*; usually bipinnate, pinnules rather narrowed below, veins ending at tip of term. teeth or if pinnule emarginate in the 2 teeth *not in the notch*. Sporules prickly.—*a. C. anthriscifolia* (Roth); pinnules ovate acute cut, segments oblong toothed. *Cystea fragilis* Sm., *N.* 155.—*b. C. cynapifolia* (Roth); pinnules obovate cut, segments obovate toothed or retuse at the end. *C. fragilis* E. B. 1587.—*c. C. angustata* (Sm.); pinnules lanceolate acute cut, segments lanceolate-oblong cut, teeth acute. *N.* 156.—*β. C. dentata* (Sm.); frond often only subbipinnate, pinnules broadest below blunt bluntly toothed, veins as in *a.* *H. F.* 23. *N.* 154. *C. fragilis* Roth. Clusters more marginal, often ultimately confluent. In well-grown plants the pinnules are suddenly widened to their full extent just above their narrow stalklike base. Spores warted.—*γ. C. Dickieana* (Sim), pinnæ usually overlapping, pinnules broad blunt, veins reaching the emarginate end, clusters scattered, spores verrucose.—*δ. C. alpina* (Desv.); frond bipinnate, pinnæ ovate, pinnules ovate deeply pinnatifid with broadly and shortly linear segments partly cloven, veins reaching the blunt end. *E. B.* 163. *H. F.* 24. Fronds much divided but compact and close.—I have placed these plants under one species with much doubt.—Rocks and walls. *γ.* In a damp cave by the sea near Aberdeen. *δ.* Teesdale. *Mr. J. Backhouse!* *P.* VII. VIII. E. S. I.

2. *C. montana* (Link); fronds triangular tripinnate, pinnæ and pinnules spreading, lobes pinnatifid with linear notched segments.—*H. F.* 25. *S.* 24.—Fronde in shape like those of *Polypodium Robertianum*, small, short, very finely divided. Lower pair of pinnæ much the largest; their lower larger than their upper pinnules. Stipe long, slender. Rhizome creeping extensively, black.—Breadalbane and Grampian Mountains. *P.* VII. VIII. ?  
S.

### Tribe III. *Asplenieæ*.

#### 7. *ATHYRIUM* Roth. Lady Fern.

1. *A. Filix-fœm'ina* (Roth); frond lanceolate pinnate or pinnatifid, pinnæ linear, pinnules linear-oblong deeply serrate or pinnatifid.—*H. F.* 35.—*a.* *A. rhœticum* (Roth); frond bipinnate pinnules toothed narrowly triangular-lanceolate convex remote not connected toothed, upper ones minute confluent, clusters ultimately confluent. *N.* 245. *A. convexum* Newm. Pinnules long, narrow, with deflexed edges, enclosing the clusters; their attachment broad; segments gradually decreasing from the base of pinnule.—*β.* *A. Filix-fœmina* (Roth); frond bipinnate, pinnules pinnatifid oblong-lanceolate bluntish flat remote not connected, segments oblong patent with lateral and terminal sharp teeth. *N.* 237. Frond much divided dark green. Pinnules much narrowed at their base. Clusters distinct. *A. latifolium* (Bab.) is apparently only an extreme state of this.—*γ.* *A. molle* (Roth); frond pinnate, pinnules toothed oblong blunt or slightly pointed flat remote all connected by wing of midrib or lower ones distinct, segments ovate bidentate lowest with 3 teeth uppermost with 1 tooth. *N.* 245. Frond scarcely more than pinnate, bright green. Pinnules attached by a broad decurrent base. Clusters distinct, in 2 rows. Sometimes (*A. trifidum* Roth?) the pinnules are much less connected, moderately cut, have a narrow attachment, and lobes with more but connivent teeth.—There are innumerable subvarieties.—Wet shady places. *P.* VI. VII. E. S. I.

#### 8. *ASPLENIUM* Linn. Spleenwort.

\* *Ultimate subdivisions with a distinct midvein.*  
*ASPLENIUM* Newm.

[1. *A. fontanum* (Presl); frond linear-lanceolate bipinnate, pinnæ oblong-ovate, pinnules obovate-cuneate with few spinous-mucronate teeth.—*E. B.* 2024. *H. F.* 34.—Fronde about 4 in. long.—A very doubtful native. Stations want confirmation, to see if it was planted. Formerly on Amersham Church, Bucks !

Wybourn, Westm. *Hudson*. Northumberland. *J. Backhouse*.  
Tan y Bwlch and Tremadoc, Merionethshire. Ashford, Hants!  
P. VI.—IX.] E.

2. *A. lanceolatum* (Huds.); fronds lanceolate bipinnate, pinnules obovate deeply and sharply toothed or lobed, clusters short nearly marginal.—*E. B.* 240. *H. F.* 32. *N.* 249.—FronDS sometimes nearly linear and simply pinnate, always narrowed at the base. Clusters oblong, ultimately rather confluent into roundish masses.—Rocks and walls, rather rare. P. VI.—IX.  
E. I.

3. *A. Adiantum-nigrum* (L.); fronds ovate-triangular or triangular-prolonged twice or thrice pinnate, pinnæ and pinnules triangular sharply toothed, clusters long central.—*E. B.* 1950. *N.* 225.—Clusters 2 or 3 times as long as in the preceding, placed near the midrib and ultimately confluent in oblong masses often covering the whole under surface of the pinnule.—*a*; fronds about as long as the stipe ovate-triangular, pinnæ and pinnules triangular-prolonged, ultimate subdivisions blunt.—*β*. *A. Serpentina* (Tausch); pinnules triangular very broad, lobes ovate blunt.—*γ*. *A. acutum* (Bory); fronds much shorter than the stipe triangular-prolonged, pinnæ and pinnules lanceolate-attenuate, ultimate subdivisions very acute. *N.* 231.—Rocks and walls. *β*. "Serpentine rocks of Cabrach, Aberdeenshire." *T. Moore*. *γ*. South-west of Ireland. P. VI.—IX. *Black Spleenwort*.  
E. S. I.

4. *A. Trichomanes* (L.); frond linear pinnate, pinnæ roundish-ovate crenate, veins forked below the clusters.—*E. B.* 576. *H. F.* 29. *N.* 285.—*Rachis* black, keeled beneath. Pinnæ scarcely oblique; both edges rounded and crenate except at the base, upper often bluntly auricled below.—*β*. *A. anceps* (Sol. ?); pinnæ oblong blunt wedgeshaped below upper edge and end crenate-dentate lower entire. Pinnæ oblique, lower edge nearly straight; lower pinna much the smaller.—A curious variety is occasionally found with its pinnæ deeply but irregularly pinatifid with linear notched segments.—Rocks and walls. *β*. Killarney. P. V.—X. *Common Spleenwort*.  
E. S. I.

5. *A. viride* (Huds.); fronds linear pinnate, pinnæ roundish-ovate or rhomboidal crenate, veins simple or forked beyond the sori.—*E. B.* 2257. *H. F.* 30.—*Rachis* green, not keeled. Sori at length confluent.—Rocks and mountains. P. VI.—X.  
E. S. I.

6. *A. marinum* (L.); fronds linear simply pinnate, pinnæ stalked ovate or oblong serrate unequal and wedgeshaped at

the base.—*E. B.* 392. *H. F.* 31. *N.* 275.—Varying greatly in size. Sori not confluent.—Maritime rocks. *P. VI.—X.*  
E. S. I.

\*\* *Ultimate subdivisions without a distinct midvein.*  
AMESIUM Newm.

7. *A. Ruta-murária* (L.); fronds bipinnate *pinnules rhomboid-wedgeshaped* notched or toothed at the end, *indusium jagged*.—*E. B.* 150. *H. F.* 28. *N.* 261.—Fronds 3—4 in. long.—Rocks and old walls. *P. V.—IX. Wall-Rue.* E. S. I.

8. *A. german'icum* (Weiss); fronds simply and alternately pinnate, *pinnæ narrow-wedgeshaped*, blunt the lowermost ternate, *indusium entire* at the edge.—*H. F.* 27. *N.* 265. *A. alternifolium* Sm., *E. B.* 2258.—Fronds 3—4 in. long.—Rocks, very rare. *P. VI.—IX.* E. S.

9. *A. septentrionåle* (Hull); fronds 2- or 3-cleft, *pinnæ very long-lanceolate bifid*, *indusium entire*.—*E. B.* 1017. *H. F.* 26. *N.* 269.—*Pinnæ very narrow*, narrowing gradually downwards, with 1 or 2 short bifid lateral teeth, and bifid at the end.—Dry clefts of rock, rare. *P. VI.—X.* E. S.

#### 9. SCOLOPEN'DRIUM Sm. Hart's-tongue.

1. *S. vulgäre* (Sym.); frond oblong strapshaped smooth simple with a cordate base, *stipe shaggy*.—*E. B.* 1150. *H. F.* 37. *Phyllitis N.* 289.—Fronds 1—2 feet long, acute, often crisped and multifid.—Damp shady places. *P. VII. VIII.* E. S. I.

#### 10. CETE'RACH Willd. Rustyback.

1. *C. officinarum* (Willd.); fronds pinnatifid covered beneath with dense scales, *pinnæ alternate or opposite blunt sessile*.—*Notolepum N.* 293. *Asplenium L., H. F.* 36. *E. B.* 1244.—Fronds 3—6 in. long, green and smooth above, wholly covered by very many scales beneath, amongst which the capsules are almost hidden.—Old walls and rocks. *P. IV.—X.* E. S. I.

#### 11. GYMNOGRAM'ME Desv.

[*G. leptophylla* (Desv.); fronds pinnate or bipinnate glabrous, *pinnules wedgeshaped deeply lobed*.—*H. F.* 1. *N. ed.* 3. 11. *S.* 48.—Frond 1—4 in. high. *Stipe purple.* *Pinnæ and pinnules alternate*.—Banks. Jersey. *A. IV.*]

Tribe IV. *Adiantea*.12. BLECH'NUM *Linn.* Hard Fern.

1. *B. boreale* (Sw.); barren fronds pinnatifid with broadly-linear rather blunt pinnæ, fertile frond pinnate with linear acute pinnæ.—*E. B.* 1159. *H. F.* 40. *Lomaria Spicant N.* 89.—Each lateral vein of the fertile pinnæ extends halfway to the edge, then turns at right angles and proceeds up the pinna until it reaches the next vein. Capsules attached in a continuous row to the longitudinal portions of the combined lateral veins.—Stony and heathy places. *P. VII.* E. S. I.

13. PTE'RIS *Roth.* Brakes or Bracken.

1. *P. aquilina* (L.); fronds tripartite, branches bipinnate, pinnules linear-lanceolate the lower ones usually pinnatifid, segments oblong blunt.—*E. B.* 1679. *H. F.* 38. *N.* 93.—Fronds annual, 1—5 feet high, very much divided, with spreading branches. Capsules attached to the marginal vein, lying upon a fine membrane and covered by the membranous continuation of the epidermis. Inferior pinnules pinnatifid or sinuate or entire.—Woods and heaths. *P. VII.* E. S. I.

14. ADIAN'TUM *Linn.* Maiden-hair.

1. *A. Capillus-Veneris* (L.); frond irregular, branches and roundish-wedged-shaped lobed thin pinnules alternate, lobes of the fertile pinnules terminated by a transversely linear-oblong reflexed lobe covering several roundish clusters, sterile lobes serrate.—*E. B.* 1564. *H. F.* 41. *N.* 83.—Rhizome blackish, shaggy. Fronds 6—12 in. high. Stipe and rachis slender, nearly black. Pinnules not jointed to the partial stalks.—Damp rocks near the sea in the south and west. *P. V.—IX.* E. I.

Tribe V. *Hymenophyllea*.15. TRICHOM'ANES *Linn.*

1. *T. radicans* (Sw.); fronds 3 or 4 times pinnatifid glabrous, segments uniform linear, involucre solitary in the axils of the upper segments, receptacle at first included ultimately very prominent.—*E. B.* 1417. *T. speciosum* Willd., *N.* 305.—Fronds rather triangular, very much divided, 4—8 in. long, formed of hard wiry branched ribs each with a rather membranous wing. Rhizome black, downy, very long. Involucres scarcely winged.

—The form called *Andrewsii* has lanceolate fronds and winged involucre. *N.* 315.—Very damp shady places, rare. Formerly at Bellbank, Yorkshire! (exactly Bolton's t. 30). *N.* Wales! Arran! (S.). Killarney! *P.* IX. X. E. S. I.

### 16. HYMENOPHYLLUM Sm. Filmy-Fern.

1. *H. tunbridgen'se* (Sm.); fronds pinnate, pinnæ distichous, segments linear undivided or bifid spinously-serrate, involucre compressed spinously-serrate, rachis broadly winged.—*E. B.* 162. *H. F.* 43. *N.* 321.—Slender, delicate and small. Rhizome very long, threadshaped. Pinnæ, rachis, and involucre in the same plane. Inv.-valves adpressed throughout the greater part of their length, slightly gibbous at the base.—Amongst moss in damp and shady places. *P.* VII. E. S. I.

2. *H. Wilsoni* (Hook.); fronds pinnate, pinnæ recurved, segments linear undivided or bifid spinously-serrate, involucre inflated entire, rachis slightly bordered.—*E. B. S.* 2686. *H. F.* 44. *H. unilaterale* Bory?—Resembling the preceding, but the pinnæ curve backwards and the involucre forwards. Inv.-valves convex or gibbous throughout, touching only by their edges which are quite entire.—Amongst moss in damp and shady places. *P.* VII. E. S. I.

## Suborder II. Osmundaceæ. Tribe VI. Osmundææ.

### 17. OSMUN'DA Linn. Flowering-Fern.

1. *O. regalis* (L.); fronds bipinnate, pinnules oblong nearly entire dilated and slightly auricled at the base, clusters paniced terminal.—*E. B.* 209. *H. F.* 45. *N.* 331.—Fronds erect or drooping, 1—8 feet high.—Boggy places. *P.* VII.—IX. E. S. I.

## Subord. III. Ophioglossaceæ. Tr. VII. Ophioglosseæ.

### 18. BOTRICH'IUM Sw. Moon-wort.

1. *B. Lunária* (Sw.); frond pinnate solitary, pinnæ lunate or fanshaped notched or crenate.—*E. B.* 318. *H. F.* 48. *N.* 137.—Height 3—6 in. Pinnæ with veins radiating from the petiole, sometimes deeply notched. Fronds usually solitary, but sometimes two on the same stalk.— $\beta$ . *B. rutaceum* (Sw.); frond triangular-rhomboidal pinnatifid, pinnæ 7—9 linear incise-serrate decreasing upwards. *N. ed.* 3. 332. See *Milde Fil. Europ.* 195.—Pastures.  $\beta$ . Sands of Barry, Forfar. *P.* V.—VII. E. S. I.



19. OPHIOGLOS'SUM *Linn.* Adder's-tongue.

1. *O. vulgatum* (L.); frond ovate blunt, epidermal cells flexuose, spores tubercled.—*E. B.* 108. *H. F.* 46. *N.* 349.—Height 4—12 in., erect. Spike clubshaped, usually rather longer than the frond, sometimes very long.—*β. ambiguum* (C. & G.); rhizome often producing two fronds from the same joining, frond oblong-lanceolate narrowed below 2—4 in. high.—Pastures. *β.* Orkney. Scilly. *P. V.* VI. *E. S. I.*

2. *O. lusitanicum* (L.); frond linear-lanceolate, epidermal cells straight, spores smooth.—*H. F.* 47. *N.* ed. 3. 331. *S.* 47.—Height 1—2 inches, erect.—Horn Head, Co. Donegal. Petit Bo Bay, Guernsey. *P. I.* *I.*

## Order CI. MARSILEACEÆ.

Creeping plants with alternate erect leaves, circinate in bud. Fructification consisting of globular coriaceous axillary bodies with 3 or 4 cells and containing sacs including either other bodies that germinate or loose granules.

1. PILULARIA. Capsules solitary, nearly sessile, globose, coriaceous, 2—4-celled. Cells containing bodies of two kinds—granules, and membranes containing minute grains.

1. PILULA'RIA *Linn.* Pillwort.

1. *P. globulifera* (L.); ped. erect, caps. 4-celled.—*E. B.* 521. *H. F.* 57. *N.* 393.—Rhizome slender, creeping, producing leaves and roots at regular intervals. L. very slender, erect. Caps. pubescent, slightly stalked, axillary, nearly spherical, hairy.—Margins of ponds and lakes. *P. VI.* VII. *E. S. I.*

## Order CII. LYCOPODIACEÆ.

Leafy plants with simple imbricate leaves, or stemless with erect subulate leaves. Fructification of axillary sessile capsules with 2 or 3 valves and no ring, including minute powdery matter or spores.

\* *Capsules not opening.—Leaves radical.*

1. ISOËTES. Caps. in pouches formed of the swollen bases of the leaves. Spores of two kinds attached to filiform receptacles; those of the outer leaves large, of the inner very small.

\*\* *Capsules bursting.*—*With leafy stems.*

2. LYCOPODIUM. Caps. of one kind, 1-celled, containing many minute spores.
3. SELAGINELLA. Caps. of two kinds, small containing many minute spores, or larger and containing about 4 large spores.

1. ISOËTES *Linn.* Quillwort.

- \* No persistent leaf-bases. Caps. not wholly covered by the membranous edge of the pouch. Back of swollen leaf-base smooth.—Aquatic.

1. *I. lacus'tris* (L.); l. subulate roundish-quadrangular with 4 longitudinal jointed tubes upright dark green, *larger spores bluntly tubercled rather mealy*, tubercles overtopped by the valve-edges.—*E. B.* 1084. *H. F.* 55. *R.* vii. 1.—Corm with longitudinal furrows. L. slender, broad and flat at the base, but elsewhere between cylindrical and quadrangular, 2—6 in. long.—*I. Morei* (Moore) is a form with exceedingly long leaves. *J. of B.* xvi. t. 199.—Usually cæspitose on the sandy and stony bottom of lakes and pools in hilly districts. P. VI. E. S. I.

2. *I. echinos'pora* (Dur. !); l. subulate roundish-quadrangular with 4 longitudinal jointed tubes patent pale green, *larger spores very acutely tubercled*, tubercles overtopping the valve-edges.—*J. of B.* i. t. 1.—Much like Sp. 1. Corm not furrowed. L. turning yellow, less rigid, flattened and dilated below. Spores covered with long acute spine-like tubercles, not mealy.—Usually solitary on the muddy bottom of pools and lakes in hilly districts. Llanberis, N. Wales. Loch of Park near Aberdeen. Ben Voirlich, Dumbartonshire. Gap of Dunloe, Kerry. P. VI. E. S. I.

- \*\* Corm more or less covered by the persistent hardened leaf-bases. Caps. wholly covered by the membranous edge of the pouch. Swollen leaf-base with a central longitudinal rugose band on the back.—Terrestrial.

[*I. Hys'trix* (Dur.); l. filiform plane-convex obscurely tubular, persistent l.-bases short blackish each with 2 long horns and an interm. tooth, larger spores white and bluntly tubercled.—*I. Duriei* H. F. 56.—Corm. small, rarely if ever quite naked. L. very slender, ultimately very much enlarged at the base to enclose the capsule, 1—2 in. long, annual.—Dampish sandy and stony places. L'Ancrese, Guernsey. *Mr. G. Wolsey.* P. V. VI.]

## 2. LYCOPO'DIUM Linn. Club-moss.

*St. creeping, prostrate. Caps. in term. spikes, with bractlike leaves.* LEPIDOTIS Palis.

1. *L. clavatum* (L.); l. scattered imbricate incurved hair-pointed, *spikes stalked* 2 or 3 together cylindrical, scales ovate-triangular membranous finely incise-serrate.—*E. B.* 224. *H. F.* 49. *N.* 353.—*St.* long. Branches short, ascending. Spikes on long stalks, pale yellow. Scales on the ped. irregularly disposed in whorls.—Heaths. *P.* VII. VIII. *Common Club-moss.* *E. S. I.*

2. *L. annotinum* (L.); l. loosely scattered lanceolate mucronate serrulate, *spikes sessile* solitary terminal, scales roundish shortly acuminate membranous and jagged—*E. B.* 1727. *N.* 361.—*St.* very long. Branches rather long, erect, each year's growth marked by a constriction. Spikes cylindrical, greenish yellow, not persistent.—Stony mountains. Very rare in Caernarvonshire and Cumberland. Common in the Highlands of Scotland. *P.* VIII. *E. S.*

3. *L. alpinum* (L.); l. *in four rows*, imbricate acute keeled entire, spikes sessile solitary terminal, scales ovate-lanceolate flat, branches erect clustered forked level-topped.—*E. B.* 234. *H. F.* 53. *N.* 365.—*St.* long, prostrate; also a subterranean rhizome. Fertile branches usually twice dichotomous, each division ending in a short cylindrical yellowish-green spike rather thicker than the branch.—Stony moors. *P.* VIII. *Savin-leaved Club-moss.* *E. S. I.*

4. *L. inundatum* (L.); l. secund linear subulate, spikes terminal sessile leafy solitary upon short erect branches.—*E. B.* 239. *H. F.* 51. *N.* 369.—*St.* short, prostrate, rooting. Branches few, simple, fertile. *Bracts subulate* from a dilated base.—Boggy heaths. *P.* VIII. IX. *E. S. I.*

\*\*\* *St. decumbent below, then erect. Caps. in axils of upper leaves, solitary. No spikes. L. all alike.* PLANANTHUS Palis.

5. *L. Selágo* (L.); l. *in eight rows crowded* uniform linear-lanceolate acuminate, caps. not spiked but in the axils of the common leaves, *st. erect* forked level-topped.—*E. B.* 233. *H. F.* 54. *N.* 375.—*St.* short, erect or slightly decumbent, densely leafy. No separate spikes. At the extremity of the stems a few curious viviparous buds may usually be found; they are well illustrated by Mr. Newman (p. 378). Occasionally the stems in sheltered situations become much lengthened.—Heaths, chiefly on mountains. *P.* VI.—VIII. *Fir Club-moss.* *E. S. I.*

3. SELAGINEL'LA *Spring.*

1. *S. spinulo'sa* (A. Br.); l. uniform scattered lanceolate ciliate, spikes terminal solitary sessile leafy upon short erect branches.—*E. B.* 1148. *N.* 371.—St. prostrate, much branched, rooting, slender. Flowering branches simple, short, erect. Small spores muricate in 2-valved reniform caps.; large spores papillose in 3—4-valved capsules.—Boggy spots chiefly in mountainous districts. P. VIII. E. S. I.

B. Stems of one or more parallel tubes, verticillately branched. Nucules and globules on the branches.

## Order CIII. CHARACEÆ.

Leafless branched plants with stems formed of one or more parallel tubes. Two kinds of fructification; round red globules (anthers?) formed of 8 valves, enclosing cells containing granular matter and spiral filaments; oval nucules formed of 1 cell with 5 filaments folded spirally round it and containing minute granules which appear at last to unite into a single seed.—Plants aquatic. The position of this Order is very doubtful<sup>1</sup>.

1. NITELLA. Globules and nucules at the forking of the branchlets. Crown of nucule of 10 cells in 2 rows lying upon each other, the upper smaller, usually deciduous.—No stipulodes. Stem of one tube.
- 2: CHARA. Globule taking the place of central bract, below the nucule. Crown of nucule of 5 equal cells in one row, persistent.—Stipulodes in 1 or 2 whorls. Usually with cortical cells.

1. NITEL'LA *Ag.*

\* *Globules terminal in the forking of the branchlets. Nucules below the globules. Segments nearly equal.*—NITELLA Braun.

† Branchlets only once divided into 1-jointed segments.

1. *N. flex'ilis* (Ag.); monœcious, st. slender flexible transparent, branchlets pointed not mucronate, fertile walls usually lax, nucules 2 or 3 together with 8 or 9 spires.—*Atl. Fl. Par.* 40.—Usually slender, flexible, light green, often encrusted.—Ponds. VI. VII. E.

<sup>1</sup> See Messrs. Groves' valuable paper in the *J. of B.* xviii. and xix., where are figures of nearly all the species.

2. *N. opaca* (Ag.); dioecious, st. slender flexible transparent, branchlets mucronate, fertile whorls usually dense, *nucules* 1—3 together *naked with 6 blunt prominent spires*.—*J. of B.* xviii. t. 210.—Rather slender, turning nearly black. Whorls with nucules dense, with globules usually more lax.—Ponds and ditches. A. V. VI. E. S. I.

3. *N. capitata* (Ag.); dioecious, st. slender flexible transparent, branchlets mucronate, fertile whorls . . . , *nucules with acute prominent spires and a mucilaginous coating*.—Small, slender, yellowish brown.—Llyn Idwal, Caern. (teste A. Braun!). Stouting, Kent. *A. Braun.* Cornwall. *A. Bennett.* A. E.

[*N. syncar'pa* (Lutet); more slender than *C. opaca* with simple branchlets on the female plant, and *nucules with a mucilaginous coating and faintly marked spires*; is probably a native.]

†† Branchlets repeatedly divided; terminal segments of 2 or more cells, last usually like a mucro.—Monœcious.

4. *N. transhucens* (Ag.); st. thick equal flexible transparent, last segments of branchlets of 2 or 3 cells, sterile branchlets only once divided (at the end) into 2—4 *minute rays*, nucules small usually in threes with 7 or 8 spires.—*E. B.* 1851.—A very large plant. Sterile branchlets appear simple; the rays resemble minute points at their end.—Deep water. A. VIII. E. S. I.

5. *N. mucronata* (Kütz.); st. equal flexible transparent, branchlets nearly equally forked or trifid *tipped with a minute acutely conical cell, nucules and globules together* at the forkings of the branchlets.—*Atl. Fl. Par.* 40.—Rather thick. Dark green. Secondary branchlets once or twice forked, terminal subdivisions rather shorter than the others. Nucules with 7 or 8 spires.—Marsh-ditch near W. Grinstead, Suss. A. VII. E.

6. *N. gracilis* (Ag.); st. equal flexible transparent, slender *branchlets in lax whorls* twice or thrice nearly equally forked their *terminal segments 2- or 3-celled* strongly mucronate, nucules and globules each solitary but together at the forkings of the branchlets.—*E. B.* 2140. *Atl. Fl. Par.* 41.—Very small and slender. Nucules subglobose with 6 or 7 spires.—St. Leonard's Forest, Suss. Glen Cullen, Co. Dublin. A. IX. E. I.

7. *N. tenuis'sima* (Kütz.); st. equal flexible transparent, *branchlets in very dense compact subglobose whorls* twice or thrice equally forked in 3—6 segments of 2 or 3 cells their terminal cell very slender and acute, nucules and globules each solitary but

together at the forkings of the branchlets.—*Atl. Fl. Par.* 41.—Very small and slender with long internodes, 1—4 in. high, usually dark green. Branchlets *forming little globose compact masses*, often much incrustated. Nucules globose with 9 spires, minute but much exceeding the diameter of the branchlets; globules much larger.—Peaty ditches in Fens of Cambridgeshire. A. VIII. E.

\*\* *Globules lateral on the forkings of the branchlets, amongst the nucules. Segments unequal. Monœcious.*—TOLYPELLA Leon.

8. *N. glomerata* (Chevall.); st. stoutish, sterile branchlets 6—12 simple blunt 3—5-jointed, fertile branchlets many densely crowded once unequally 3—4-forked with blunt 2—5-jointed divisions, globules usually stalked with one or more nucules of 8 or 9 faint spires.—*Atl. Fl. Par.* 41.—Much incrustated and brittle.—Brackish ponds and ditches chiefly. A. IV. E. I.

9. *N. prolifera* (Kütz.); st. stout, sterile branchlets 6—20 simple acute 3—5-jointed, fertile branchlets many densely crowded divided into 3—4 acute 2—4-jointed divisions, globules sessile (?) with one or more nucules of 9 or 10 spires.—A very large plant. Branchlets even 7 inches long and very stout.—Very rare. Henfield, Suss. Glasnevin, Dublin. E. I.

10. *N. intricata* (A. Br.!) ; st. stout, sterile branchlets 6—10 once or twice divided acute, fertile branchlets many densely crowded divided once or twice into 4—5-celled acute divisions, globules stalked with one or more nucules of 8—9 spires.—Known by its divided branchlets.—Ponds and ditches. A. IV. V. E. I.

#### CHA'RA Ag.

\* Stem composed of a single tube.

A. *Stipulodes rudimentary, in one ring. Diœcious.*

1. *C. stelligera* (Bauer); st. thick equal transparent its lower nodes being *white starlike knobs*, branchlets 4—6 of 2 or 3 long joints bearing 1 or 2 unequal 1-jointed bracts, stipulodes exceedingly small, nucules with 9 spires subglobose.—*N. obtusa J. of B.* xix. t. 216.—Very large, resembling *N. translucens*. Nucules with very thick spires and very small crown.—Filby Broad, Norf. VIII. E.

B. *Stipulodes long, in one ring. Monœcious.*—  
LYCHNOTHAMNUS Rupr.

2. *C. alopecuroides* (Del.); st. opaque, 3—5-jointed branchlets 6—8 in a whorl their terminal joints much the shortest and

forming an acute point, stipulodes needleshaped, bracts 5 or 6 at each node equal exceeding the oval nucule of 10—12 spires.—*J. of B. i. t. 7.*—Dark green, 4—8 in. high. Not incrusted.—In brine-pits at Newton, Isle of Wight. A. VIII. E.

\*\* Stem of one central and many cortical tubes (cells).

† *St. with one row of cortical cells to each branchlet. Dioecious.*

3. *C. crinita* (Wallr.); st. slender coarsely striate densely beset with setaceous patent clustered spines, branchlets 8—10 short, stipulodes long, bracts 7—10 whorled slender equal usually exceeding the narrowly oblong nucules of 10—12 spires.—*C. canescens J. of B. xviii. t. 208.*—St. rigid, but little branched, rarely incrusted, pale green. Branchlets stout, 6—8-jointed.—Burdoch Pool near Falmouth. E.

†† *St. with 2 rows of cortical cells to each branchlet.—*  
a. Monœcious.

4. *C. fœtida* (Br.); st. finely striate, primary cortical cells less prominent than secondary and bearing the few small spines, branchlets 6—9 their upper part without cortical cells, stipulodes small blunt, bracts on inner side of branchlet exceeding the ovoid nucule of 12 or 13 spires.—*E. B. 336.* *C. vulgaris L.* in part.—Primary cortical cells collapse as they dry and so place the spines in furrows. Very variable; (*C. longibracteata* Kütz.) branchlets and bracts much longer; (*C. decipiens* Desv.) spines many spreading long deciduous, secondary cortical cells very prominent; (*C. subverticillata* Nordst. MS.) branchlets spreading with long upper naked joints, not incrusted and few spines, dark green (hardly *C. atrovirens* Lowe); (*C. crassicaulis* Schl.) stout, branchlets stout connivent often  $\frac{1}{2}$  naked, bracts short ovate, spines obsolete.—Universal. V. VI. E. S. I.

5. *C. contraria* (A. Br.); st. finely striate, primary cortical cells more prominent than the secondary and bearing the spines or tubercles; branchlets 6—9 coated nearly throughout, stipulodes blunt, bracts 4 (♀) scarcely exceeding the nucules of 12 or 13 spires and a conical contracted crown.—Much like *C. fœtida*. Spines more acute, ultimately on ridges.—Frensham Little Pond, Surrey (1881). *Mr. W. H. Beeby. VII. E.*

6. *C. hispida* (L.); st. stout thickened upwards rough spirally striate, secondary cortical cells larger and more prominent than the primary, many setaceous spreading spines, branchlets 7—9-jointed, stipulodes prominent, bracts on both sides of the branchlet 8—10 exceeding the ovoid nucules of 10 or 11 spires.—*E. B. 463.*—Usually very large, much incrusted,

often 3 or 4 feet long. Sterile branchlets 3—4 inches long. Spines variable in quantity and length.—Ponds and ditches. VI. E. S. I.

7. *C. polyacantha* (Braun); st. stout, *primary cortical cells more prominent* than secondary, many setaceous fascicled spreading spines, 8—10 branchlets of 6—8 joints, stipulodes long slender, bracts 6—10 exceeding the ovoid nucules of 12 or 13 spires.—*J. of B.* xviii. t. 208. *Atl. Fl. Par.* t. 38. B. f. 3.—Very spinous, much incrustated; smaller, more spinous, with shorter branchlets but longer internodes than *C. hispida*.—Ponds and ditches. VI. E. S. I.

8. *C. baltica* (Fr.); st. stout finely striate flexible *primary cortical cells more prominent* than secondary, spines many slender spreading, branchlets 6—9 patent many-jointed, stipulodes slender, bracts whorled long exceeding oblong nucules of 10—15 spires and spreading crown.—Pale green, *not incrustated. Moderately spinous.* Upper part of branches not coated. Bracts very long. Lowest nodes with calcareous bulbils.—Ponds. Kynance Vale, Cornwall. VIII. E.

#### b. Diccious.

9. *C. tomentosa* (L.); st. thickened upwards twisted, *primary cortical cells very prominent*, spines few short stout scattered, branchlets 5—7 rather incurved of 4—6 long joints the upper joints not coated but inflated, *stipulodes usually short and thick*, bracts usually 5 very thick the lateral exceeding the ovoid nucules of 12—14 spires the others falling short of them.—*Hook. Icon. Pl.* 532.—Opaque, greenish white when fresh, usually much incrustated. Spines rather whorled, very short, acute. Upper joints of branchlets remarkably inflated and lengthened in our plant.—Belvidere Lake and near Portumna, Ireland. I.

††† *St. with 3 rows of cortical cells to each branchlet.*

10. *C. aspera* (W.); *diccious*, st. slender flexible with *slender acute spreading spines*, branchlets 6—9 rather incurved or straight, *stipulodes prominent*, bracts whorled nearly equal exceeding the ovoid nucules of 13 spires and *spreading crown*.—*E. B. S.* 2738.—Small, in dense masses. Sometimes with smooth calcareous bulbils on lower nodes. Length and number of spines variable. Joints very long.—Lakes and pools. VII. VIII. E. S. I.

11. *C. con'nivens* (Braun); *diccious*, st. slender brittle with spines, branchlets usually 8 strongly incurved, *stipulodes scarcely visible*, bracts much falling short of nucules scarcely apparent



except with them, nucules small ovoid with 12—14 spires and conical crown.—*J. of B.* xviii. t. 207.—Light green. 12 inches high.—Rare. In south of England. VII. VIII. E.

12. *C. frag'ilis* (Desv.); *monœcious*, st. slender finely striate *no spines*, branchlets 6—9 of 7—9 joints the upper 1—3 joints not coated, lower stylopodes very short, bracts usually 4 usually about equalling the ovoid nucules of 12—14 spires and rather conical crown.—*C. pulchella* *E. B. S.* 2824.—Slender, green, usually not incrustated. Very variable.—β. *C. capillacea* (Thuil.); bracts and stipulodes much longer.—γ. *C. Hedwigii* (Ag.); stronger and larger with very short bracts. *E. B. S.* 2762.—δ. *C. fulcrata* (Gant.); bracts shorter than nucules, stipulodes obsolete. *C. verrucosa* (Itz.); small, 2—4 in. high, often with calcareous bulbils at base.—Slow and stagnant waters. VI.—VIII. E. S. I.

13. *C. fragifera* (Dur.); *diœcious*, st. very finely striate *no spines*, branchlets slender 6—9 of many joints upper joints not coated, stipulodes very small, bracts 3—5 half as long as nucule very short with granule, nucules ovoid of 11—13 spires and short blunt crown.—*J. of B.* xv. t. 192.—Often flexible and exceedingly slender, 6—12 inches long, bright green, not incrustated, bearing large compound bulbils at its base.—West of Cornwall in pools. VII. VIII. E.

---

“Quamquam multas observaverim plantas et sedulo quidem, tamen non confido me semper veritatem invenisse.”—LINK.

---



# ALPHABETICAL INDEX

OF

## THE ORDERS AND GENERA.

\* \* The names in *italics* are synonymous.

- |  |   |   |
|--|---|---|
| <p>Acanthus, 271.<br/>Acer, 70.<br/>Aceraceæ, 70.<br/>Aceras, 316.<br/>Achillea, 193.<br/><i>Achyrophorus</i>, 210.<br/><i>Acinos</i>, 279.<br/>Aconitum, 14.<br/>Acorus, 377.<br/><i>Acrostichum</i>, 455.<br/>Actæa, 14.<br/>Actinocarpus, 358.<br/>Adiantum, 463.<br/>Adonis, 5.<br/>Adoxa, 170.<br/><i>Ægopodium</i>, 157.<br/><i>Æthusa</i>, 161.<br/><i>Agruphis</i>, 367.<br/>Agrimonia, 102.<br/><i>Agrostemma</i>, 54.<br/>Agrostis, 426.<br/>Aira, 429.<br/><i>Aira</i>, 428.<br/><i>Airopsis</i>, 430.<br/>Ajuga, 285.<br/>Alchemilla, 103.<br/>Alisma, 357.<br/><i>Alisma</i>, 358.<br/>Alismaceæ, 356.<br/>Alliaria, 30.<br/>Allium, 364.<br/><i>Allosorus</i>, 453.<br/>Alnus, 334.<br/>Alopecurus, 423.<br/>Alsine, 57.<br/><i>Alsine</i>, 58.<br/>Alsineæ, 48.<br/>Althæa, 66.<br/>Alyssum, 32.<br/>Amaranthaceæ, 297.</p> | <p>Amaranthus, 297.<br/>Amaryllidaceæ, 355.<br/>Ambrosiaceæ, 226.<br/>Amentiferæ, 321.<br/><i>Amesium</i>, 462.<br/><i>Ammophila</i>, 425.<br/>Amygdaleæ, 100.<br/><i>Anacamptis</i>, 345.<br/>Anacharis, 339.<br/>Anagallis, 292.<br/>Anchusa, 248.<br/>Andromeda, 233.<br/><i>Androsæmum</i>, 68.<br/><i>Anemagrostis</i>, 426.<br/>Anemone, 4.<br/>Angelica, 163.<br/>Antennaria, 192.<br/>Anthemis, 194.<br/><i>Anthericum</i>, 363.<br/>Anthoxanthum, 421.<br/><i>Anthriscus</i>, 166.<br/>Anthyllis, 90.<br/>Antirrhinum, 261.<br/><i>Apargia</i>, 211.<br/>Apera, 426.<br/><i>Aphanes</i>, 103.<br/>Apium, 156.<br/>Apocynaceæ, 238.<br/>Aquifoliaceæ, 237.<br/>Aquilegia, 14.<br/>Arabis, 27.<br/><i>Arabis</i>, 29.<br/>Araceæ, 377.<br/>Arbutus, 232.<br/><i>Arbutus</i>, 233.<br/>Archangelica, 163.<br/>Arctium, 202.<br/>Arctostaphylos, 232.<br/>Arenaria, 57.<br/><i>Arenaria</i>, 56.</p> | <p>Aristolochia, 313.<br/>Aristolochiaceæ, 313.<br/>Artmeria, 295<br/>Armoracia, 34.<br/>Arnoseris, 209.<br/>Arrhenatherum, 431.<br/>Artemisia, 196.<br/>Arthrolobium, 97.<br/>Arum, 377.<br/><i>Arundo</i>, 425, 426.<br/>Asarum, 313.<br/>Asparagaceæ, 359.<br/>Asparagus, 359.<br/>Asperugo, 247.<br/>Asperula, 172.<br/><i>Aspidium</i>, 458.<br/>Asplenium, 460.<br/>Aster, 188.<br/>Astragalus, 91.<br/><i>Astragalus</i>, 91.<br/>Astrantia, 155.<br/><i>Athamanta</i>, 162.<br/>Athyrium, 460.<br/>Atriplex, 302.<br/><i>Atriplex</i>, 304.<br/>Atropa, 254.<br/>Avena, 430.<br/><i>Avena</i>, 430.<br/>Azalea, 235.<br/><i>Baothyron</i>, 392.<br/>Ballota, 284.<br/>Balsaminaceæ, 74.<br/>Barbarea, 26.<br/><i>Barkhausia</i>, 214.<br/>Bartsia, 266.<br/><i>Bartsia</i>, 267.<br/><i>Batrachium</i>, 5.<br/>Bellis, 189.<br/>Berberidaceæ, 15.<br/>Berberis, 15.</p> |
|--|---|---|

- Berula*, 159.  
 Beta, 301.  
*Betonica*, 283.  
*Betula*, 333.  
*Bidens*, 201.  
*Blechnum*, 463.  
*Blysmus*, 393.  
 Boraginaceæ, 246.  
*Borago*, 248.  
*Botrychium*, 464.  
*Brachypodium*, 444.  
*Brassica*, 30.  
*Brea*, 207.  
*Briza*, 438.  
*Bromus*, 441.  
*Bromus*, 442.  
*Bryonia*, 138.  
*Bucetum*, 440.  
*Buffonia*, 54.  
*Bunium*, 158.  
*Bupleurum*, 159.  
*Butomæ*, 357.  
*Butomus*, 358.  
*Buxus*, 314.  
*Cakile*, 38.  
*Calamagrostis*, 426.  
*Calamintha*, 278.  
 Callitrichaceæ, 318.  
*Callithriche*, 318.  
*Calluna*, 233.  
*Caltha*, 13.  
*Calycifloræ*, 77.  
*Calystegia*, 244.  
*Camelina*, 34.  
*Campanula*, 228.  
*Campanula*, 230.  
 Campanulaceæ, 227.  
 Cannabinaceæ, 320.  
 Caprifoliaceæ, 170.  
*Capsella*, 37.  
*Cardamine*, 28.  
*Carduus*, 206.  
*Carduus*, 208.  
*Carex*, 395.  
*Carlina*, 201.  
*Carpinus*, 335.  
*Carum*, 158.  
 Caryophyllaceæ, 47.  
*Castanea*, 334.  
*Catabrosa*, 438.  
*Caucalis*, 165.  
 Celastraceæ, 77.  
*Centaurea*, 203.  
*Centranthus*, 177.  
*Centunculus*, 293.  
*Cephalanthera*, 351.  
*Cerastium*, 60.  
*Cerastium*, 60.  
*Cerasus*, 100.  
 Ceratophyllaceæ, 317.  
*Ceratophyllum*, 317.  
*Ceterach*, 462.  
 Chærophyllum, 166.  
*Chamagrostis*, 421.  
*Chara*, 470.  
 Characeæ, 468.  
*Cheiranthus*, 25.  
*Chelidonium*, 19.  
 Chenopodiaceæ, 297.  
*Chenopodium*, 299.  
*Cherleria*, 57.  
*Chlora*, 240.  
*Chrysanthemum*, 196.  
*Chrysocoma*, 189.  
*Chrysosplenium*, 149.  
*Cicendia*, 242.  
*Cichorium*, 209.  
*Cicuta*, 156.  
*Cineraria*, 200.  
*Circæa*, 137.  
*Cirsium*, 206.  
 Cistaceæ, 40.  
*Cistus*, 40.  
*Cladium*, 389.  
*Claytonia*, 139.  
*Clematis*, 3.  
*Clinopodium*, 279.  
*Cnicus*, 206.  
*Cochlearia*, 33.  
*Colchicum*, 368.  
*Comarum*, 105.  
 Compositæ, 180.  
 Coniferæ, 335.  
*Conium*, 167.  
*Convallaria*, 360.  
*Convallaria*, 360.  
 Convolvulaceæ, 244.  
*Convolvulus*, 244.  
*Conyza*, 190.  
*Corallorhiza*, 351.  
*Coriandrum*, 168.  
 Cornaceæ, 168.  
*Cornus*, 169.  
 Corollifloræ, 169.  
*Corrigiola*, 140.  
*Corydalis*, 19.  
*Corylus*, 335.  
*Corynephorus*, 428.  
*Cotoneaster*, 130.  
*Cotyledon*, 144.  
*Crambe*, 38.  
 Crassulaceæ, 141.  
*Cratægus*, 130.  
*Crepis*, 214.  
*Crithmum*, 163.  
*Crocus*, 354.  
 Cruciferæ, 21.  
 Cryptogameæ, 448.  
*Cryptogramme*, 453.  
*Cucubalus*, 50.  
 Cucurbitaceæ, 138.  
*Cuscuta*, 245.  
*Cyclamen*, 291.  
*Cynodon*, 420.  
*Cynoglossum*, 248.  
*Cynosurus*, 438.  
 Cyperaceæ, 387.  
*Cyperus*, 388.  
*Cypripedium*, 352.  
*Cystea*, 459.  
*Cystopteris*, 459.  
*Dabeocia*, 235.  
*Digitalis*, 438.  
*Daphne*, 312.  
*Datura*, 254.  
*Daucus*, 164.  
*Delphinium*, 14.  
*Dentaria*, 28.  
*Deschampsia*, 429.  
*Dianthus*, 50.  
 Dicotyledones, 1.  
 Dictyogenæ, 337.  
*Digitalis*, 261.  
*Digitaria*, 419.  
 Dioscoreaceæ, 338.  
*Diotis*, 196.  
*Diplostaxis*, 32.  
 Dipsacaceæ, 179.  
*Dipsacus*, 179.  
*Doronicum*, 197.

- Draba*, 32.  
*Drosera*, 44.  
*Droseraceæ*, 44.  
*Dryas*, 125.  
*Echinochloa*, 420.  
*Echinophora*, 167.  
*Echium*, 249.  
*Elæagnaceæ*, 311.  
*Elatinaceæ*, 47.  
*Elatine*, 47.  
*Eleocharis*, 390.  
*Elodea*, 339.  
*Elodes*, 70.  
*Elymus*, 446.  
*Elymus*, 446.  
*Empetraceæ*, 313.  
*Empetrum*, 313.  
*Endymion*, 367.  
*Epilobium*, 133.  
*Epimedium*, 15.  
*Epipactis*, 349.  
*Epipogon*, 351.  
*Equisetaceæ*, 448.  
*Equisetum*, 448.  
*Eranthis*, 13.  
*Erica*, 233.  
*Ericaceæ*, 230.  
*Erigeron*, 188.  
*Erinus*, 261.  
*Eriocaulaceæ*, 375.  
*Eriocaulon*, 375.  
*Eriophorum*, 394.  
*Erodium*, 74.  
*Ervum*, 92.  
*Eryngium*, 156.  
*Erysimum*, 30.  
*Erythraea*, 240.  
*Eufragia*, 266.  
*Euonymus*, 77.  
*Eupatorium*, 187.  
*Euphorbia*, 314.  
*Euphorbiaceæ*, 314.  
*Euphrasia*, 267.  
*Exacum*, 242.  
*Fagopyrum*, 311.  
*Fagus*, 334.  
*Falcatula*, 89.  
*Festuca*, 439.  
*Filago*, 191.  
*Filices*, 451.  
*Floridæ*, 338.  
*Fœniculum*, 162.  
*Fragaria*, 106.  
*Frankenia*, 45.  
*Frankeniaceæ*, 45.  
*Fraxinus*, 238.  
*Fritillaria*, 362.  
*Fumaria*, 20.  
*Fumariaceæ*, 19.  
*Gagea*, 363.  
*Galanthus*, 356.  
*Galeobdolon*, 281.  
*Galeopsis*, 282.  
*Galinsoga*, 201.  
*Galium*, 173.  
*Gastridium*, 428.  
*Genista*, 82.  
*Gentiana*, 242.  
*Gentianaceæ*, 239.  
*Geraniaceæ*, 71.  
*Geranium*, 71.  
*Geum*, 125.  
*Gladiolus*, 354.  
*Glaucium*, 18.  
*Glaux*, 292.  
*Glechoma*, 280.  
*Glumiferae*, 387.  
*Glyceria*, 435.  
*Glyceria*, 437.  
*Gnaphalium*, 192.  
*Gnaphalium*, 191.  
*Goodyera*, 348.  
*Gramineæ*, 411.  
*Gymnadenia*, 346.  
*Gymnogramme*, 462.  
*Gymnospermæ*, 335.  
*Gypsocallis*, 234.  
*Habenaria*, 346.  
*Haloragaceæ*, 137.  
*Haloscias*, 162.  
*Hedera*, 168.  
*Hederaceæ*, 168.  
*Helianthemum*, 40.  
*Helleborus*, 13.  
*Helminthia*, 212.  
*Helosciadium*, 156.  
*Heracleum*, 164.  
*Herminium*, 348.  
*Herniaria*, 140.  
*Hesperis*, 29.  
*Hieracium*, 216.  
*Hieracium*, 216.  
*Hierochloe*, 422.  
*Himantoglossum*, 345.  
*Hippocrepis*, 97.  
*Hippophaë*, 311.  
*Hippuris*, 138.  
*Holcus*, 428.  
*Holoschœnus*, 393.  
*Holosteum*, 58.  
*Honkeneja*, 56.  
*Hordeum*, 446.  
*Hottonia*, 289.  
*Humulus*, 320.  
*Hutchinsia*, 35.  
*Hydrocharidaceæ*, 339.  
*Hydrocharis*, 339.  
*Hydrocotyle*, 155.  
*Hymenophyllum*, 464.  
*Hyoscyamus*, 254.  
*Hypericaceæ*, 67.  
*Hypericum*, 68.  
*Hypochæris*, 209.  
*Hypopitys*, 237.  
*Iberis*, 36.  
*Ilex*, 238.  
*Illecebrum*, 140.  
*Impatiens*, 75.  
*Inula*, 189.  
*Inula*, 190.  
*Iridaceæ*, 352.  
*Iris*, 353.  
*Isatis*, 38.  
*Isnardia*, 136.  
*Isoetes*, 466.  
*Isoplepis*, 393.  
*Jasione*, 228.  
*Juncaceæ*, 368.  
*Juncagineæ*, 357.  
*Juncus*, 369.  
*Juncus*, 369.  
*Juniperus*, 337.  
*Knappia*, 421.  
*Knautia*, 180.  
*Kobresia*, 395.  
*Kœleria*, 431.  
*Labiatae*, 271.  
*Lactuca*, 212.  
*Lagurus*, 427.  
*Lamium*, 280.  
*Lapsana*, 209.  
*Lastrea*, 456.  
*Lathraea*, 257.  
*Lathyrus*, 95.

- Lavatera*, 66.  
*Leersia*, 425.  
 Leguminosæ, 78.  
*Lemna*, 378.  
 Lemnaceæ, 378.  
 Lentibulariaceæ, 286.  
*Leontodon*, 210.  
*Leonurus*, 282.  
*Lepidium*, 36.  
*Lepidotis*, 467.  
*Lepidotis*, 467.  
*Lepigonum*, 63.  
*Lepturus*, 446.  
*Leucojum*, 356.  
*Ligusticum*, 162.  
*Ligustrum*, 238.  
 Liliaceæ, 361.  
*Lilium*, 362.  
*Limnanthemum*, 243.  
*Limosella*, 264.  
 Linaceæ, 76.  
*Linaria*, 261.  
 Linnæa, 172.  
*Linosyris*, 189.  
*Linum*, 76.  
*Liparis*, 352.  
*Listera*, 349.  
*Lithospermum*, 250.  
*Littorella*, 297.  
*Lloydia*, 363.  
*Lobelia*, 227.  
 Lobeliæ, 227.  
*Loiseleuria*, 235.  
*Lolium*, 447.  
*Lomaria*, 453.  
*Lonicera*, 171.  
 Loranthaceæ, 169.  
*Lotus*, 90.  
*Luzula*, 373.  
*Ludwigia*, 136.  
*Lychnis*, 53.  
*Lychnothamnus*, 470.  
*Lycium*, 254.  
 Lycopodiaceæ, 465.  
*Lycopodium*, 467.  
*Lycopsis*, 249.  
*Lycopus*, 276.  
*Lysimachia*, 291.  
 Lythraceæ, 132.  
*Lythrum*, 132.  
*Maianthemum*, 360.  
*Malachium*, 60.  
*Malaxis*, 351.  
*Malaxis*, 352.  
*Malcolmia*, 29.  
*Malva*, 65.  
 Malvaceæ, 65.  
*Marrubium*, 284.  
 Marsileaceæ, 465.  
*Maruta*, 194.  
*Matricaria*, 195.  
*Matthiola*, 25.  
*Meconopsis*, 18.  
*Medicago*, 83.  
*Melampyrum*, 264.  
 Melanthaceæ, 367.  
*Melica*, 432.  
*Melilotus*, 84.  
*Melissa*, 279.  
*Melittis*, 280.  
*Mentha*, 274.  
*Menyanthes*, 243.  
*Menziesia*, 235.  
*Mercurialis*, 317.  
*Mertensia*, 250.  
*Mespilus*, 130.  
*Meum*, 162.  
*Mibora*, 421.  
*Milium*, 425.  
*Mimulus*, 265.  
*Moehringia*, 57.  
*Moenchia*, 62.  
*Molinia*, 432.  
*Moneses*, 237.  
 Monochlamydeæ, 297.  
 Monocotyledones, 337.  
*Monotropa*, 237.  
*Montia*, 139.  
*Mulgedium*, 214.  
*Muscari*, 367.  
*Myosotis*, 250.  
*Myosurus*, 5.  
*Myrica*, 333.  
*Myriophyllum*, 137.  
*Myrrhis*, 167.  
*Naiadaceæ*, 386.  
*Naias*, 386.  
*Narcissus*, 355.  
*Nardus*, 424.  
*Narthecium*, 368.  
*Nasturtium*, 25.  
*Nasturtium*, 34.  
*Neotinea*, 346.  
*Neottia*, 349.  
*Nepeta*, 280.  
*Nephrodium*, 458.  
*Nitella*, 468.  
*Nuphar*, 16.  
*Nymphæa*, 16.  
 Nymphæaceæ, 15.  
*Obione*, 304.  
*Odontites*, 267.  
*Oenanthe*, 160.  
*Oenothera*, 136.  
 Oleaceæ, 238.  
 Onagraceæ, 133.  
*Onobrychis*, 97.  
*Ononis*, 82.  
*Onopordum*, 205.  
 Ophioglossaceæ, 464.  
*Ophioglossum*, 465.  
*Ophrys*, 347.  
*Oporinia*, 211.  
 Orchidaceæ, 340.  
*Orchis*, 343.  
*Orchis*, 347.  
*Origanum*, 277.  
*Ornithogalum*, 363.  
*Ornithogalum*, 363.  
*Ornithopus*, 97.  
 Orobanchaceæ, 254.  
*Orobanche*, 255.  
*Orobus*, 96.  
 Orontææ, 377.  
*Osmunda*, 464.  
 Osmundaceæ, 464.  
 Oxalidaceæ, 75.  
*Oxalis*, 75.  
*Oxycoccus*, 236.  
*Oxyria*, 307.  
*Oxytropis*, 91.  
*Pæonia*, 15.  
*Panicum*, 420.  
*Papaver*, 17.  
 Papaveraceæ, 16.  
*Parietaria*, 319.  
*Paris*, 338.  
*Parnassia*, 149.  
 Paronychiaceæ, 139.  
*Pastinaca*, 163.  
*Pedicularis*, 265.  
*Peplis*, 132.

- Peristylus*, 346.  
 Petasites, 187.  
 Petroselinum, 157.  
 Peucedanum, 163.  
 Phalaris, 421.  
*Phelipæa*, 257.  
*Phellandrium*, 161.  
 Phleum, 422.  
 Phragmites, 425.  
*Phyllitis*, 462.  
 Phyllodoce, 235.  
 Physospermum, 167.  
 Phyteuma, 228.  
 Picris, 211.  
 Pilularia, 465.  
 Pimpinella, 158.  
 Pinguicula, 287.  
 Pinus, 337.  
*Pisum*, 96.  
 Plananthus, 467.  
 Plantaginaceæ, 295.  
 Plantago, 295.  
*Platanthera*, 347.  
 Plumbaginaceæ, 293.  
 Poa, 432.  
 Polemoniaceæ, 243.  
 Polemonium, 243.  
 Polycarpon, 63.  
 Polygala, 46.  
 Polygalaceæ, 45.  
 Polygonaceæ, 304.  
 Polygonatum, 360.  
 Polygonum, 308.  
*Polygonum*, 311.  
 Polypodiaceæ, 451.  
 Polypodium, 454.  
 Polypogon, 427.  
 Polystichum, 458.  
 Pomeæ, 130.  
 Populus, 332.  
*Porrum*, 364.  
 Portulacæ, 138.  
 Potamogeton, 379.  
 Potamogetonaceæ, 379.  
 Potentilla, 104.  
 Poterium, 102.  
*Prenanthes*, 313.  
 Primula, 290.  
 Primulaceæ, 288.  
 Prunella, 280.  
 Prunus, 100.  
 Psamma, 425.  
 Pteris, 463.  
 Pulicaria, 190.  
 Pulmonaria, 249.  
*Pyrethrum*, 195.  
 Pyrola, 236.  
*Pyrola*, 237.  
 Pyrus, 130.  
 Quercus, 335.  
 Radiola, 77.  
 Ranunculaceæ, 1.  
 Ranunculus, 5.  
 Raphanus, 38.  
 Reseda, 39.  
 Resedaceæ, 39.  
 Rhamnaceæ, 78.  
 Rhamnus, 78.  
 Rhinanthus, 266.  
*Rhodiola*, 141.  
 Rhynchospora, 389.  
 Ribes, 144.  
 Ribesiaceæ, 144.  
*Robertsonia*, 145.  
 Roemeria, 18.  
 Romulea, 354.  
 Rosa, 126.  
 Rosaceæ, 97.  
 Roseæ, 101.  
*Rottboellia*, 446.  
 Rubia, 176.  
 Rubiaceæ, 172.  
 Rubus, 106.  
 Rumex, 304.  
 Ruppia, 385.  
 Ruscus, 360.  
 Sagina, 54.  
 Sagittaria, 358.  
 Salicornia, 302.  
 Salix, 323.  
 Salsola, 299.  
 Salvia, 276.  
 Sambucus, 172.  
 Samolus, 293.  
 Sanguisorba, 102.  
 Sanicula, 155.  
 Santalaceæ, 312.  
 Saponaria, 51.  
 Sarothamnus, 82.  
 Saussurea, 201.  
 Saxifraga, 145.  
 Saxifragaceæ, 145.  
 Scabiosa, 180.  
 Scandix, 166.  
*Schedonorus*, 441.  
 Scheuchzeria, 358.  
*Schoberia*, 299.  
 Schœnus, 389.  
 Scilla, 364.  
*Scilla*, 367.  
 Scirpus, 390.  
*Scirpus*, 393.  
 Scleranthæ, 64.  
 Scleranthus, 64.  
 Sclerochloa, 436.  
 Scolopendrium, 462.  
 Scrophularia, 263.  
 Scrophulariaceæ, 258.  
 Scutellaria, 279.  
 Sedum, 141.  
 Selaginella, 468.  
*Selinum*, 163.  
 Sempervivum, 143.  
 Senebiera, 37.  
 Senecio, 198.  
 Serrafalcus, 442.  
 Serratula, 203.  
 Seseli, 162.  
 Sesleria, 424.  
 Setaria, 220.  
 Sherardia, 172.  
 Sibbaldia, 104.  
 Sibthorpia, 267.  
 Silaus, 162.  
 Silene, 51.  
 Sileneæ, 50.  
 Siler, 164.  
 Silybum, 208.  
 Simethis, 367.  
 Sinapis, 31.  
*Sinapis*, 31.  
 Sison, 157.  
*Sison*, 157.  
 Sisymbrium, 29.  
*Sisymbrium*, 30.  
 Sisyrinchium, 353.  
 Sium, 159.  
 Smilacina, 360.  
 Smyrnum, 167.  
 Solanaceæ, 252.  
 Solanum, 253.  
 Solidago, 189.

- Sonchus*, 213.  
*Sonchus*, 214.  
*Sorbus*, 131.  
*Sparganium*, 376.  
*Spartina*, 420.  
*Spartium*, 82.  
*Spatularia*, 146.  
*Specularia*, 230.  
*Spergella*, 56.  
*Spergula*, 64.  
*Spergula*, 56.  
*Spergularia*, 63.  
*Spiræa*, 101.  
*Spiranthes*, 348.  
*Stachys*, 283.  
*Statice*, 293.  
*Stellaria*, 58.  
*Stipa*, 425.  
*Stratiotes*, 339.  
*Sturmia*, 352.  
*Sturmia*, 421.  
*Suæda*, 299.  
*Subularia*, 37.  
*Symphytum*, 249.  
*Tamariscaceæ*, 132.  
*Tamarix*, 133.  
*Tamus*, 338.  
*Tanacetum*, 197.  
*Taraxacum*, 213.  
*Taxus*, 336.  
*Teesdalia*, 36.  
*Teucrium*, 285.  
*Thalamifloræ*, 1.  
*Thalictrum*, 3.  
*Thesium*, 312.  
*Thlaspi*, 35.  
*Thlaspi*, 37.  
*Thrinicia*, 210.  
*Thymelaceæ*, 312.  
*Thymus*, 277.  
*Thymus*, 278.  
*Tilia*, 67.  
*Tiliaceæ*, 66.  
*Tillæa*, 141.  
*Tofieldia*, 368.  
*Tolypella*, 470.  
*Tordylium*, 164.  
*Torilis*, 165.  
*Tormentilla*, 105.  
*Tragopogon*, 210.  
*Trichomanes*, 463.  
*Trichonema*, 354.  
*Trientalis*, 292.  
*Trifolium*, 85.  
*Triglochin*, 359.  
*Trigonella*, 89.  
*Trilliaceæ*, 337.  
*Trinia*, 157.  
*Triodia*, 431.  
*Trisetum*, 430.  
*Triticum*, 444.  
*Triticum*, 437.  
*Trollius*, 13.  
*Tulipa*, 362.  
*Turritis*, 27.  
*Tussilago*, 188.  
*Tussilago*, 187.  
*Typha*, 376.  
*Typhaceæ*, 375.  
*Ulex*, 81.  
*Ulmaceæ*, 320.  
*Ulmus*, 321.  
*Umbelliferæ*, 149.  
*Urtica*, 320.  
*Urticaceæ*, 319.  
*Utricularia*, 287.  
*Vaccinium*, 235.  
*Valeriana*, 177.  
*Valeriana*, 177.  
*Valerianaceæ*, 176.  
*Valerianella*, 178.  
*Vella*, 34.  
*Verbascum*, 259.  
*Verbena*, 286.  
*Verbenaceæ*, 286.  
*Veronica*, 268.  
*Viburnum*, 171.  
*Vicia*, 92.  
*Villarsia*, 243.  
*Vinca*, 239.  
*Viola*, 41.  
*Violaceæ*, 41.  
*Viscum*, 170.  
*Vulpia*, 439.  
*Wahlenbergia*, 230.  
*Wolffia*, 379.  
*Woodsia*, 455.  
*Xanthium*, 226.  
*Zannichellia*, 385.  
*Zostera*, 386.



# INDEX

TO THE  
POPULAR ENGLISH NAMES.

---

- Abele, 333.  
 Aconite, 13.  
 Adder's-tongue, 465.  
 Agrimony, 102.  
 Alder, 334.  
 Alexanders, 167.  
 Alkanet, 248.  
 Allseed, 77.  
 Alsike Clover, 88.  
 Anemone, 4.  
 Archangel, 281.  
 Arrow-grass, 359.  
 Arrow-head, 358.  
 Asarabacca, 313.  
 Ash, 238.  
 Aspen, 333.  
 Asphodel, 368.  
 Avens, 125.  
 Awl-wort, 37.  
 Bald-money, 162.  
 Balm, 279.  
 Balm, bastard, 280.  
 Balsam, 75.  
 Bane-berry, 14.  
 Barberry, 15.  
 Barley, 446.  
 Barren Strawberry,  
     105.  
 Barrenwort, 15.  
 Basil, 279.  
 Bastard Pimpernel,  
     293.  
 Bear's-foot, 13.  
 Bed-straw, 173.  
 Beech, 334.  
 Beet, 301.  
 Bell flower, 228, 229.  
 Bent, 425, 426.  
 Betony, 283.  
 Bilberry, 235.  
 Bindweed, 244.  
 Birch, 333.  
 Bird-Cherry, 100.  
 Bird's-foot, 97.  
 Bird's-foot Trefoil,  
     90.  
 Bird's-nest, 349.  
 Bird's-nest, yellow,  
     237.  
 Birthwort, 313.  
 Bitter-Cress, 28.  
 Bitter-sweet, 253.  
 Black Alder, 78.  
 Black Bindweed, 310.  
 Black Bryony, 338.  
 Black grass, 424.  
 Black Saltwort, 292.  
 Blackthorn, 100.  
 Bladder Campion, 52.  
 Bladder-fern, 459.  
 Bladderwort, 287.  
 Blinks, 139.  
 Blue-bell, 367.  
 Bluebottle, 204.  
 Bog-Asphodel, 368.  
 Bog-rush, 389.  
 Borage, 248.  
 Box, 314.  
 Bracken, 463.  
 Brakes, 463.  
 Bramble, 106.  
 Brandy-bottle, 15.  
 Brooklime, 268.  
 Brook-weed, 293.  
 Broom, 82.  
 Broom-rape, 255.  
 Bryony, black, 338.  
 Bryony, red, 138.  
 Buckbean, 243.  
 Buckthorn, 78.  
 Buck-wheat, 311.  
 Bugle, 285.  
 Bugloss, 249.  
 Bugloss, Viper's, 249.  
 Bullace, 100.  
 Bulrush, 392.  
 Burdock, 202.  
 Burnet, 102.  
 Burnet-Saxifrage, 158.  
 Bur-reed, 376.  
 Butcher's Broom, 360.  
 Butterbur, 187.  
 Butterwort, 287.  
 Cabbage, 30.  
 Calamint, 278.  
 Campion, 53, 54.  
 Canary-grass, 421.  
 Candytuft, 36.  
 Caper-Spurge, 317.  
 Caraway, 158.  
 Carrot, 164.  
 Catchfly, 51.  
 Cat-Mint, 280.  
 Cat's-ear, 209.  
 Cat's-foot, 193.  
 Cat's-tail grass, 422.  
 Celandine, 19.  
 Celery, 156.  
 Centaury, 240.  
 Chamomile, 194, 195,  
     196.  
 Charlock, 30.  
 Charlock, jointed, 37.  
 Cherry, 100, 101.  
 Chervil, 166.  
 Chestnut, 334.  
 Chickweed, 60.  
 Chickweed, great, 60.  
 Chicory, 209.  
 Chives, 366.


- Cicely, 167.  
 Cinque-foil, 104, 105.  
 Clary, 277.  
 Cleavers, 174.  
 Cloudberry, 124.  
 Clover, 85, 88.  
 Club-moss, 467.  
 Club-rush, 390.  
 Cock's-foot grass, 438.  
 Cock's-head, 97.  
 Coltsfoot, 188.  
 Columbine, 14.  
 Comfrey, 249.  
 Coralwort, 28.  
 Coriander, 168.  
 Corn-Cockle, 54.  
 Corn-Marigold, 196.  
 Corn Rose, 17.  
 Corn-salad, 178.  
 Cotton-grass, 394.  
 Cotton Thistle, 205.  
 Cotton-weed, 196.  
 Couch-grass, 445.  
 Cowbane, 156.  
 Cowberry, 236.  
 Cow-parsnep, 164.  
 Cowslip, 290.  
 Cow-wheat, 264, 265.  
 Crab-tree, 130.  
 Cranberry, 236.  
 Crane's bill, 71.  
 Crested Hair-grass, 431.  
 Crosswort, 173.  
 Crowberry, 313.  
 Crowfoot, 5, 10, 12.  
 Cuckoo-pint, 377.  
 Cudweed, 191.  
 Currant, 144.  
 Cushion-pink, 53.  
 Cut-grass, 425.  
 Daffodil, 355.  
 Daisy, 189.  
 Dames' Violet, 29.  
 Dandelion, 213.  
 Danewort, 171.  
 Darnel, 447.  
 Dead-nettle, 280.  
 Deadly Nightshade, 254.  
 Devil's-bit, 180.  
 Dewberry, 123.  
 Dittander, 37.  
 Dock, 304.  
 Dodder, 245.  
 Dog's-tail grass, 438.  
 Dog-wood, 169.  
 Dove's-foot, 73.  
 Dropwort, 102.  
 Duckweed, 378.  
 Dutch Clover, 88.  
 Dwale, 254.  
 Dyer's-weed, 82.  
 Elder, 171.  
 Elecampane, 190.  
 Elm, 321.  
 Enchanter's Night-shade, 137.  
 Eryngo, 156.  
 Evening Primrose, 136.  
 Eye-bright, 266, 267.  
 Fat Hen, 300.  
 Feather-grass, 425.  
 Felwort, 236.  
 Fennel, 162.  
 Fern, Bladder, 459.  
 Fern, Holly, 458.  
 Fern, Lady, 460.  
 Fern, Male, 457.  
 Fern, Marsh, 456.  
 Fern, Mountain, 456.  
 Fescue-grass, 439.  
 Feverfew, 195.  
 Figwort, 263.  
 Filmy Fern, 464.  
 Fiorin-grass, 427.  
 Fir, 337.  
 Flag, 353.  
 Flax, 76.  
 Flax-seed, 77.  
 Flea-bane, 188.  
 Flixweed, 29.  
 Flote-grass, 436.  
 Flowering-fern, 464.  
 Flowering-rush, 358.  
 Fluellin, 262.  
 Fool's Parsley, 161.  
 Forget-me-not, 250.  
 Foxglove, 261.  
 Fox-tail grass, 423.  
 Fritillary, 362.  
 Frog-bit, 339.  
 Fumitory, 19, 20.  
 Furze, 81.  
 Gale, 333.  
 Galingale, 388.  
 Garlic, 364.  
 Gean, 101.  
 Gentian, 242.  
 Gipsy-wort, 276.  
 Gladdon, 353.  
 Glasswort, 302.  
 Globe-flower, 13.  
 Goat's-beard, 211.  
 Gold-of-pleasure, 34.  
 Golden-rod, 189.  
 Golden Samphire, 190.  
 Golden Saxifrage, 149.  
 Goldilocks, 11, 189.  
 Gooseberry, 144.  
 Goose-foot, 299.  
 Goose-grass, 174.  
 Gorse, 81.  
 Gout-weed, 157.  
 Grape-Hyacinth, 367.  
 Grass-wrack, 386.  
 Gromwell, 250.  
 Ground-Ivy, 280.  
 Ground-Pine, 286.  
 Groundsel, 198.  
 Guelder-rose, 171.  
 Hairbell, 229.  
 Hair-grass, 429.  
 Hammersedge, 410.  
 Hard fern, 463.  
 Hare's-ear, 159.  
 Hare's-foot, 86.  
 Hare's-tail grass, 427.  
 Hart's-tongue, 462.  
 Hartwort, 164.  
 Hautboy, 106.  
 Hawkbit, 210.  
 Hawk's-beard, 214.  
 Hawkweed, 216.  
 Hawthorn, 130.  
 Hazel, 335.  
 Heartsease, 44.  
 Heath, 233, 235.  
 Heath-grass, 431.  
 Hedge Mustard, 29.  
 Hedge Parsley, 165.  
 Hellebore, 13.

- Helleborine, 349.  
 Hemlock, 167.  
 Hemp-Agrimony, 187.  
 Hemp-nettle, 282.  
 Henbane, 254.  
 Henbit, 281.  
 Hen's-foot, 165.  
 Herb Paris, 338.  
 High-taper, 260.  
 Holly, 238.  
 Holly Fern, 458.  
 Holy grass, 422.  
 Honewort, 157.  
 Honeysuckle, 172.  
 Hop, 320.  
 Horehound, 284.  
 Horehound white, 284.  
 Hornbeam, 335.  
 Horned Pondweed, 385.  
 Horned Poppy, 18.  
 Hornwort, 317.  
 Horsebane, 161.  
 Horse-radish, 34.  
 Horseshoe Vetch, 97.  
 Horsetail, 448.  
 Hound's-tongue, 248.  
 Houseleek, 143.  
 Hyacinth, Grape, 367.  
 Ivy, 168.  
 Jack-by-the-Hedge, 30.  
 Jacob's Ladder, 243.  
 Juniper, 337.  
 Knapweed, 203, 204.  
 Knapwell, 64.  
 Knot-grass, 310.  
 Lady Fern, 460.  
 Lady's Finger, 90.  
 Lady's Mantle, 103.  
 Lady's Slipper, 352.  
 Lady's Smock, 28.  
 Lady's Tresses, 348.  
 Lamb's Lettuce, 178.  
 Lamb's Tongue, 296.  
 Larkspur, 14.  
 Leopard'sbane, 197.  
 Lettuce, 212.  
 Lily, 362.  
 Lily of the Valley, 360.  
 Lime-tree, 67.  
 Ling, 233.  
 Liquorice, 92.  
 Live-long, 142.  
 Lizard Orchis, 345.  
 London Pride, 145.  
 London Rocket, 29.  
 Loosestrife, 291.  
 Loosestrife, purple, 132.  
 Lop-grass, 443.  
 Louse-wort, 265.  
 Loveage, 162.  
 Lucerne, 83.  
 Lungwort, 249.  
 Madder, 172, 176.  
 Madwort, 247.  
 Maiden-hair, 463.  
 Male Fern, 457.  
 Mallow, 65.  
 Maple, 70.  
 Mare's-tail, 138.  
 Marjoram, 277.  
 Marram, 425.  
 Marsh-Cinquefoil, 105.  
 Marsh Fern, 456.  
 Marsh-Mallow, 66.  
 Marsh-Marigold, 13.  
 Masterwort, 163.  
 Matfellow, 204.  
 Mat-grass, 424.  
 Meadow-grass, 432.  
 Meadow-Rue, 3, 4.  
 Meadow-Saffron, 368.  
 Meadow-sweet, 101.  
 Medick, 83, 84.  
 Medlar, 130.  
 Melic, 432.  
 Melilot, 84.  
 Mercury, 317.  
 Mezereon, 312.  
 Mignonette, 39.  
 Milk-Thistle, 208.  
 Milk-vetch, 91.  
 Milkwort, 46.  
 Millefoil, 194.  
 Millet-grass, 425.  
 Mint, 274.  
 Mistletoe, 170.  
 Moneywort, 292.  
 Monk's-hood, 14.  
 Monk's-Rhubarb, 307.  
 Moonwort, 464.  
 Moor-grass, 424.  
 Moschatel, 170.  
 Motherwort, 282.  
 Mountain-Ash, 131.  
 Mountain Fern, 456.  
 Mountain-Sorrel, 307.  
 Mouse-ear, 60.  
 Mousetail, 5.  
 Mudwort, 264.  
 Mugwort, 197.  
 Mullein, 259.  
 Musk-Mallow, 65.  
 Mustard, 31.  
 Navelwort, 144.  
 Needle Whin, 82.  
 Nettle, 320.  
 Nettle, Dead, 280.  
 Nightshade, 253, 254.  
 Nightshade, Enchanter's, 137.  
 Nipplewort, 209.  
 Nit-grass, 428.  
 Oak, 33£.  
 Oat, 430.  
 Oat-grass, 431.  
 Orache, 302.  
 Orchis, 343, 345, 346, 347, 348.  
 Orpine, 142.  
 Osier, 324.  
 Ox-eye, 196.  
 Oxlip, 290.  
 Ox-tongue, 212.  
 Pæony, 15.  
 Paigle, 290.  
 Pansy, 44.  
 Parsley, 157.  
 Parsley Piert, 103.  
 Parsnep, 163.  
 Pasque-flower, 4.  
 Pea, Sea, 96.  
 Pear-tree, 130.  
 Pearlwort, 54.  
 Pellitory, 319.  
 Penny Cress, 35.  
 Penny-royal, 276.  
 Pennywort, 155.  
 Pepperwort, 36.  
 Periwinkle, 239.  
 Pheasant's-eye, 5.

- Pig-nut, 158.  
 Pilewort, 11.  
 Pillwort, 465.  
 Pimpernel, 292.  
 Pink, 50.  
 Pipewort, 375.  
 Plantain, 295.  
 Ploughman's Spike-  
   nard, 190.  
 Polypody, 454.  
 Pond-weed, 378, 385.  
 Poplar, 332.  
 Poppy, 17, 18.  
 Prickwood, 77.  
 Primrose, 290.  
 Privet, 238.  
 Purslane, Sea, 304.  
 Purslane, Water, 132.  
 Quaking-grass, 438.  
 Quill-wort, 466.  
 Quinancy-wort, 173.  
 Radish, 38.  
 Ragged Robin, 54.  
 Ragwort, 198.  
 Rampons, 230.  
 Ramsons, 366.  
 Red Bryony, 138.  
 Red Rattle, 266.  
 Reed, 425, 426.  
 Reed-grass, 421.  
 Reed-mace, 376.  
 Rest-harrow, 82.  
 Ribwort, 269.  
 Rock-brake, 454.  
 Rock-cress, 27.  
 Rock-rose, 40.  
 Rocket, 26.  
 Rose, 126.  
 Rose-bay, 134.  
 Rose-root, 141.  
 Rowan-tree, 131.  
 Rupturewort, 140.  
 Rush, 369.  
 Rustyback, 462.  
 Rye-grass, 447.  
 Sage, 276.  
 Sainfoin, 97.  
 Sallow, 331.  
 Salsify, 211.  
 Saltwort, 299.  
 Samphire, 163.  
 Sand-spurry, 63.  
 Sandwort, 57.  
 Sanicle, 155.  
 Sauce-alone, 30.  
 Saw-wort, 203.  
 Saxifrage, 145.  
 Scabious, 180.  
 Scorpion-grass, 250.  
 Scotch Fir, 337.  
 Scottish Asphodel, 368.  
 Scottish Lovage, 162.  
 Scurvy-grass, 33.  
 Sea-Blite, 299.  
 Sea-Buckthorn, 311.  
 Sea-heath, 45.  
 Sea-Holly, 156.  
 Sea-Kale, 38.  
 Sea-Lavender, 293.  
 Sea-Pea, 96.  
 Sea-Purslane, 304.  
 Sea-Radish, 39.  
 Sea-Rocket, 38.  
 Sedge, 389.  
 Self-heal, 280.  
 Service-tree, 130, 131.  
 Setterwort, 13.  
 Sheep's Fescue Grass,  
   440.  
 Shepherd's Needle,  
   166.  
 Shepherd's Purse, 37.  
 Shepherd's Rod, 180.  
 Shore-weed, 297.  
 Silver-weed, 104.  
 Skull-cap, 279.  
 Sloe, 100.  
 Smallage, 156.  
 Small-reed, 426.  
 Snakeweed, 308.  
 Snapdragon, 261.  
 Sneezewort, 193.  
 Snowdrop, 356.  
 Snowflake, 356.  
 Soapwort, 51.  
 Soft-grass, 428.  
 Solomon's Seal, 360.  
 Sorrel, 307.  
 Sorrel, Mountain, 307.  
 Sorrel, Wood, 75.  
 Sowbane, 300.  
 Sow-bread, 291.  
 Sowthistle, 213.  
 Spearwort, 10, 11.  
 Speedwell, 268.  
 Spindle-tree, 77.  
 Spleenwort, 460, 461.  
 Spurge, 314.  
 Spurry, 64.  
 Squill, 364.  
 St. John's wort, 68.  
 St. Patrick's Cabbage,  
   145.  
 Star of Bethlehem, 363.  
 Star-thistle, 205.  
 Starwort, 188.  
 Stitchwort, 58, 60.  
 Stock, 25.  
 Stonecrop, 141.  
 Stonewort, 157.  
 Storksbill, 74.  
 Strapwort, 140.  
 Strawberry, 105, 106.  
 Strawberry-tree, 232.  
 Succory, 209.  
 Sulphurwort, 162.  
 Sundew, 44.  
 Sweet Cicely, 167.  
 Sweet Flag, 377.  
 Swine's Cress, 38.  
 Swine's Succory, 209.  
 Sycamore, 71.  
 Tamarisk, 133.  
 Tansy, 197.  
 Tares, 92.  
 Tassel-Pondweed, 385.  
 Teasel, 179.  
 Thistle, 205, 206, 207,  
   208.  
 Thorn-apple, 254.  
 Thorough-wax, 160.  
 Thrift, 295.  
 Thrumwort, 358.  
 Thyme, 277.  
 Timothy-grass, 423.  
 Toad-flax, 261, 263.  
 Toothwort, 257.  
 Traveller's Joy, 3.  
 Tree-mallow, 66.  
 Trefoil, 90.  
 Tulip, 362.  
 Turk's-cap Lily, 362.  
 Tutsan, 68.

- Tway-blade, 349.  
 Valerian, 177.  
 Vernal-grass, 421.  
 Vervain, 286.  
 Vetch, 92.  
 Vetchling, 95.  
 Violet, 41.  
 Viper's Bugloss, 249.  
 Wallflower, 25.  
 Wall-Mustard, 32.  
 Wall-Pepper, 143.  
 Wall-Rue, 462.  
 Wart-Cress, 37.  
 Water-Avens, 126.  
 Water-Cress, 25.  
 Water-Dropwort, 160.  
 Water-Fennel, 6.  
 Water-Hemlock, 156.  
 Water-Lily, 16.  
 Water-Milfoil, 137.  
 Water-Pepper, 309.  
 Water-Plantain, 357.  
 Water-Soldier, 339.  
 Water-Starwort, 318.  
 Water-Thyme, 339.  
 Water-Violet, 289.  
 Waterwort, 47.  
 Way-bread, 297.  
 Wayfaring tree, 171.  
 Weld, 39.  
 Welsh Poppy, 18.  
 Wheat-grass, 444.  
 Whin, 81, 82.  
 White Bottle, 52.  
 Whitlow-grass, 32.  
 Whortleberry, 236.  
 Willow, 323.  
 Willow Herb, 133.  
 Wind-flower, 5.  
 Wind-grass, 426.  
 Winter-green, 236.  
 Woad, 38.  
 Woodbine, 172.  
 Woodruff, 173.  
 Wood-Rush, 373.  
 Wood-Sage, 285.  
 Wood-Sorrel, 75.  
 Worm-seed, 30.  
 Wormwood, 196.  
 Woundwort, 283.  
 Yarrow, 193.  
 Yellow Rattle, 266.  
 Yellow wort, 240.  
 Yew, 336.  
 Yorkshire Fog, 428.

THE END.





## Other Books on Botany published by Mr. Van Voorst.

- ~~~~~
- In one large 8vo Volume. £3 3s.
- BRITISH WILD FLOWERS.** Illustrated by JOHN E. SOWERBY. Described, with an Introduction and a Key to the Natural Orders, by C. PIERPOINT JOHNSON. Containing a coloured figure of each of the 1780 plants comprised in the British Flora.
- A MANUAL OF THE BRITISH MARINE ALGÆ,** containing Generic and Specific Descriptions of all the known British Species of Sea-weeds, with Plates to illustrate all the Genera. By Professor W. H. HARVEY, M.D., M.R.I.A. 8vo, 21s., coloured copies £1 11s. 6d.
- BRITISH POISONOUS PLANTS.** Illustrated by JOHN E. SOWERBY. Described by CHARLES JOHNSON, Botanical Lecturer at Guy's Hospital, and C. P. JOHNSON. Second Edition, containing the principal Poisonous Fungi. Post 8vo, with 32 Coloured Plates, 9s. 6d.
- THE BRITISH FERNS AT ONE VIEW.** By BERTHOLD SEEMANN, Ph.D., F.L.S. Illustrated by WALTER FITCH, F.L.S. An eight-page folded sheet of Coloured Figures, with Magnified Views of the Generic characters, and Descriptions of the Genera. Cloth, 6s.
- FERNS OF THE BRITISH ISLES.** Described and Photographed by SY. C. 8vo, cloth, with 20 Plates, giving Photographs of all the Species, Woodcuts showing the Fructification, and an Illustrated Glossary of Terms. 10s. 6d.
- AN ILLUSTRATED KEY TO THE NATURAL ORDERS OF BRITISH WILD FLOWERS.** By JOHN E. SOWERBY. With 9 Plates, containing 112 Coloured Figures. 8vo, 7s. 6d.
- MANUAL of BRITISH BOTANY;** containing the Flowering Plants and Ferns, arranged according to their Natural Orders. By C. C. BABINGTON, M.A., F.R.S., F.L.S., &c. Eighth Edition, Corrected throughout. 12mo, 10s. 6d., cloth; or copies on thin paper, for the pocket, 12s., roan.
- FLORA OF CAMBRIDGESHIRE;** or, a Catalogue of Plants found in the County of Cambridge. By Professor BABINGTON. 12mo, 7s.
- FLORA OF EASTBOURNE;** being an Introduction to the Flowering Plants, Ferns, &c. of the Cuckmere District, East Sussex. By F. C. S. ROPER, F.L.S. &c. Post 8vo, with a Map, 4s. 6d.
- FLORA OF SURREY :** From the Manuscripts of the late J. D. SALMON, F.L.S., and other Sources. Compiled by J. A. BREWER. 12mo, with 2 Maps, 7s. 6d.
- CONTRIBUTIONS TOWARDS A CYBELE HIBERNICA.** By DAVID MOORE, Ph.D. &c., and A. G. MORE, F.L.S. Post 8vo, 10s. 6d.
- THE BRITISH RUBI :** An attempt to discriminate the Species of *Rubus* known to inhabit the British Isles. By Professor BABINGTON. Fcap. 8vo, 5s.

*Books on Botany (continued).*

---

**PRINCIPLES OF THE ANATOMY AND PHYSIOLOGY OF THE VEGETABLE CELL.** By HUGO VON MOHL. Translated, with the Author's permission, by ARTHUR HENFREY, F.R.S. &c. 8vo, with an Illustrative Plate and numerous Woodcuts, 7s. 6d.

Also by PROFESSOR HENFREY,

**THE VEGETATION OF EUROPE, ITS CONDITIONS AND CAUSES.** Fcap. 8vo, 5s.

**THE RUDIMENTS OF BOTANY.** A familiar Introduction to the Study of Plants. Second Edition, Fcap. 8vo, with Illustrative Woodcuts, 3s. 6d.

**A SET OF SIX COLOURED DIAGRAMS ILLUSTRATIVE OF THE RUDIMENTS OF BOTANY:** for Schools and Lectures. 15s.

---

**RYMER JONES'S GENERAL OUTLINE** of the ORGANIZATION of the ANIMAL KINGDOM, and MANUAL of COMPARATIVE ANATOMY. Fourth Edition. 571 Engravings. £1 11s. 6d.

**ATTFIELD'S CHEMISTRY: GENERAL, MEDICAL, and PHARMACEUTICAL;** including the Chemistry of the British Pharmacopœia. Ninth Edition. Illustrated. 15s.

**FRANKLAND'S LECTURE NOTES** for CHEMICAL STUDENTS. Second Edition. Vol. I. (Inorganic), 4s. Vol. II. (Organic), 5s.

**GUTHRIE'S ELEMENTS** of HEAT and of NON-METALLIC CHEMISTRY. 7s.

**ELIOT and STORER'S MANUAL** of INORGANIC CHEMISTRY; arranged to facilitate the Experimental Demonstration of the Facts and Principles of the Science. 10s. 6d.

**GREVILLE WILLIAMS'S HANDBOOK** of CHEMICAL MANIPULATION. Illustrated. 15s.

**CHURCH'S LABORATORY GUIDE:** A Manual of Practical Chemistry for Colleges and Schools, specially arranged for Agricultural Students. Fourth Edition. 6s. 6d.

**GRIFFITH'S ELEMENTARY TEXT-BOOK** of the MICROSCOPE; Including a Description of the Methods of preparing and Mounting Objects, &c. With 12 Coloured Plates. 7s. 6d.

In 1 vol. of 900 pages, Illustrated by 48 Plates and 812 Woodcuts, £2 12s. 6d.  
**THE MICROGRAPHIC DICTIONARY;** a Guide to the Examination and Investigation of the Structure and Nature of Microscopic Objects. By Dr. GRIFFITH and Professor HENFREY. Third Edition. Edited by J. W. GRIFFITH, M.D. &c., and Professor MARTIN DUNCAN, M.D. Lond., F.R.S., &c., assisted by the Rev. M. J. BERKELEY, M.A., F.L.S., and T. RUPERT JONES, F.R.S., Professor of Geology, Sandhurst, &c.

Fourth Edition. In Monthly parts, 2s. 6d. each. Anticipated to be completed in 21 Monthly parts.

---

JOHN VAN VOORST, 1 PATERNOSTER ROW.









